# Regularity of stable solutions of *p*-Laplace equations through geometric Sobolev type inequalities

Daniele Castorina

Manel Sanchón

#### Abstract

In this paper we prove a Sobolev and a Morrey type inequality involving the mean curvature and the tangential gradient with respect to the level sets of the function that appears in the inequalities. Then, as an application, we establish *a priori* estimates for semi-stable solutions of  $-\Delta_p u = g(u)$  in a smooth bounded domain  $\Omega \subset \mathbb{R}^n$ . In particular, we obtain new  $L^r$  and  $W^{1,r}$  bounds for the extremal solution  $u^*$  when the domain is strictly convex. More precisely, we prove that  $u^* \in L^{\infty}(\Omega)$  if  $n \leq p + 2$  and  $u^* \in L^{\frac{np}{n-p-2}}(\Omega) \cap W_0^{1,p}(\Omega)$  if n > p + 2.

**Keywords.** Geometric inequalities, mean curvature of level sets, Schwarz symmetrization, *p*-Laplace equations, regularity of stable solutions

### **1** Introduction

The aim of this paper is to obtain *a priori* estimates for semi-stable solutions of *p*-Laplace equations. We will accomplish this by proving some geometric type inequalities involving the functionals

$$I_{p,q}(v;\Omega) := \left( \int_{\Omega} \left( \frac{1}{p'} |\nabla_{T,v}| \nabla v|^{p/q} | \right)^q + |H_v|^q |\nabla v|^p \, dx \right)^{1/p}, \quad p,q \ge 1$$
(1.1)

where  $\Omega$  is a smooth bounded domain of  $\mathbb{R}^n$  with  $n \geq 2$  and  $v \in C_0^{\infty}(\overline{\Omega})$ . Here, and in the rest of the paper,  $H_v(x)$  denotes the mean curvature at x of the hypersurface  $\{y \in \Omega :$  $|v(y)| = |v(x)|\}$  (which is smooth at points  $x \in \Omega$  satisfying  $\nabla v(x) \neq 0$ ), and  $\nabla_{T,v}$  is the

D. Castorina: Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain; e-mail: castorina@mat.uab.cat

M. Sanchón: Departament de Matemàtica Aplicada i Anàlisi, Universitat de Barcelona, Gran Via 585, 08007 Barcelona, Spain; e-mail: msanchon@maia.ub.es

Mathematics Subject Classification (2010): Primary 35K57, 35B65 ; Secondary 35J60

tangential gradient along a level set of |v|. We will prove a Morrey's type inequality when n and a Sobolev inequality when <math>n > p + q (see Theorem 1.2 below).

Then, as an application of these inequalities, we establish  $L^r$  and  $W^{1,r}$  a priori estimates for semi-stable solutions of the reaction-diffusion problem

$$\begin{cases}
-\Delta_p u = g(u) & \text{in } \Omega, \\
u > 0 & \text{in } \Omega, \\
u = 0 & \text{on } \partial\Omega.
\end{cases}$$
(1.2)

Here, the diffusion is modeled by the *p*-Laplace operator  $\Delta_p$  (remember that  $\Delta_p u := \operatorname{div}(|\nabla u|^{p-2}\nabla u)$ ) with p > 1, while the reaction term is driven by any positive  $C^1$  non-linearity *g*.

As we will see, these estimates will lead to new  $L^r$  and  $W^{1,r}$  bounds for the extremal solution  $u^*$  of (1.2) when  $g(u) = \lambda f(u)$  and the domain  $\Omega$  is strictly convex. More precisely, we prove that  $u^* \in L^{\infty}(\Omega)$  if  $n \leq p+2$  and  $u^* \in L^{\frac{np}{n-p-2}}(\Omega) \cap W_0^{1,p}(\Omega)$  if n > p+2.

#### **1.1 Geometric Sobolev inequalities**

Before we establish our Sobolev and Morrey type inequalities we will state that the functional  $I_{p,q}$  defined in (1.1) decreases (up to a universal multiplicative constant) by Schwarz symmetrization. Given a Lipschitz continuous function v and its Schwarz symmetrization  $v^*$  it is well known that

$$\int_{B_R} |v^*|^r \, dx = \int_{\Omega} |v|^r \, dx \quad \text{for all } r \in [1, +\infty]$$

and

$$\int_{B_R} |\nabla v^*|^r \, dx \le \int_{\Omega} |\nabla v|^r \, dx \quad \text{for all } r \in [1,\infty).$$

Our first result establishes that  $I_{p,q}(v^*; B_R) \leq CI_{p,q}(v; \Omega)$  for some universal constant C depending only on n, p, and q.

**Theorem 1.1.** Let  $\Omega$  be a smooth bounded domain of  $\mathbb{R}^n$  with  $n \geq 2$  and  $B_R$  the ball centered at the origin and with radius  $R = (|\Omega|/|B_1|)^{1/n}$ . Let  $v \in C_0^{\infty}(\overline{\Omega})$  and  $v^*$  its Schwarz symmetrization. Let  $I_{p,q}$  be the functional defined in (1.1) with  $p, q \geq 1$ . If n > q + 1 then there exists a universal constant C depending only on n, p, and q, such that

$$\left(\int_{B_R} \frac{1}{|x|^q} |\nabla v^*|^p \, dx\right)^{1/p} = I_{p,q}(v^*; B_R) \le C I_{p,q}(v; \Omega). \tag{1.3}$$

Note that the Schwarz symmetrization of v is a radial function, and hence, its level sets are spheres. In particular, the mean curvature  $H_{v^*}(x) = 1/|x|$  and the tangential gradient  $\nabla_{T,v^*} |\nabla v^*|^{p/q} = 0$ . This explains the equality in (1.3).

A related result was proved by Trudinger [18] when q = 1 for the class of mean convex functions (*i.e.*, functions for which the mean curvature of the level sets is nonnegative). More precisely, he proved Theorem 1.1 replacing the functional  $I_{p,q}$  by

$$\tilde{I}_{p,q}(v;\Omega) := \left(\int_{\Omega} |H_v|^q |\nabla v|^p \, dx\right)^{1/p} \tag{1.4}$$

and considering the Schwarz symmetrization of v with respect to the perimeter instead of the classical one like us (see Definition 2.1 below). In order to define this symmetrization (with respect to the perimeter) it is essential to know that the mean curvature  $H_v$  of the level sets of |v| is nonnegative. Then using an Aleksandrov-Fenchel inequality for mean convex hypersurfaces (see [17]) he proved Theorem 1.1 for this class of functions when q = 1.

We prove Theorem 1.1 using two ingredients. The first one is the classical isoperimetric inequality:

$$n|B_1|^{1/n}|D|^{(n-1)/n} \le |\partial D| \tag{1.5}$$

for any smooth bounded domain D of  $\mathbb{R}^n$ . The second one is a geometric Sobolev inequality, due to Michael and Simon [12] and to Allard [1], on compact (n - 1)-hypersurfaces M without boundary which involves the mean curvature H of M: for every  $q \in [1, n - 1)$ , there exists a constant A depending only on n and q such that

$$\left(\int_{M} |\phi|^{q^{\star}} d\sigma\right)^{1/q^{\star}} \le A \left(\int_{M} |\nabla\phi|^{q} + |H\phi|^{q} d\sigma\right)^{1/q}$$
(1.6)

for every  $\phi \in C^{\infty}(M)$ , where  $q^* = (n-1)q/(n-1-q)$  and  $d\sigma$  denotes the area element in M. Using the classical isoperimetric inequality (1.5) and the geometric Sobolev inequality (1.6) with  $M = \{x \in \Omega : |v(x)| = t\}$  and  $\phi = |\nabla v|^{(p-1)/q}$  we will prove Theorem 1.1 with the explicit constant  $C = A^{\frac{q}{p}} |\partial B_1|^{\frac{q}{(n-1)p}}$ , being A the universal constant in (1.6).

From Theorem 1.1 and well known 1-dimensional weighted Sobolev inequalities it is easy to prove Morrey and Sobolev geometric inequalities involving the functional  $I_{p,q}$ . Indeed, by Theorem 1.1 and since Schwarz symmetrization preserves the  $L^r$  norm, it is sufficient to prove the existence of a positive constant  $\overline{C}$  independent of  $v^*$  such that

$$||v^*||_{L^r(B_R)} \le \overline{C}I_{p,q}(v^*; B_R).$$

Using this argument we prove the following geometric inequalities.

**Theorem 1.2.** Let  $\Omega$  be a smooth bounded domain of  $\mathbb{R}^n$  with  $n \ge 2$  and  $v \in C_0^{\infty}(\overline{\Omega})$ . Let  $I_{p,q}$  be the functional defined in (1.1) with  $p, q \ge 1$  and

$$p_q^\star := \frac{np}{n - (p + q)}$$

Assume n > q + 1. The following assertions hold:

(a) If n then

$$\|v\|_{L^{\infty}(\Omega)} \le C_1 |\Omega|^{\frac{p+q-n}{np}} I_{p,q}(v;\Omega)$$

$$(1.7)$$

for some constant  $C_1$  depending only on n, p, and q.

(b) If n > p + q, then

$$\|v\|_{L^{r}(\Omega)} \le C_{2}|\Omega|^{\frac{1}{r} - \frac{1}{p_{q}^{\star}}} I_{p,q}(v;\Omega) \quad \text{for every } 1 \le r \le p_{q}^{\star},$$
 (1.8)

where  $C_2$  is a constant depending only on n, p, q, and r.

(c) If n = p + q, then

$$\int_{\Omega} \exp\left\{\left(\frac{|v|}{C_3 I_{p,q}(v;\Omega)}\right)^{p'}\right\} dx \le \frac{n}{n-1} |\Omega|, \quad \text{where } p' = p/(p-1), \quad (1.9)$$

for some positive constant  $C_3$  depending only on n and p.

Cabré and the second author [6] proved recently Theorem 1.2 under the assumption  $q \ge p$  using a different method (without the use of Schwarz symmetrization). More precisely, they proved the theorem replacing the functional  $I_{p,q}(v;\Omega)$  by the one defined in (1.4),  $\tilde{I}_{p,q}(v;\Omega)$ . Therefore, our geometric inequalities are only new in the range  $1 \le q < p$ .

**Open Problem 1.** Is Theorem 1.2 true for the range  $1 \le q < p$  and replacing the functional  $I_{p,q}(v; \Omega)$  by the one defined in (1.4),  $\tilde{I}_{p,q}(v; \Omega)$ ?

This question has a posive answer for the class of mean convex functions. Trudinger [18] proved this result for this class of functions when q = 1 and can be easily extended for every  $q \ge 1$ . However, to our knowledge, for general functions (without mean convex level sets) it is an open problem.

#### **1.2 Regularity of semi-stable solutions**

The second part of the paper deals with *a priori* estimates for semi-stable solutions of problem (1.2). Remember that a regular solution  $u \in C_0^1(\overline{\Omega})$  of (1.2) is said to be *semi-stable* if the second variation of the associated energy functional at u is nonnegative definite, *i.e.*,

$$\int_{\Omega} |\nabla u|^{p-2} \left\{ |\nabla \phi|^2 + (p-2) \left( \nabla \phi \cdot \frac{\nabla u}{|\nabla u|} \right)^2 \right\} - g'(u)\phi^2 \, dx \ge 0 \tag{1.10}$$

for every  $\phi \in H_0$ , where  $H_0$  denotes the space of admissible functions (see Definition 4.1 below). The class of semi-stable solutions includes local minimizers of the energy functional as well as minimal and extremal solutions of (1.2) when  $g(u) = \lambda f(u)$ .

Using an appropriate test function in (1.10) we prove the following *a priori* estimates for semi-stable solutions. This result extends the ones in [3] and [6] for the Laplacian case (p = 2) due to Cabré and the second author.

**Theorem 1.3.** Let g be any  $C^{\infty}$  function and  $\Omega \subset \mathbb{R}^n$  any smooth bounded domain. Let  $u \in C_0^1(\overline{\Omega})$  be a semi-stable solution of (1.2), i.e., a solution satisfying (1.10). The following assertions hold:

(a) If  $n \le p + 2$  then there exists a constant C depending only on n and p such that

$$\|u\|_{L^{\infty}(\Omega)} \le s + \frac{C}{s^{2/p}} |\Omega|^{\frac{p+2-n}{np}} \left( \int_{\{u \le s\}} |\nabla u|^{p+2} dx \right)^{1/p} \quad \text{for all } s > 0.$$
(1.11)

(b) If n > p + 2 then there exists a constant C depending only on n and p such that

$$\left(\int_{\{u>s\}} \left(|u|-s\right)^{\frac{np}{n-(p+2)}} dx\right)^{\frac{n-(p+2)}{np}} \le \frac{C}{s^{2/p}} \left(\int_{\{u\le s\}} |\nabla u|^{p+2} dx\right)^{1/p}$$
(1.12)

for all s > 0. Moreover, there exists a constant C depending only on n, p, and r such that

$$\int_{\Omega} |\nabla u|^r \, dx \le C \left( |\Omega| + \int_{\Omega} |u|^{\frac{np}{n-(p+2)}} \, dx + \|g(u)\|_{L^1(\Omega)} \right) \tag{1.13}$$

for all  $1 \le r < r_1 := \frac{np^2}{(1+p)n-p-2}$ .

To prove (1.11) and (1.12) we use the semi-stability condition (1.10) with the test function  $\phi = |\nabla u|\eta$  to obtain

$$\int_{\Omega} \left( \frac{4}{p^2} |\nabla_{T,u}| \nabla u|^{p/2} |^2 + \frac{n-1}{p-1} H_u^2 |\nabla u|^p \right) \eta^2 \, dx \le \int_{\Omega} |\nabla u|^p |\nabla \eta|^2 \, dx \tag{1.14}$$

for every Lipschitz function  $\eta$  in  $\overline{\Omega}$  with  $\eta|_{\partial\Omega} = 0$ . Then, taking  $\eta = T_s u = \min\{s, u\}$ , we obtain (1.11) and (1.12) when  $n \neq p+2$  by using the Morrey and Sobolev inequalities established in Theorem 1.2 with q = 2. The critical case n = p + 2 is more involved. In order to get (1.11) in this case, we take another explicit test function  $\eta = \eta(u)$  in (1.14) and use the geometric Sobolev inequality (1.6). The gradient estimate established in (1.13) will follow by using a technique introduced by Bénilan *et al.* [2] to get the regularity of entropy solutions for *p*-Laplace equations with  $L^1$  data (see Proposition 4.2).

The rest of the introduction deals with the regularity of extremal solutions. Let us recall the problem and some known results in this topic. Consider

$$\begin{cases} -\Delta_p u = \lambda f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$
(1.15)<sub>\lambda</sub>

where  $\lambda$  is a positive parameter and f is a  $C^1$  positive increasing function satisfying

$$\lim_{t \to +\infty} \frac{f(t)}{t^{p-1}} = +\infty.$$
 (1.16)

Cabré and the second author [5] proved the existence of an extremal parameter  $\lambda^* \in (0, \infty)$  such that problem  $(1.15)_{\lambda}$  admits a minimal regular solution  $u_{\lambda} \in C_0^1(\overline{\Omega})$  for  $\lambda \in (0, \lambda^*)$  and admits no regular solution for  $\lambda > \lambda^*$ . Moreover, every minimal solution  $u_{\lambda}$  is a semi-stable for  $\lambda \in (0, \lambda^*)$ .

For the Laplacian case (p = 2), the limit of minimal solutions

$$u^{\star} := \lim_{\lambda \uparrow \lambda^{\star}} u_{\lambda}$$

is a weak solution of the extremal problem  $(1.15)_{\lambda^*}$  and it is known as extremal solution. Nedev [13] proved, in the case of convex nonlinearities, that  $u^* \in L^{\infty}(\Omega)$  if  $n \leq 3$  and  $u^* \in L^r(\Omega)$  for all  $1 \leq r < n/(n-4)$  if  $n \geq 4$ . Recently, Cabré [3], Cabré and the second author [6], and Nedev [14] proved, in the case of convex domains and general nonlinearities, that  $u^* \in L^{\infty}(\Omega)$  if  $n \leq 4$  and  $u^* \in L^{\frac{2n}{n-4}}(\Omega) \cap H^1_0(\Omega)$  if  $n \geq 5$ .

For arbitrary p > 1 it is unknown if the limit of minimal solutions  $u^*$  is a (weak or entropy) solution of  $(1.15)_{\lambda^*}$ . In the affirmative case, it is called the *extremal solution of*  $(1.15)_{\lambda^*}$ . However, in [15] it is proved that the limit of minimal solutions  $u^*$  is a weak solution (in the distributional sense) of  $(1.15)_{\lambda^*}$  whenever  $p \ge 2$  and f satisfies the additional condition:

there exists 
$$T \ge 0$$
 such that  $(f(t) - f(0))^{1/(p-1)}$  is convex for all  $t \ge T$ . (1.17)

Moreover,

$$u^* \in L^\infty(\Omega)$$
 if  $n$ 

and

$$u^* \in L^r(\Omega)$$
, for all  $r < \tilde{r}_0 := (p-1)\frac{n}{n - (p+p')}$ , if  $n \ge p + p'$ .

This extends previous results of Nedev [13] for the Laplacian case (p = 2) and convex nonlinearities.

Our next result improves the  $L^q$  estimate in [13, 15] for strictly convex domains. We also prove that  $u^*$  belongs to the energy class  $W_0^{1,p}(\Omega)$  independently of the dimension extending an unpublished result of Nedev [14] for p = 2 to every  $p \ge 2$  (see also [6]).

**Theorem 1.4.** Let f be an increasing positive  $C^1$  function satisfying (1.16). Assume that  $\Omega$  is a smooth strictly convex domain of  $\mathbb{R}^n$ . Let  $u_{\lambda} \in C_0^1(\overline{\Omega})$  be the minimal solution of  $(1.15)_{\lambda}$ . There exists a constant C independent of  $\lambda$  such that:

- (a) If  $n \le p + 2$  then  $||u_{\lambda}||_{L^{\infty}(\Omega)} \le C ||f(u_{\lambda})||_{L^{1}(\Omega)}^{1/(p-1)}$ .
- (b) If n > p + 2 then  $\|u_{\lambda}\|_{L^{\frac{np}{n-p-2}}(\Omega)} \le C \|f(u_{\lambda})\|_{L^{1}(\Omega)}^{1/(p-1)}$ . Moreover  $\|u_{\lambda}\|_{W_{0}^{1,p}(\Omega)} \le C'$ where C' is a constant depending only on  $n, p, \Omega, f$  and  $\|f(u_{\lambda})\|_{L^{1}(\Omega)}$ .

Assume, in addition,  $p \ge 2$  and that (1.17) holds. Then

(i) If  $n \leq p+2$  then  $u^* \in L^{\infty}(\Omega)$ . In particular,  $u^* \in C_0^1(\overline{\Omega})$ .

(*ii*) If 
$$n > p + 2$$
 then  $u^* \in L^{\frac{np}{n-p-2}}(\Omega) \cap W^{1,p}_0(\Omega)$ .

**Remark 1.5.** If  $f(u_{\lambda})$  is bounded in  $L^{1}(\Omega)$  by a constant independent of  $\lambda$ , then parts (a) and (b) will lead automatically to the assertions (i) and (ii) stated in the theorem (without the requirement that  $p \geq 2$  and (1.17) hold true). However, as we said before, the estimate  $f(u^{\star}) \in L^{1}(\Omega)$  is unknown in the general case, *i.e.*, for arbitrary positive and increasing nonlinearities f satisfying (1.16) and arbitrary p > 1.

**Open Problem 2.** Is it true that  $f(u^*) \in L^1(\Omega)$  for arbitrary positive and increasing nonlinearities f satisfying (1.16)?

Under assumptions  $p \ge 2$  and (1.17) it is proved in [15] that  $f(u^*) \in L^r(\Omega)$  for all  $1 \le r < n/(n-p')$  when  $n \ge p'$  and  $f(u^*) \in L^{\infty}(\Omega)$  if n < p'. In particular, one has  $f(u^*) \in L^1(\Omega)$  independently of the dimension n and the parameter p > 1. As a consequence, assertions (i) and (ii) follow immediately from parts (a) and (b) of the theorem.

To prove the  $L^r$  *a priori* estimates stated in part (*a*) and (*b*) we make three steps. First, we use the strict convexity of the domain  $\Omega$  to prove that

$$\{x \in \Omega : \operatorname{dist}(x, \partial \Omega) < \varepsilon\} \subset \{x \in \Omega : u_{\lambda}(x) < s\}$$

for a suitable *s*. This is done using a moving plane procedure for *p*-Laplace equations (see Proposition 3.1 below). Then, we prove that the Morrey and Sobolev type inequalities stated in Theorem 1.2 for smooth functions, also hold for regular solutions of (1.2) when  $1 \le q \le 2$ . Finally, taking a test function  $\eta$  related to dist $(\cdot, \partial \Omega)$  in (1.14) and proceeding as in the proof of Theorem 1.3 we will obtain the  $L^r$  *a priori* estimates established in the theorem.

The energy estimate established in parts (ii) and (b) of Theorem 1.4 follows by extending the arguments of Nedev [14] for the Laplacian case (see also Theorem 2.9 in [6]). First, using a Pohožaev identity we obtain

$$\int_{\Omega} |\nabla u_{\lambda}|^{p} dx \leq \frac{1}{p'} \int_{\partial \Omega} |\nabla u_{\lambda}|^{p} x \cdot \nu d\sigma, \quad \text{for all } p > 1 \text{ and } \lambda \in (0, \lambda^{\star}), \quad (1.18)$$

where  $d\sigma$  denotes the area element in  $\partial\Omega$  and  $\nu$  is the outward unit normal to  $\Omega$ . Then, using the strict convexity of the domain (as in the  $L^r$  estimates) and standard regularity estimates for  $-\Delta_p u = \lambda f(u_\lambda(x))$  in a neighborhood of the boundary, we are able to control the right hand side of (1.18) by a constant whose dependence on  $\lambda$  is given by a function of  $||f(u_\lambda)||_{L^1(\Omega)}$ . **Remark 1.6.** Let us compare our regularity results with the sharp ones proved by Cabré, Capella, and the second author in [4] when  $\Omega$  is the unit ball  $B_1$  of  $\mathbb{R}^n$ . In the radial case, the extremal solution  $u^*$  of  $(1.15)_{\lambda^*}$  is bounded if the dimension n . Moreover, $if <math>n \ge p + \frac{4p}{p-1}$  then  $u^* \in W_0^{1,r}(B_1)$  for all  $1 \le r < \overline{r}_1$ , where

$$\bar{r}_1 := \frac{np}{n - 2\sqrt{\frac{n-1}{p-1}} - 2}.$$

In particular,  $u^* \in L^r(B_1)$  for all  $1 \le r < \overline{r}_0$ , where

$$\bar{r}_0 := rac{np}{n - 2\sqrt{rac{n-1}{p-1}} - p - 2}.$$

It can be shown that these regularity results are sharp by taking the exponential and power nonlinearities.

Note that the  $L^r(\Omega)$ -estimate established in Theorem 1.4 differs with the sharp exponent  $\bar{r}_0$  defined above by the term  $2\sqrt{\frac{n-1}{p-1}}$ . Moreover, observe that  $\bar{r}_1$  is larger than p and tends to it as n goes to infinity. In particular, the best expected regularity independent of the dimension n for the extremal solution  $u^*$  is  $W_0^{1,p}(\Omega)$ , which is the one we obtain in Theorem 1.4.

### **1.3** Outline of the paper

The paper is organized as follows. In section 2 we prove Theorem 1.1 and the geometric type inequalities stated in Theorem 1.2. In section 3 we prove that Theorem 1.2 holds for solutions of (1.2) when  $1 \le q \le 2$ . Moreover we give boundary estimates when the domain is strictly convex. In section 4, we present the semi-stability condition (1.10) and the space of admissible functions  $H_0$ . The rest of the section deals with the regularity of semi-stable solutions proving Theorems 1.3 and 1.4.

### 2 Geometric Hardy-Sobolev type inequalities

In this section we prove Theorems 1.1 and 1.2. As we said in the introduction, the geometric inequalities established in Theorem 1.2 are new for the range  $1 \le q < p$  since the case  $q \ge p$  was proved in [6]. However, we will give the proof in all cases using Schwarz symmetrization, giving an alternative proof for the known range of parameters  $q \ge p$ .

We start recalling the definition of Schwarz symmetrization of a compact set and of a Lipschitz continuous function.

**Definition 2.1.** We define the *Schwarz symmetrization of a compact set*  $D \subset \mathbb{R}^n$  as

$$D^* := \begin{cases} B_R(0) \text{ with } R = (|D|/|B_1|)^{1/n} & \text{if } D \neq \emptyset, \\ \emptyset & \text{if } D = \emptyset. \end{cases}$$

$$v^*(x) := \sup\{t \in \mathbb{R} : x \in \Omega^*_t\}.$$

Equivalently, we can define the Schwarz symmetrization of v as

$$v^*(x) = \inf\{t \ge 0 : V(t) < |B_1| |x|^n\},\$$

where  $V(t) := |\Omega_t| = |\{x \in \Omega : |v(x)| > t\}|$  denotes the distribution function of v.

The first ingredient in the proof of Theorem 1.1 is the isoperimetric inequality for functions v in  $W_0^{1,1}(\Omega)$ :

$$n|B_1|^{1/n}V(t)^{(n-1)/n} \le P(t) := \frac{d}{dt} \int_{\{|v| \le t\}} |\nabla v| \, dx \qquad \text{for a.e. } t > 0, \qquad (2.1)$$

where P(t) stands for the perimeter in the sense of De Giorgi (the total variation of the characteristic function of  $\{x \in \Omega : |v(x)| > t\}$ ).

The second ingredient is the following Sobolev inequality on compact hypersurfaces without boundary due to Michael and Simon [12] and to Allard [1].

**Theorem 2.2** ([1, 12]). Let  $M \subset \mathbb{R}^n$  be a  $C^{\infty}$  immersed (n - 1)-dimensional compact hypersurface without boundary and  $\phi \in C^{\infty}(M)$ . If  $q \in [1, n - 1)$ , then there exists a constant A depending only on n and q such that

$$\left(\int_{M} |\phi|^{q^{\star}} d\sigma\right)^{1/q^{\star}} \le A \left(\int_{M} |\nabla\phi|^{q} + |H\phi|^{q} d\sigma\right)^{1/q},$$
(2.2)

where H is the mean curvature of M,  $d\sigma$  denotes the area element in M, and  $q^{\star} = \frac{(n-1)q}{n-1-q}$ .

As we said in the introduction it is well known that Schwarz symmetrization preserves the  $L^r$ -norm and decreases the  $W^{1,r}$ -norm. Let us prove that it also decreases (up to a multiplicative constant) the functional  $I_{p,q}$  defined in (1.1) using the isoperimetric inequality (2.1) and the geometric inequality (2.2) applied to  $M = M_t = \{x \in \Omega : |v(x)| = t\}$  and  $\phi = |\nabla v|^{(p-1)/q}$ .

Proof of Theorem 1.1. Let  $v \in C_0^{\infty}(\overline{\Omega})$ ,  $p \ge 1$ , and  $1 \le q < n - 1$ . By Sard's theorem, almost every  $t \in (0, \|v\|_{L^{\infty}(\Omega)})$  is a regular value of |v|. By definition, if t is a regular value of |v|, then  $|\nabla v(x)| > 0$  for all  $x \in \Omega$  such that |v(x)| = t. Therefore,  $M_t := \{x \in \Omega : |v(x)| = t\}$  is a  $C^{\infty}$  immersed (n - 1)-dimensional compact hypersurface of  $\mathbb{R}^n$ without boundary for every regular value t. Applying inequality (2.2) to  $M = M_t$  and  $\phi = |\nabla v|^{(p-1)/q}$  we obtain

$$\left(\int_{M_t} |\nabla v|^{(p-1)\frac{q^{\star}}{q}} \, d\sigma\right)^{q/q^{\star}} \le A^q \int_{M_t} \left|\nabla_{T,v} |\nabla v|^{\frac{p-1}{q}}\right|^q + |H_v|^q |\nabla v|^{p-1} \, d\sigma \tag{2.3}$$

for a.e.  $t \in (0, ||v||_{L^{\infty}(\Omega)})$ , where  $H_v$  denotes the mean curvature of  $M_t$ ,  $d\sigma$  is the area element in  $M_t$ , A is the constant in (2.2) which depends only on n and q, and

$$q^\star := \frac{(n-1)q}{n-1-q}$$

Recall that V(t), being a nonincreasing function, is differentiable almost everywhere and, thanks to the coarea formula and that almost every  $t \in (0, ||v||_{L^{\infty}(\Omega)})$  is a regular value of |v|, we have

$$-V'(t) = \int_{M_t} \frac{1}{|\nabla v|} d\sigma \quad \text{and} \quad P(t) = \int_{M_t} d\sigma \quad \text{for a.e. } t \in (0, \|v\|_{L^{\infty}(\Omega)}).$$

Therefore applying Jensen inequality and then using the isoperimetric inequality (2.1), we obtain

$$\left(\int_{M_t} |\nabla v|^{(p-1)\frac{q^*}{q}+1} \frac{d\sigma}{|\nabla v|}\right)^{\frac{q}{q^*}} \ge \frac{P(t)^{p-1+\frac{q}{q^*}}}{\left(-V'(t)\right)^{p-1}} \ge \frac{(A_1V(t)^{\frac{n-1}{n}})^{p-1+\frac{q}{q^*}}}{\left(-V'(t)\right)^{p-1}} \tag{2.4}$$

for a.e.  $t \in (0, ||v||_{L^{\infty}(\Omega)})$ , where  $A_1 := n|B_1|^{1/n}$ .

Note that for radial functions the inequalities in (2.4) are equalities. Therefore, since the Schwarz symmetrization  $v^*$  of v is a radial function and it satisfies (2.3), with an equality and with constant  $A = |\partial B_1|^{-1/(n-1)}$ , we obtain

$$\left(\int_{\{|v^*|=t\}} |\nabla v^*|^{(p-1)\frac{q^*}{q}} \, d\sigma\right)^{q/q^*} = |\partial B_1|^{-\frac{q}{n-1}} \int_{\{v^*=t\}} |H_{v^*}|^q |\nabla v^*|^{p-1} \, d\sigma$$
  
$$= \frac{(A_1 V(t)^{\frac{n-1}{n}})^{p-1+\frac{q}{q^*}}}{(-V'(t))^{p-1}}.$$
(2.5)

for a.e.  $t \in (0, ||v||_{L^{\infty}(\Omega)})$ . Here, we used that  $V(t) = |\{|v| > t\}| = |\{|v^*| > t\}|$  for a.e.  $t \in (0, ||v||_{L^{\infty}(\Omega)})$ .

Therefore, from (2.3), (2.4), and (2.5), we obtain

$$|\partial B_1|^{-\frac{q}{n-1}} \int_{\{v^*=t\}} |H_{v^*}|^q |\nabla v^*|^{p-1} \, d\sigma \le A^q \int_{M_t} \left| \nabla_{T,v} |\nabla v|^{\frac{p-1}{q}} \right|^q + |H_v|^q |\nabla v|^{p-1} \, d\sigma,$$

for a.e.  $t \in (0, \|v\|_{L^{\infty}(\Omega)})$ . Integrating the previous inequality with respect to t on  $(0, \|v\|_{L^{\infty}(\Omega)})$  and using the coarea formula we obtain inequality (1.3), with the explicit constant  $C = A^{\frac{q}{p}} |\partial B_1|^{\frac{q}{(n-1)p}}$ , proving the result.

**Remark 2.3.** We obtained the explicit admissible constant  $C = A^{\frac{q}{p}} |\partial B_1|^{\frac{q}{(n-1)p}}$  in (1.3), where A is the universal constant appearing in (2.2).

We prove Theorem 1.2 using Theorem 1.1 and known results on one dimensional weighted Sobolev inequalities.

Proof of Theorem 1.2. Let  $v \in C_0^{\infty}(\overline{\Omega})$  and  $v^*$  its Schwarz symmetrization. Recall that  $v^*$  is defined in  $B_R$  with  $R = (|\Omega|/|B_1|)^{1/n}$ .

(a) Assume 1 + q < n < p + q. Using Hölder inequality we obtain

$$v^{*}(s) = \int_{s}^{R} |(v^{*})'(\tau)| d\tau$$
  

$$\leq \left( \int_{0}^{R} |(v^{*})'(\tau)|^{p} \tau^{-q} \tau^{n-1} d\tau \right)^{1/p} \left( \int_{s}^{R} \tau^{\frac{1+q-n}{p-1}} d\tau \right)^{1/p'}$$
(2.6)

for a.e.  $s \in (0, R)$ . In particular,

$$v^{*}(s) \leq |\partial B_{1}|^{-1/p} \left(\frac{p-1}{p+q-n}\right)^{1/p'} \left(\frac{|\Omega|}{|B_{1}|}\right)^{\frac{p+q-n}{np}} I_{p,q}(v^{*}; B_{R})$$

for a.e.  $s \in (0, R)$ . We conclude this case, by Theorem 1.1, noting that  $||v||_{L^{\infty}(\Omega)} = v^*(0)$ .

(b) Assume n > p+q. We use the following 1-dimensional weighted Sobolev inequality:

$$\left(\int_{0}^{R} |\varphi(s)|^{p_{q}^{\star}} s^{n-1} \, ds\right)^{1/p_{q}^{\star}} \le C(n, p, q) \left(\int_{0}^{R} s^{-q} |\varphi'(s)|^{p} s^{n-1} \, ds\right)^{1/p} \tag{2.7}$$

with optimal constant

$$C(n,p,q) := \left(\frac{p-1}{n-(p+q)}\right)^{1/p'} n^{-1/p_q^{\star}} \left[\frac{\Gamma\left(\frac{np}{p+q}\right)}{\Gamma\left(\frac{n}{p+q}\right)\Gamma\left(1+\frac{n(p-1)}{p+q}\right)}\right]^{\frac{p+q}{np}}$$
(2.8)

stated in [18]. Applying inequality (2.7) to  $\varphi = v^*$  and noting that the  $L^{p_q^*}$ -norm is preserved by Schwarz symmetrization, we obtain

$$|\partial B_1|^{-1/p_q^{\star}} \left( \int_{\Omega} |v|^{p_q^{\star}} \, dx \right)^{1/p_q^{\star}} \le C(n, p, q) |\partial B_1|^{-1/p} \left( \int_{B_R} |x|^{-q} |\nabla v^{\star}|^p \, dx \right)^{1/p}.$$

Using Theorem 1.1 again we prove (1.8) for  $r = p_q^*$ . The remaining cases,  $1 \le r < p_q^*$ , now follow easily from Hölder inequality.

(c) Assume n = p + q. From (2.6) and Theorem 1.1 we obtain

$$v^{*}(s) \leq \left(\int_{0}^{R} |(v^{*})'(\tau)|^{p} \tau^{-q} \tau^{n-1} d\tau\right)^{1/p} \left(\int_{s}^{R} \tau^{-1} d\tau\right)^{1/p'} \\ \leq |\partial B_{1}|^{-1/p} C I_{p,q}(v;\Omega) \left(\ln\left(\frac{R}{s}\right)\right)^{1/p'}$$

for a.e.  $s \in (0, R)$ . Equivalently

$$\exp\left\{\left(\frac{v^*(s)}{|\partial B_1|^{-1/p}CI_{p,q}(v;\Omega)}\right)^{p'}\right\}|\partial B_1|s^{n-1}\leq \frac{R}{s}|\partial B_1|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^{n-1}|s^$$

for a.e.  $s \in (0, R)$ . Integrating the previous inequality with respect to s in (0, R) we obtain

$$\int_{B_R} \exp\left\{\left(\frac{v^*}{|\partial B_1|^{-1/p} CI_{p,q}(v;\Omega)}\right)^{p'}\right\} dx \le |\partial B_1| \frac{R^n}{n-1} = \frac{n}{n-1} |\Omega|.$$

We conclude the proof noting that the integral in inequality (1.9) is preserved under Schwarz symmetrization.

**Remark 2.4.** Note that we obtained explicit admissible constants  $C_1$ ,  $C_2$ , and  $C_3$  in inequalities of Theorem 1.2. More precisely, we obtained

$$C_{1} = |\partial B_{1}|^{-\frac{1}{p}} \left(\frac{p-1}{p+q-n}\right)^{\frac{1}{p'}} \left(\frac{|\Omega|}{|B_{1}|}\right)^{\frac{p+q-n}{np}} A^{\frac{q}{p}} |\partial B_{1}|^{\frac{q}{(n-1)p}},$$
$$C_{2} = C(n,p,q) |\partial B_{1}|^{\frac{1}{p_{q}^{*}} - \frac{1}{p}} A^{\frac{q}{p}} |\partial B_{1}|^{\frac{q}{(n-1)p}},$$

and

$$C_3 = |\partial B_1|^{-\frac{1}{p}} A^{\frac{n-p}{p}} |\partial B_1|^{\frac{n-p}{(n-1)p}},$$

where A is the universal constant appearing in (2.2) and C(n, p, q) is defined in (2.8).

All the constants  $C_i$  depend only on n, p, and q. However, the best constant A in (2.2) is unknown (even for mean convex hypersurfaces). Behind this Sobolev inequality there is the following geometric isoperimetric inequality

$$|M|^{\frac{n-2}{n-1}} \le A_2 \int_M |H(x)| \, d\sigma.$$
 (2.9)

Here,  $M \subset \mathbb{R}^n$  is a  $C^{\infty}$  immersed (n-1)-dimensional compact hypersurface without boundary and H is the mean curvature of M as in Theorem 2.2. The best constant in (2.9) is also unknown even for mean convex hypersurfaces.

## **3** Properties of solutions of *p*-Laplace equations

In this section, we first establish an *a priori*  $L^{\infty}$  estimate in a neighborhood of the boundary  $\partial\Omega$  for any regular solution u of (1.2) when the domain  $\Omega$  is strictly convex. More precisely, we prove that there exists positive constants  $\varepsilon$  and  $\gamma$ , depending only on the domain  $\Omega$ , such that

$$\|u\|_{L^{\infty}(\Omega_{\varepsilon})} \leq \frac{1}{\gamma} \|u\|_{L^{1}(\Omega)}, \quad \text{where } \Omega_{\varepsilon} := \{x \in \Omega : \operatorname{dist}(x, \partial\Omega) < \varepsilon\}.$$
(3.1)

Then, we establish that the geometric inequalities of Theorem 1.2 still hold for solutions of (1.2) in the smaller range  $1 \le q \le 2$ . In the next section, these two ingredients will allow us to obtain *a priori* estimates for semi-stable solutions.

Let  $u \in W_0^{1,p}(\Omega)$  be a weak solution (*i.e.*, a solution in the distributional sense) of the problem

$$\begin{cases}
-\Delta_p u = g(u) & \text{in } \Omega, \\
u > 0 & \text{in } \Omega, \\
u = 0 & \text{on } \partial\Omega,
\end{cases}$$
(3.2)

where  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ , with  $n \ge 2$ , and g is any positive smooth nonlinearity.

We say that  $u \in W_0^{1,p}(\Omega)$  is a *regular solution* of (3.2) if it satisfies the equation in the distributional sense and  $g(u) \in L^{\infty}(\Omega)$ . By well known regularity results for degenerate elliptic equations, one has that every regular solution u belongs to  $C^{1,\alpha}(\Omega)$  for some  $\alpha \in (0, 1]$  (see [8, 16]). Moreover,  $u \in C^1(\overline{\Omega})$  (see [11]). This is the best regularity that one can hope for solutions of p-Laplace equations. Therefore, equation (3.2) is always meant in a distributional sense.

We prove the boundary *a priori* estimate (3.1) through a moving plane procedure for the *p*-Laplacian which is developed in [9].

**Proposition 3.1.** Let  $\Omega$  be a smooth bounded domain of  $\mathbb{R}^n$  and g any positive smooth function. Let u be any positive regular solution of (3.2).

If  $\Omega$  is strictly convex, then there exist positive constants  $\varepsilon$  and  $\gamma$  depending only on the domain  $\Omega$  such that for every  $x \in \Omega$  with  $dist(x, \partial \Omega) < \varepsilon$ , there exists a set  $I_x \subset \Omega$ with the following properties:

$$|I_x| \ge \gamma$$
 and  $u(x) \le u(y)$  for all  $y \in I_x$ .

As a consequence,

$$\|u\|_{L^{\infty}(\Omega_{\varepsilon})} \leq \frac{1}{\gamma} \|u\|_{L^{1}(\Omega)}, \quad \text{where } \Omega_{\varepsilon} := \{x \in \Omega : \operatorname{dist}(x, \partial\Omega) < \varepsilon\}.$$
(3.3)

*Proof.* First let us observe that from the regularity of the solution u up to the boundary  $\partial \Omega$  and the fact that  $\Delta_p u \leq 0$ , we can apply the generalized Hopf boundary lemma [19] to see that the normal derivative  $\frac{\partial u}{\partial \nu} < 0$  on  $\partial \Omega$ . Thus, if we let  $Z_u := \{x \in \Omega : \nabla u(x) = 0\}$  be the critical set of u, we have that  $Z_u \cap \partial \Omega = \emptyset$ . By the compactness of both sets, there exists  $\varepsilon_0 > 0$  such that  $Z_u \cap \Omega_{\varepsilon} = \emptyset$  for any  $\varepsilon \leq \varepsilon_0$ .

We will now prove that this neighborhood of the boundary is in fact independent of the solution u. In order to begin a moving plane argument we need some notations: let  $e \in S^{n-1}$  be any direction and for  $\lambda \in \mathbb{R}$  let us consider the hyperplane

$$T = T_{\lambda,e} = \{ x \in \mathbb{R}^n : x \cdot e = \lambda \}$$

and the corresponding cap

$$\Sigma = \Sigma_{\lambda, e} = \{ x \in \Omega : x \cdot e < \lambda \}.$$

Set

$$a(e) = \inf_{x \in \Omega} x \cdot e$$

and for any  $x \in \Omega$ , let  $x' = x_{\lambda,e}$  be its reflection with respect to the hyperplane T, *i.e.*,

$$x' = x + (\lambda - 2x \cdot e) e.$$

For any  $\lambda > a(e)$  the cap

$$\Sigma' = \{ x \in \Omega : x' \in \Sigma \}$$

is the (non-empty) reflected cap of  $\Sigma$  with respect to T.

Furthermore, consider the function  $v(x) = u(x') = u(x_{\lambda,e})$ , which is just the reflected of u with respect to the same hyperplane. By the boundedness of  $\Omega$ , for  $\lambda - a(e)$  small, we have that the corresponding reflected cap  $\Sigma'$  is contained in  $\Omega$ . Moreover, by the strict convexity of  $\Omega$ , there exists  $\lambda_0 = \lambda_0(\Omega)$  (independent of e) such that  $\Sigma'$  remains in  $\Omega$  for any  $\lambda \leq \lambda_0$ .

Let us then compare the function u and its reflection v for such values of  $\lambda$  in the cap  $\Sigma$ . First of all, both functions solve the same equation since  $\Delta_p$  is invariant under reflection; secondly, on the hyperplane T the functions coincide, whereas for any  $x \in \partial \Sigma \cap \partial \Omega$  we have that u(x) = 0 and that v(x) = u(x') > 0, since the reflection  $x' \in \Omega$ . Hence we can see that:

$$\Delta_p(u) + f(u) = \Delta_p(v) + f(v) \text{ in } \Sigma, \quad u \le v \text{ on } \partial \Sigma.$$

Again by the boundedness of  $\Omega$ , if  $\lambda - a(e)$  is small, the measure of the cap  $\Sigma$  will be small. Therefore, from the Comparison Principle in small domains (see [9]) we have that  $u \leq v$  in  $\Sigma$ . Moreover, by Strong Comparison Principle and Hopf Lemma, we see that  $u \leq v$  in  $\Sigma_{\lambda,e}$  for any  $a(e) < \lambda \leq \lambda_0$ . In particular, this spells that u(x) is nondecreasing in the *e* direction for all  $x \in \Sigma$ .

Now, fix  $x_0 \in \partial\Omega$  and let  $e = \nu(x_0)$  be the unit normal to  $\partial\Omega$  at  $x_0$ . By the convexity assumption  $T_{a(\nu(x_0)),\nu(x_0)} \cap \partial\Omega = \{x_0\}$ . If we let  $\theta \in S^{n-1}$  be another direction close to the outer normal  $\nu(x_0)$ , the reflection of the caps  $\Sigma_{\lambda,\theta}$  with respect to the hyperplane  $T_{\lambda,\theta}$ (which is close to the tangent one) would still be contained in  $\Omega$  thanks to its strict convexity. So the above argument could be applied also to the new direction  $\theta$ . In particular, we see that we can get a neighborhood  $\Theta$  of  $\nu(x_0)$  in  $S^{n-1}$  such that u(x) is nondecreasing in every direction  $\theta \in \Theta$  and for any x such that  $x \cdot \theta < \frac{\lambda_0}{2}$ .

By eventually taking a smaller neighborhood  $\Theta$ , we may assume that

$$|x \cdot (\theta - \nu(x_0))| < \lambda_0/8$$

for any  $x \in \Sigma_{\lambda_0,\theta}$  and  $\theta \in \Theta$ . Moreover, noticing that

$$x \cdot \theta = x \cdot (\theta - \nu(x_0)) + x \cdot \nu(x_0)$$

and

$$\frac{\lambda_0}{2} = \frac{\lambda_0}{8} + \frac{3\lambda_0}{8} > x \cdot \theta > \frac{\lambda_0}{8} - \frac{\lambda_0}{8} = 0$$

it is then easy to see that u is nondecreasing in any direction  $\theta \in \Theta$  on  $\Sigma_0 = \{x \in \Omega : \frac{\lambda_0}{8} < x \cdot \nu(x_0) < \frac{3\lambda_0}{8}\}.$ 

Finally, let us choose  $\varepsilon = \frac{\lambda_0}{8}$ . Fix any point  $x \in \Omega_{\varepsilon}$  and let  $x_0$  be its projection onto  $\partial \Omega$ . From the above arguments we see that

$$u(x) \le u(x_0 - \varepsilon \nu(x_0)) \le u(y)$$

for any  $y \in I_x$ , where  $I_x \subset \Sigma_0$  is a truncated cone with vertex at  $x_1$ , opening angle  $\Theta$ , and height  $\frac{\lambda_0}{4}$ . Hence, we have obtained that there exists a positive constant  $\gamma = \gamma(\Omega, \varepsilon)$  such that  $|I_x| \ge \gamma$  and  $u(x) \le u(y)$  for any  $y \in I_x$ . Finally, choosing  $x_{\varepsilon}$  as the maximum of uin  $\Omega_{\varepsilon}$ , we get

$$\|u\|_{L^{\infty}(\Omega_{\varepsilon})} = u_{\varepsilon}(x_{\varepsilon}) \le \frac{1}{\gamma} \int_{I_{x_{\varepsilon}}} u(y) \, dy \le \frac{1}{\gamma} \|u\|_{L^{1}(\Omega)}$$

which proves (3.3).

We will now prove that inequalities in Theorem 1.2 are also valid for a positive solution u of (3.2) in the smaller range  $1 \le q \le 2$ . To do this, we will construct an approximation of u through smooth functions and see that, thanks to strong uniform estimates on this approximation, we can pass to the limit in all of the inequalities.

**Proposition 3.2.** Let  $\Omega$  be a smooth bounded domain of  $\mathbb{R}^n$  and g any positive smooth function. Let u be any positive regular solution of (3.2). If  $1 \le q \le 2$ , then inequalities in Theorem 1.2 hold for v = u. Given s > 0, the same holds true also for v = u - s and  $\Omega$  replaced by  $\Omega_s := \{x \in \Omega : u > s\}$ .

*Proof.* Let  $Z_u = \{x \in \Omega : \nabla u(x) = 0\}$ . Recall that by standard elliptic regularity  $u \in C^{\infty}(\Omega \setminus Z_u)$  and that  $|Z_u| = 0$  by [9]. Therefore, u is smooth almost everywhere in  $\Omega$ . Let  $x \in \Omega \setminus Z_u$  and observe that for the mean curvature  $H_u$  of the level set passing through x we have the following explicit expression

$$-(n-1)H_u = \operatorname{div}\left(\frac{\nabla u}{|\nabla u|}\right) = \frac{\Delta u}{|\nabla u|} - \frac{\langle D^2 u \nabla u, \nabla u \rangle}{|\nabla u|^3}$$
(3.4)

whereas for the tangential gradient term we have

$$\nabla_{T,u} |\nabla u| = \frac{D^2 u \nabla u}{|\nabla u|} - \frac{\langle D^2 u \nabla u, \nabla u \rangle \nabla u}{|\nabla u|^3}, \tag{3.5}$$

where all the terms in these expressions are evaluated at x. Hence, there exists a positive constant C = C(n, p, q) such that

$$\left(\frac{1}{p'}|\nabla_{T,u}|\nabla u|^{\frac{p}{q}}|\right)^{q} + |H_{u}|^{q}|\nabla u|^{p} \le C|D^{2}u|^{q}|\nabla u|^{p-q} \quad \text{for a.e. } x \in \Omega.$$
(3.6)

From [9] we recall the following important estimate: for any  $1 \le q \le 2$  there holds

$$\int_{\Omega} |D^2 u|^q |\nabla u|^{p-q} \, dx < \infty. \tag{3.7}$$

Thanks to (3.6) and (3.7), all of the integrals in the geometric Hardy-Sobolev inequalities are well defined for any  $1 \le q \le 2$ .

However, since the solution u is not smooth around  $Z_u$ , we need to regularize u in a neighborhood of the critical set in order to apply the inequalities of Theorem 1.2. We will now describe an approximation argument due to Canino, Le, and Sciunzi [7] for the  $p(\cdot)$ -Laplacian (in our case  $p(x) \equiv p$  constant).

**Lemma 3.3** ([7]). Let  $D \subset \Omega$  be an open set,  $1 \leq q \leq 2$ , and  $\varepsilon \in (0, 1)$ . Let  $u \in C^1(\overline{\Omega})$ be a solution of (1.2) and h := g(u). If  $h_{\varepsilon} \in C^{\infty}(\overline{D})$  is any sequence converging to h in  $C^1(\overline{D})$  as  $\varepsilon \downarrow 0$ , then the unique solution  $v_{\varepsilon}$  of the following regularized problem

$$\begin{cases} -\operatorname{div}\left(\left(\varepsilon^{2}+|\nabla v_{\varepsilon}|^{2}\right)^{\frac{p-2}{2}}\nabla v_{\varepsilon}\right) &= h_{\varepsilon}(x) \quad \text{in } D, \\ v_{\varepsilon} &= u \quad \text{on } \partial D. \end{cases}$$
(3.8)

tends to u strongly in  $W^{1,p}(B)$ . Moreover, there exists a constant C independent of  $\varepsilon$  such that

$$\int_{D} |D^{2} v_{\varepsilon}|^{q} (\varepsilon^{2} + |\nabla v_{\varepsilon}|^{2})^{\frac{p-q}{2}} dx \le C$$

and

$$\lim_{\varepsilon \to 0} \int_D |D^2 v_\varepsilon|^q (\varepsilon^2 + |\nabla v_\varepsilon|^2)^{\frac{p-q}{2}} dx = \int_D |D^2 u|^q |\nabla u|^{p-q} dx.$$
(3.9)

Let  $v_{\varepsilon} \in C^{\infty}(D)$  be the unique solution of (3.8) and let us consider a smooth cutoff function  $\eta$  with compact support contained in  $\Omega$  and such that  $\eta \equiv 1$  on D. We can construct a smooth regularization  $u_{\varepsilon}$  of u defining  $u_{\varepsilon} := (1 - \eta)u + \eta v_{\varepsilon}$ . We can then apply Theorem 1.2 to any  $u_{\varepsilon}$  to get the appropriate inequality (a), (b), or (c). From [8, 11] and standard elliptic regularity we know that the regularization  $u_{\varepsilon}$  will converge to u, as  $\varepsilon \downarrow 0$ , both in  $C^1(\overline{\Omega})$  and  $C^2(\overline{\Omega} \setminus Z_u)$ . Hence we can easily pass to the limit as  $\varepsilon \downarrow 0$  in the left hand side of (1.7) and (1.8).

In order to see that also the remaining terms  $I_{p,q}(u_{\varepsilon}; \Omega)$  which involve tangential gradient and mean curvature behave well under this approximation the argument is the following. Splitting the domain  $\Omega$  and recalling that  $u_{\varepsilon} \equiv v_{\varepsilon}$  in D we have that:

$$I_{p,q}(u_{\varepsilon};\Omega) = I_{p,q}(u_{\varepsilon};D) + I_{p,q}(u_{\varepsilon};\Omega \setminus D) = I_{p,q}(v_{\varepsilon};D) + I_{p,q}(u_{\varepsilon};\Omega \setminus D).$$

Clearly, from the  $C^2$  convergence we have that  $I_{p,q}(u_{\varepsilon}; \Omega \setminus D) \to I_{p,q}(u; \Omega \setminus D)$  as  $\varepsilon \downarrow 0$ . Therefore we can concentrate on the convergence of  $I_{p,q}(v_{\varepsilon}; D)$ .

From (3.4), (3.5), and through a simple expansion of  $(\varepsilon^2 + |\nabla v_{\varepsilon}|^2)^{\frac{p-q}{2}}$  around  $\varepsilon = 0$ , we see that for a sufficiently small  $\varepsilon_0 > 0$  there exists a constant  $K = K(n, p, q, \varepsilon_0) > 0$  such that for any  $\varepsilon \leq \varepsilon_0$  we have

p-Laplace equations and geometric Sobolev inequalities

$$\left(\frac{1}{p'}|\nabla_{T,v_{\varepsilon}}|\nabla v_{\varepsilon}|^{\frac{p}{q}}|\right)^{q} + |H_{v_{\varepsilon}}|^{q}|\nabla v_{\varepsilon}|^{p} \le K |D^{2}v_{\varepsilon}|^{q} (\varepsilon^{2} + |\nabla v_{\varepsilon}|^{2})^{\frac{p-q}{2}}.$$
(3.10)

Moreover, by the fact that  $v_{\varepsilon} \to u$  in  $C^2(D \setminus Z_u)$  and  $|Z_u| = 0$ , almost everywhere in D we have

$$\lim_{\varepsilon \to 0} \left( \frac{1}{p'} |\nabla_{T, v_{\varepsilon}} |\nabla v_{\varepsilon}|^{\frac{p}{q}} | \right)^{q} + |H_{v_{\varepsilon}}|^{q} |\nabla v_{\varepsilon}|^{p} = \left( \frac{1}{p'} |\nabla_{T, u} |\nabla u|^{\frac{p}{q}} | \right)^{q} + |H_{u}|^{q} |\nabla u|^{p}.$$
(3.11)

Now, thanks to (3.9), (3.10), and (3.11), by dominated convergence theorem we see that:

$$\lim_{\varepsilon \to 0} \int_D \left( \frac{1}{p'} |\nabla_{T, v_\varepsilon} |\nabla v_\varepsilon|^{\frac{p}{q}} | \right)^q + |H_{v_\varepsilon}|^q |\nabla v_\varepsilon|^p \, dx$$
$$= \int_D \left( \frac{1}{p'} |\nabla_{T, u} |\nabla u|^{\frac{p}{q}} | \right)^q + |H_u|^q |\nabla u|^p \, dx.$$

Thus, the assertions of Theorem 1.2 hold for v = u.

To conclude the proof let us fix any s > 0 and consider v = u - s on  $\Omega_s = \{x \in \Omega : u > s\}$ . It is clear that the integrands in the inequalities remain unchanged in this case, so the only problem comes from the fact  $\Omega_s$  might not be smooth. If this is the case, let us consider two sequences  $\varepsilon_n \to 0$  and  $s_n \to s$ , with the corresponding regularizations of v given by  $v_n := v_{\varepsilon_n} = u_{\varepsilon_n} - s_n$ . Thanks to the smoothness of any  $v_n$  and Sard Lemma, we can choose each  $s_n$  as a regular value of  $v_n$ , so that the level set  $\{v_n > 0\} = \{u_n > s_n\}$  is smooth. Moreover, from the  $C^1$  convergence, it is clear that for the characteristic functions we have  $\chi_{\{u_n > s_n\}} \to \chi_{\{u > s\}}$ . Hence we can conclude the proof using the same dominated convergence argument as above.

# 4 Regularity of stable solutions. Proof of Theorems 1.3 and 1.4

We are now ready to establish  $L^r$  and  $W^{1,r}$  *a priori* estimates of semi-stable solutions to *p*-Laplace equations proving Theorems 1.3 and 1.4.

Before the proof our regularity results let us recall some known facts on the linearized operator associated to (1.2) and semi-stable solutions.

### 4.1 Linearized operator and semi-stable solutions

This subsection deals with the linearized operator at any regular semi-stable solution  $u \in C_0^1(\overline{\Omega})$  of

where  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^n$ , with  $n \geq 2$ , and g is any positive  $C^1$  nonlinearity.

The linearized operator  $L_u$  associated to (4.1) at u is defined by duality as

$$L_u(v,\phi) := \int_{\Omega} |\nabla u|^{p-2} \left\{ \nabla v \cdot \nabla \phi + (p-2) \left( \nabla v \cdot \frac{\nabla u}{|\nabla u|} \right) \left( \nabla \phi \cdot \frac{\nabla u}{|\nabla u|} \right) \right\} dx$$
$$- \int_{\Omega} g'(u) v \phi \, dx$$

for all  $(v, \phi) \in H_0 \times H_0$ , where the Hilbert space  $H_0$  is defined according to [9] as follows.

**Definition 4.1.** Let  $u \in C_0^1(\overline{\Omega})$  be a regular semi-stable solution of (4.1). We introduce the following weighted  $L^2$ -norm of the gradient

$$|\phi| := \left(\int_{\Omega} \rho |\nabla \phi|^2 dx\right)^{1/2}$$
 where  $\rho := |\nabla u|^{p-2}$ .

According to [9], the space

$$H^1_{\rho}(\Omega) := \{\phi \in L^2(\Omega) \text{ weakly differentiable } : \ |\phi| < +\infty\}$$

is a Hilbert space and is the completion of  $C^{\infty}(\Omega)$  with respect to the  $|\cdot|$ -norm.

We define the Hilbert space  $H_0$  of admissible test functions as

$$H_0 := \left\{ \begin{array}{ll} \{\phi \in H_0^1(\Omega): \ |\phi| < +\infty\} & \mbox{if} \quad 1 < p \le 2 \\ \\ \mbox{the closure of } C_0^\infty(\Omega) \mbox{ in } H_\rho^1(\Omega) & \mbox{if} \quad p > 2. \end{array} \right.$$

Note that for  $1 , <math>H_0$  is a subspace of  $H_0^1(\Omega)$  and since

$$\int_{\Omega} |\nabla \phi|^2 \le \|\nabla u\|_{L^{\infty}(\Omega)}^{2-p} |\phi|^2,$$

we see that  $(H_0, |\cdot|)$  is a Hilbert space. For p > 2, the weight  $\rho = |\nabla u|^{p-2}$  is in  $L^{\infty}(\Omega)$  and satisfies  $\rho^{-1} \in L^1(\Omega)$ , as shown in [9].

Now, thanks to the above definition, the operator  $L_u$  is well defined for  $\phi \in H_0$  and, therefore, the semistability of the solution u reads as

$$L_u(\phi,\phi) = \int_{\Omega} |\nabla u|^{p-2} \left\{ |\nabla \phi|^2 + (p-2) \left( \nabla \phi \cdot \frac{\nabla u}{|\nabla u|} \right)^2 \right\} - g'(u)\phi^2 \, dx \ge 0, \quad (4.2)$$

for every  $\phi \in H_0$ .

On the one hand, considering  $\phi = |\nabla u|\eta$  as a test function in the semistability condition (4.2) for u, we obtain

$$\int_{\Omega} \left[ (p-1) |\nabla u|^{p-2} |\nabla_{T,u}| \nabla u||^2 + B_u^2 |\nabla u|^p \right] \eta^2 \, dx \le (p-1) \int_{\Omega} |\nabla u|^p |\nabla \eta|^2 \, dx \quad (4.3)$$

for any Lipschitz continuous function  $\eta$  with compact support. Here,  $B_u^2$  denotes the  $L^2$ norm of the second fundamental form of the level set of |u| through x (*i.e.*, the sum of the squares of its principal curvatures). The fact that  $\phi = \eta |\nabla u|$  is an admissible test function derives from the estimate (3.7), whereas the computations behind (4.3) are done in [10] (see Theorem 2.5 [10]).

On the other hand, noting that  $(n-1)H_u^2 \leq B_u^2$  and

$$|\nabla u|^{p-2} |\nabla_{T,u}| \nabla u||^2 = \frac{4}{p^2} |\nabla_{T,u}| \nabla u|^{\frac{p}{2}}|^2,$$

we obtain the key inequality to prove our regularity results for semi-stable solutions

$$\int_{\Omega} \left( \frac{4}{p^2} |\nabla_{T,u}| \nabla u|^{p/2} |^2 + \frac{n-1}{p-1} H_u^2 |\nabla u|^p \right) \eta^2 \, dx \le \int_{\Omega} |\nabla u|^p |\nabla \eta|^2 \, dx \tag{4.4}$$

for any Lipschitz continuous function  $\eta$  with compact support.

#### 4.2 *A priori* estimates of stable solutions. Proof of Theorem 1.3

In order to prove the gradient estimate (1.13) established in Theorem 1.3 (b) we will use the following result. Its proof is based on a technique introduced by Bénilan *et al.* [2] to obtain the regularity of entropy solutions for *p*-Laplace equations with  $L^1$  data.

**Proposition 4.2.** Assume  $n \ge 3$  and  $h \in L^1(\Omega)$ . Let u be the entropy solution of

$$\begin{cases} -\Delta_p u = h(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$
(4.5)

Let  $r_0 \ge (p-1)n/(n-p)$ . If  $\int_{\Omega} |u|^{r_0} dx < +\infty$ , then the following a priori estimate holds:

$$\int_{\Omega} |\nabla u|^r \, dx \le r |\Omega| + \left(\frac{r_1}{r} - 1\right)^{-1} \left(\int_{\Omega} |u|^{r_0} \, dx + \|h\|_{L^1(\Omega)}\right)$$

for all  $r < r_1 := pr_0/(r_0 + 1)$ .

**Remark 4.3.** Bénilan *et al.* [2] proved the existence and uniqueness of entropy solutions to problem (4.5). Moreover, they proved that  $|\nabla u|^{p-1} \in L^r(\Omega)$  for all  $1 \le r < n/(n-1)$ and  $|u|^{p-1} \in L^r(\Omega)$  for all  $1 \le r < n/(n-p)$ . Proposition 4.2 establishes an improvement of the previous gradient estimate knowing an *a priori* estimate of  $\int_{\Omega} |u|^{r_0} dx$  for some  $r_0 > (p-1)n/(n-p)$ .

*Proof of Proposition* 4.2. Multiplying (4.5) by  $T_s u = \max\{-s, \min\{s, u\}\}$  we obtain

$$\int_{\{|u| \le s\}} |\nabla u|^p \, dx = \int_{\Omega} h(x) T_s u \, dx \le s \|h\|_{L^1(\Omega)}.$$

Let  $t = s^{(r_0+1)/p}$ . From the previous inequality, recalling that  $V(s) = |\{x \in \Omega : |u| > s\}|$ , we deduce

$$\begin{split} s^{r_0}|\{|\nabla u| > t\}| &\leq s^{r_0} \int_{\{|\nabla u| > t\} \cap \{|u| \le s\}} \left(\frac{|\nabla u|}{t}\right)^p dx + s^{r_0} \int_{\{|u| > s\}} dx \\ &\leq \|h\|_{L^1(\Omega)} + s^{r_0} V(s) \quad \text{for a.e. } s > 0. \end{split}$$

In particular

$$t^{\frac{pr_0}{r_0+1}} |\{|\nabla u| > t\}| \le ||h||_{L^1(\Omega)} + \sup_{\tau > 0} \left\{\tau^{r_0} V(\tau)\right\} \quad \text{for a.e. } t > 0.$$
(4.6)

Moreover, since

$$\tau^{r_0}V(\tau) \le \tau^{r_0} \int_{\{|u|>\tau\}} \left(\frac{|u|}{\tau}\right)^{r_0} dx \le \int_{\Omega} |u|^{r_0} dx \quad \text{for a.e. } \tau > 0,$$

we have  $\sup_{\tau>0} \left\{ \tau^{r_0} V(\tau) \right\} \leq \int_{\Omega} |u|^{r_0} dx.$ Let  $r < r_r := \frac{nr_0}{r_0} / (r_0 + 1)$  From (4.6)

Let  $r < r_1 := pr_0/(r_0 + 1)$ . From (4.6) and the previous inequality, we have

$$\int_{\Omega} |\nabla u|^r \, dx = r \int_0^\infty t^{r-1} |\{ |\nabla u| > t\} | \, dt$$
  
$$\leq r |\Omega| + r \left( \int_{\Omega} |u|^{r_0} \, dx + ||h||_{L^1(\Omega)} \right) \int_1^\infty t^{r-1} t^{-\frac{pr_0}{r_0+1}} \, dt$$

proving the proposition.

Now, we have all the ingredients to prove the *a priori* estimates established in Theorem 1.3 for semi-stable solutions. It will follow from Theorem 1.2 and Propositions 3.2 and 4.2 choosing adequate test functions in the semistability condition (4.4).

First, we prove Theorem 1.3 when  $n \neq p + 2$ . We will take  $\eta = T_s u = \min\{s, u\}$  as a test function in (4.4) and then, thanks to Proposition 3.2, we apply our Morrey and Sobolev inequalities (1.7) and (1.8) with q = 2.

Proof of Theorem 1.3 for  $n \neq p+2$ . Assume  $n \neq p+2$ . Let  $u \in C_0^1(\overline{\Omega})$  be a semi-stable solution of (1.2). By taking  $\eta = T_s u = \min\{s, u\}$  in the semistability condition (4.4) we obtain

$$\int_{\{u>s\}} \left(\frac{4}{p^2} |\nabla_{T,u}| \nabla u|^{p/2}|^2 + \frac{n-1}{p-1} H_u^2 |\nabla u|^p\right) \, dx \le \frac{1}{s^2} \int_{\{u$$

for a.e. s > 0. In particular,

$$\min\left(\frac{4}{(n-1)p},1\right)I_{p,2}(u-s;\{x\in\Omega:u>s\})^p \le \frac{p-1}{(n-1)s^2}\int_{\{u$$

for a.e. s > 0, where  $I_{p,2}$  is the functional defined in (1.1) with q = 2. By Proposition 3.2 we can apply Theorem 1.2 with  $\Omega$  replaced by  $\{x \in \Omega : u > s\}$ , v = u - s, and q = 2. Then, the  $L^r$  estimates established in parts (a) and (b) follow directly from the Morrey and Sobolev type inequalities (1.7) and (1.8).

Finally, the gradient estimate (1.13) follows directly from Proposition 4.2 with  $r_0 = np/(n-p-2)$ .

Now, we deal with the proof of Theorem 1.3 (a) when n = p + 2. This critical case follows from Theorem 2.2 and the semistability condition (4.4) with the test function  $\eta = \eta(u)$  defined in (4.11) and (4.10) below.

*Proof of Theorem* 1.3 *when* n = p + 2. Assume n = p + 2 (and hence, n > 3). Taking a Lipschitz function  $\eta = \eta(u)$  (to be chosen later) in (4.3) and using the coarea formula we obtain

$$C \int_{0}^{\infty} \int_{\{u=t\}} \left\{ \left| \nabla_{T,u} |\nabla u|^{\frac{p-1}{2}} \right|^{2} + \left| H_{u} |\nabla u|^{\frac{p-1}{2}} \right|^{2} \right\} \eta(t)^{2} \, d\sigma dt$$

$$\leq \int_{0}^{\infty} \int_{\{u=t\}} |\nabla u|^{p+1} \, \dot{\eta}(t)^{2} \, d\sigma dt,$$
(4.7)

where  $d\sigma$  denotes the area element in  $\{u = t\}$  and C, here and in the rest of the proof, is a constant depending only on p.

To apply the Sobolev inequality (2.2) in the left hand side of the previous inequality we need to make an approximation argument. Consider the sequence  $u_k$  of smooth regularizations of u introduced in the proof of Proposition 3.2 and note that  $\{u_k = t\}$  is a smooth hypersurface for a.e.  $t \ge 0$ . Then, from the Sobolev inequality (2.2) with  $\phi = |\nabla u_k|^{\frac{p-1}{2}}$ , q = 2, and  $M = \{u_k = t\}$ , and noting that

$$(p-1)\frac{n-1}{n-3} = p+1$$
 when  $n = p+2$ ,

we obtain

$$C \int_{0}^{\infty} \left( \int_{\{u_{k}=t\}} |\nabla u_{k}|^{p+1} \right)^{\frac{n-3}{n-1}} \eta(t)^{2} \, d\sigma \, dt$$

$$\leq \int_{0}^{\infty} \int_{\{u_{k}=t\}} \left\{ \left| \nabla_{T,u_{k}} |\nabla u_{k}|^{\frac{p-1}{2}} \right|^{2} + \left| H_{u_{k}} |\nabla u_{k}|^{\frac{p-1}{2}} \right|^{2} \right\} \, \eta(t)^{2} \, d\sigma \, dt.$$

$$(4.8)$$

Now, we will pass to the limit in the previous inequality. Note that, if  $\eta$  is bounded, through a dominated convergence argument as in Proposition 3.2 we have

$$\lim_{k \to \infty} \int_0^\infty \int_{\{u_k = t\}} \left\{ \left| \nabla_{T, u_k} |\nabla u_k|^{\frac{p-1}{2}} \right|^2 + \left| H_{u_k} |\nabla u_k|^{\frac{p-1}{2}} \right|^2 \right\} \eta(t)^2 \, d\sigma dt$$
$$= \int_0^\infty \int_{\{u = t\}} \left\{ \left| \nabla_{T, u} |\nabla u|^{\frac{p-1}{2}} \right|^2 + \left| H_u |\nabla u|^{\frac{p-1}{2}} \right|^2 \right\} \eta(t)^2 \, d\sigma dt.$$

Moreover, from the  $C^1$  convergence of  $u_k$  to u we obtain

$$\lim_{k \to \infty} \int_0^\infty \left( \int_{\{u_k=t\}} |\nabla u_k|^{p+1} \right)^{\frac{n-3}{n-1}} \eta(t)^2 \, d\sigma \, dt = \int_0^\infty \left( \int_{\{u=t\}} |\nabla u|^{p+1} \right)^{\frac{n-3}{n-1}} \eta(t)^2 \, d\sigma \, dt.$$

Therefore, taking the limit as k goes to infinity in (4.8) and using (4.7), we get

$$C\int_{0}^{\infty}\psi(t)^{\frac{n-3}{n-1}}\eta(t)^{2}\,dt \leq \int_{0}^{\infty}\psi(t)\,\dot{\eta}(t)^{2}\,dt = \int_{0}^{\infty}\int_{\{u=t\}}|\nabla u|^{p+1}\,d\sigma\,\dot{\eta}(t)^{2}\,dt,\quad(4.9)$$

where

$$\psi(t) := \int_{\{u=t\}} |\nabla u|^{p+1} \, d\sigma. \tag{4.10}$$

Now, let  $\bar{M} := \|u\|_{L^{\infty}(\Omega)}$ . Given s > 0, choose

$$\eta(t) = \eta_s(t) := \begin{cases} t/s & \text{if } 0 \le t \le s, \\ \exp\left(\frac{1}{\sqrt{2}} \int_s^t \left(\frac{C\psi(\tau)^{\frac{n-3}{n-1}}}{\psi(\tau)}\right)^{\frac{1}{2}} d\tau\right) & \text{if } s < t \le \bar{M} \\ \eta(M) & \text{if } t > \bar{M}. \end{cases}$$
(4.11)

It is then clear that

$$\int_{0}^{\bar{M}} \int_{\{u=t\}} |\nabla u|^{p+1} \, d\sigma \, \dot{\eta}_{s}(t)^{2} \, dt = \frac{1}{s^{2}} \int_{\{u\leq s\}} |\nabla u|^{p+2} \, dx + \frac{C}{2} \int_{s}^{\bar{M}} \psi(t)^{\frac{n-3}{n-1}} \, \eta_{s}(t)^{2} \, dt.$$

Therefore, from (4.9) we obtain

$$\frac{C}{2} \int_{s}^{M} \psi(t)^{\frac{n-3}{n-1}} \eta_{s}(t)^{2} dt \leq \frac{1}{s^{2}} \int_{\{u \leq s\}} |\nabla u|^{p+2} dx.$$
(4.12)

Let us choose  $\alpha = \frac{2}{n-2}$ ,  $\beta = \frac{n-3}{(n-2)(n-1)}$ , and m = n-2. Note that  $\alpha, \beta > 0$ , m > 1, and  $\beta m' = 1/(n-1)$ . Moreover, using the definition of  $\eta_s$  we have

$$\frac{1}{\psi(t)^{\beta m'} \eta_s(t)^{\alpha m'}} = \sqrt{\frac{2}{C}} \frac{\dot{\eta}_s(t)}{\eta_s(t)^{\alpha m'+1}}$$
(4.13)

for all t > s. By (4.13), Hölder inequality, and (4.12), we see that

$$\begin{split} \bar{M} - s &= \int_{s}^{M} \frac{\psi(t)^{\beta} \eta_{s}(t)^{\alpha}}{\psi(t)^{\beta} \eta_{s}(t)^{\alpha}} dt \\ &\leq \left( \int_{s}^{\bar{M}} \psi(t)^{\beta m} \eta_{s}(t)^{\alpha m} dt \right)^{\frac{1}{m}} \left( \int_{s}^{\bar{M}} \frac{dt}{\psi(t)^{\beta m'} \eta_{s}(t)^{\alpha m'}} \right)^{\frac{1}{m'}} \\ &\leq \left( \int_{s}^{\bar{M}} \psi(t)^{\frac{n-3}{n-1}} \eta_{s}(t)^{2} dt \right)^{\frac{1}{n-2}} \left( \sqrt{\frac{2}{C}} \int_{s}^{\bar{M}} \frac{\dot{\eta}_{s}(t)}{\eta_{s}(t)^{m'\alpha+1}} dt \right)^{\frac{n-3}{n-2}} \\ &\leq \left( \frac{2}{Cs^{2}} \int_{\{u \leq s\}} |\nabla u|^{p+2} dx \right)^{\frac{1}{n-2}} \left( \sqrt{\frac{2}{C}} \frac{n-3}{2} \right)^{\frac{n-3}{n-2}} \end{split}$$

which is exactly (1.11) (note that n-2 = p and  $\eta(\overline{M}) \ge 1$ ).

#### 4.3 Regularity of the extremal solution. Proof of Theorem 1.4

In this subsection we will prove the *a priori* estimates for minimal and extremal solutions of  $(1.15)_{\lambda}$  stated in Theorem 1.4. Let us remark that in the proof of Theorem 1.4 we will assume the nonlinearity f to be smooth. However, if it is only  $C^1$  we can proceed with an approximation argument as in the proof of Theorem 1.2 in [3].

The  $W^{1,p}$ -estimate established in Theorem 1.4 has as main ingredient the following result.

**Lemma 4.4.** Let f be an increasing positive  $C^1$  function satisfying (1.16) and  $\lambda \in (0, \lambda^*)$ . Let  $u = u_{\lambda} \in C_0^1(\overline{\Omega})$  be the minimal solution of  $(1.15)_{\lambda}$ . The following inequality holds:

$$\int_{\Omega} |\nabla u|^p \, dx \le \left( \max_{x \in \overline{\Omega}} |x| \right) \frac{1}{p'} \int_{\partial \Omega} |\nabla u|^p \, d\sigma. \tag{4.14}$$

*Proof.* Let  $G'(t) = g(t) = \lambda f(t)$ . First, we note that

$$x \cdot \nabla u \ g(u) = x \cdot \nabla G(u) = \operatorname{div} \left( G(u) x \right) - n G(u)$$

and that almost everywhere on  $\Omega$  we can evaluate

$$\begin{aligned} x \cdot \nabla u \ \Delta_p u - \operatorname{div} & \left( x \cdot \nabla u \ |\nabla u|^{p-2} \nabla u \right) &= -|\nabla u|^{p-2} \nabla u \cdot \nabla (x \cdot \nabla u) \\ &= -|\nabla u|^p - \frac{1}{p} \nabla |\nabla u|^p \cdot x \\ &= \frac{n-p}{p} |\nabla u|^p - \frac{1}{p} \operatorname{div} \left( |\nabla u|^p x \right). \end{aligned}$$

As a consequence, multiplying  $(1.15)_{\lambda}$  by  $x \cdot \nabla u$  and integrating on  $\Omega$ , we have

$$n\int_{\Omega} G(u) \, dx - \frac{n-p}{p} \int_{\Omega} |\nabla u|^p \, dx = \frac{1}{p'} \int_{\partial\Omega} |\nabla u|^p \, x \cdot \nu \, d\sigma, \tag{4.15}$$

where  $\nu$  is the outward unit normal to  $\Omega$ .

Noting that u is an absolute minimizer of the energy functional

$$J(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p \, dx - \int_{\Omega} G(u) \, dx$$

in the convex set  $\{v \in W_0^{1,p}(\Omega) : 0 \le v \le u\}$  (see [5]), we have that  $J(u) \le J(0) = 0$ . Therefore, from (4.15) we obtain

$$\int_{\Omega} |\nabla u|^p \, dx = nJ(u) + \frac{1}{p'} \int_{\partial \Omega} |\nabla u|^p \, x \cdot \nu \, d\sigma \le \left( \max_{x \in \overline{\Omega}} |x| \right) \frac{1}{p'} \int_{\partial \Omega} |\nabla u|^p \, d\sigma$$

proving the lemma.

Finally, we prove Theorem 1.4 (using the semistability condition (4.4) with an appropriate test function), Theorem 1.2, and Lemma 4.4.

Proof of Theorem 1.4. Let  $u_{\lambda}$  be the minimal solution of  $(1.15)_{\lambda}$  for  $\lambda \in (0, \lambda^*)$ . From [5] we know that minimal solutions are semi-stable. In particular,  $u_{\lambda}$  satisfies the semistability condition (4.4) for all  $\lambda \in (0, \lambda^*)$ .

Assume that  $\Omega$  is strictly convex. Let  $\delta(x) := \operatorname{dist}(x, \partial \Omega)$  be the distance to the boundary and  $\Omega_{\varepsilon} := \{x \in \Omega : \delta(x) < \varepsilon\}$ . By Proposition 3.1 there exist positive constants  $\varepsilon$ and  $\gamma$  such that for every  $x_0 \in \Omega_{\varepsilon}$  there exists a set  $I_{x_0} \subset \Omega$  satisfying  $|I_{x_0}| > \gamma$  and

$$u_{\lambda}(x_0)^{p-1} \le u_{\lambda}(y)^{p-1}$$
 for all  $y \in I_{x_0}$ . (4.16)

Let  $x_{\varepsilon} \in \overline{\Omega}_{\varepsilon}$  be such that  $u_{\lambda}(x_{\varepsilon}) = ||u_{\lambda}||_{L^{\infty}(\Omega_{\varepsilon})}$ . Integrating with respect to y in  $I_{x_{\varepsilon}}$  inequality (4.16) and using (1.16), we obtain

$$\|u_{\lambda}\|_{L^{\infty}(\Omega_{\varepsilon})}^{p-1} \leq \frac{1}{\gamma} \int_{I_{x_{\varepsilon}}} u_{\lambda}^{p-1} dy \leq \frac{1}{\gamma} \int_{\Omega} u_{\lambda}^{p-1} dy \leq \frac{C}{\gamma} \|f(u_{\lambda})\|_{L^{1}(\Omega)},$$
(4.17)

where C, here and in the rest of the proof, is a constant independent of  $\lambda$ . Letting  $s = \left(\frac{C}{\gamma} \|f(u_{\lambda})\|_{L^{1}(\Omega)}\right)^{1/(p-1)}$ , we deduce

$$\Omega_{\varepsilon} \subset \{ x \in \Omega : u_{\lambda}(x) \le s \}.$$
(4.18)

Now, choose

$$\eta(x) := \left\{ \begin{array}{ll} \delta(x) & \text{if} \quad \delta(x) < \varepsilon \\ \varepsilon & \text{if} \quad \delta(x) \ge \varepsilon \end{array} \right.$$

as a test function in (4.4) and use (4.18) to obtain

$$\varepsilon^2 \int_{\{u_{\lambda}>s\}} \left(\frac{4}{p^2} |\nabla_{T,u_{\lambda}}| \nabla u_{\lambda}|^{p/2}|^2 + \frac{n-1}{p-1} H^2_{u_{\lambda}} |\nabla u_{\lambda}|^p\right) \, dx \le \int_{\{u_{\lambda}\le s\}} |\nabla u_{\lambda}|^p \, dx.$$

Multiplying equation  $(1.15)_{\lambda}$  by  $T_s u_{\lambda} = \min\{s, u_{\lambda}\}$  we have

$$\int_{\{u_{\lambda} < s\}} |\nabla u_{\lambda}|^p \, dx = \lambda \int_{\Omega} f(u_{\lambda}) T_s u \, dx \le \lambda^* s \|f(u_{\lambda})\|_{L^1(\Omega)} = C \|f(u_{\lambda})\|_{L^1(\Omega)}^{p'}.$$
(4.19)

Combining the previous two inequalities we obtain

$$\int_{\{u_{\lambda}>s\}} \left(\frac{4}{p^2} |\nabla_{T,u_{\lambda}}| \nabla u_{\lambda}|^{p/2}|^2 + \frac{n-1}{p-1} H^2_{u_{\lambda}} |\nabla u_{\lambda}|^p\right) \, dx \le C \|f(u_{\lambda})\|^{p'}_{L^1(\Omega)}.$$

At this point, proceeding exactly as in the proof of Theorem 1.3, we conclude the  $L^r$  estimates established in parts (a) and (b).

In order to prove the  $W^{1,p}$ -estimate of part (b), recall that by (4.15) we have

$$\int_{\Omega} |\nabla u_{\lambda}|^p \, dx \le C \int_{\partial \Omega} |\nabla u_{\lambda}|^p \, d\sigma.$$

Therefore, we need to control the right hand side of the previous inequality. Since the nonlinearity f is increasing by hypothesis we obtain

$$f(u_{\lambda}) \le f\left(C\|f(u_{\lambda})\|_{L^{1}(\Omega)}^{\frac{1}{p-1}}\right) \quad \text{in } \Omega_{\varepsilon}$$

by (4.17), where C is a constant independent of  $\lambda$ .

Now, since  $-\Delta_p u_{\lambda} = \lambda f(u_{\lambda}) \in L^{\infty}(\Omega_{\varepsilon})$  in  $\Omega_{\varepsilon}$ , it holds

$$\|u_{\lambda}\|_{C^{1,\beta}(\overline{\Omega}_{\varepsilon})} \le C'$$

for some  $\beta \in (0,1)$  by [11], where C' is a constant depending only on  $n, p, \Omega, f$ , and  $||f(u_{\lambda})||_{L^{1}(\Omega)}$  proving the assertion.

Finally, assume that  $p \ge 2$  and (1.17) holds. From [15] we know that  $f(u^*) \in L^r(\Omega)$ for all  $1 \le r < n/(n - p')$ . In particular,  $f(u^*) \in L^1(\Omega)$ . Therefore, parts (i) and (ii) follow directly from (a) and (b).

Acknowledgments. The authors were supported by grant 2009SGR345(Catalunya) and MTM2011-27739-C04 (Spain). The second author was also supported by grant MTM2008-06349-C03-01 (Spain).

### References

- [1] Allard, W.K.: On the first variation of a varifold. Ann. Math. 95, 417–491 (1972)
- [2] Bénilan, Ph., Boccardo, L., Gallouët, T., Gariepy, R., Pierre, M., Vázquez, J.L.: An L<sup>1</sup>-theory of existence and uniqueness of solutions of nonlinear elliptic equations. Ann. Sc. Norm. Super. Pisa, Cl. Sci. 22, 241–273 (1995)
- [3] Cabré, X.: Regularity of minimizers of semilinear elliptic problems up to dimension 4. Comm. Pure Appl. Math **63**, 1362–1380 (2010)
- [4] Cabré, X., Capella, A., Sanchón, M.: Regularity of radial minimizers of reaction equations involving the *p*-Laplacian. Calc. Var. PDE 34, 475–494 (2009)
- [5] Cabré, X., Sanchón, M.: Semi-stable and extremal solutions of reaction equations involving the *p*-Laplacian. Comm. Pure Appl. Anal. 6, 43–67 (2007)
- [6] Cabré, X., Sanchón, M.: Geometric-type Hardy-Sobolev inequalities and applications to the regularity of minimizers. Preprint: arXiv:1111.2801v1
- [7] Canino, A., Le, P., Sciunzi, B.: Local  $W_{loc}^{2,m(x)}$  regularity for p(x)-Laplace equations. Preprint
- [8] DiBenedetto, E.:  $C^{1+\alpha}$  local regularity of weak solutions of degenerate elliptic equations. Nonlinear Anal. **7**, 827–850 (1983)

- [9] Damascelli, L., Sciunzi, B.: Regularity, monotonicity and symmetry of positive solutions of *m*-Laplace equations. J. Differential Equations **206**, 483–515 (2004)
- [10] Farina, A., Sciunzi, B., Valdinoci, E.: Bernstein and De Giorgi type problems: new results via a geometric approach. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 7, 741–791 (2008)
- [11] Lieberman, G.M.: Boundary regularity for solutions of degenerate elliptic equations. Nonlinear Anal. 12, 1203–1219 (1988)
- [12] Michael, J.H., Simon, L.: Sobolev and meanvalue inequalities on generalized submanifolds of R<sup>n</sup>. Comm. Pure Appl. Math. 26, 361–379 (1973)
- [13] Nedev, G.: Regularity of the extremal solution of semilinear elliptic equations. C. R. Acad. Sci. Paris Sér. I Math., 330, 997–1002 (2000)
- [14] Nedev, G.: Extremal solution of semilinear elliptic equations. Preprint 2001.
- [15] Sanchón, M.: Existence and regularity of the extremal solution of some nonlinear elliptic problems related to the *p*-Laplacian. Potential Anal. **27**, 217–224 (2007)
- [16] Tolksdorf, P.: Regularity for a more general class of quasilinear elliptic equations. J. Differential Equations 51, 126–150 (1984)
- [17] Trudinger, N.S.: Isoperimetric inequalities for quermassintegrals. Ann. Inst. Henri Poincaré 11, 411–425 (1994)
- [18] Trudinger, N.S.: On new isoperimetric inequalities and symmetrization. J. reine angew. Math. 488, 203–220 (1997)
- [19] Vázquez, J.L.: A strong maximum principle for some quasilinear elliptic equations. Appl. Math. Optim. 12, 191–202 (1984)