

THE RIESZ TRANSFORM AND QUANTITATIVE RECTIFIABILITY FOR GENERAL RADON MEASURES

DANIEL GIRELA-SARRIÓN AND XAVIER TOLSA

ABSTRACT. In this paper we show that if μ is a Borel measure in \mathbb{R}^{n+1} with growth of order n , so that the n -dimensional Riesz transform \mathcal{R}_μ is bounded in $L^2(\mu)$, and $B \subset \mathbb{R}^{n+1}$ is a ball with $\mu(B) \approx r(B)^n$ such that:

- (a) there is some n -plane L passing through the center of B such that for some $\delta > 0$ small enough, it holds

$$\int_B \frac{\text{dist}(x, L)}{r(B)} d\mu(x) \leq \delta \mu(B),$$

- (b) for some constant $\varepsilon > 0$ small enough,

$$\int_B |\mathcal{R}_\mu 1(x) - m_{\mu, B}(\mathcal{R}_\mu 1)|^2 d\mu(x) \leq \varepsilon \mu(B),$$

where $m_{\mu, B}(\mathcal{R}_\mu 1)$ stands for the mean of $\mathcal{R}_\mu 1$ on B with respect to μ , then there exists a uniformly n -rectifiable set Γ , with $\mu(\Gamma \cap B) \gtrsim \mu(B)$, and so that $\mu|_\Gamma$ is absolutely continuous with respect to $\mathcal{H}^n|_\Gamma$. This result is an essential tool to solve an old question on a two phase problem for harmonic measure in a subsequent paper by Azzam, Mouroglou and Tolsa.

CONTENTS

1. Introduction	2
2. Preliminaries	5
3. The Main Lemma	6
3.1. Preliminaries and statement of the Main Lemma	6
3.2. Reduction of Theorem 1.1 to the Main Lemma	6
3.3. The proof of Lemma 3.2	7
4. The Localization Lemma	12
5. The dyadic lattice of David and Mattila	16
6. The low density cells and the stopping cells	18
7. The periodic measure $\tilde{\mu}$	21
8. The approximating measure η	28
9. Proof of the Key Lemma by contradiction	40
9.1. A variational argument and an almost everywhere inequality	40
9.2. A maximum principle.	43
9.3. The contradiction	45
10. Construction of the AD-regular measure ζ and the uniformly rectifiable set Γ in the Main Lemma	47

2010 *Mathematics Subject Classification.* 42B20, 28A75, 28A78, 49Q20.

The authors were supported by the ERC grant 320501 of the European Research Council (FP7/2007-2013), and also partially supported by 2014-SGR-75 (Catalonia) and MTM2013-44304-P (Spain). D.G. was also supported by the grant BES-2011-045708 (Spain) and X.T. by the Marie Curie ITN MAnET (FP7-607647).

1. INTRODUCTION

In the work [NToV1] it was shown that, given an n -AD-regular measure in \mathbb{R}^{n+1} , the $L^2(\mu)$ boundedness of the n -dimensional Riesz transform implies the uniform n -rectifiability of μ . In the codimension 1 case, this result solved a long standing problem raised by David and Semmes [DS1]. In the present paper we obtain a related quantitative result valid for general Radon measures in \mathbb{R}^{n+1} with growth of order n . Our result turns out to be an essential tool for the solution of an old question on harmonic measure which will appear in a subsequent work by Azzam, Mouroglou and the second author of this paper [AMT].

To state our main theorem in detail we need to introduce some notation and terminology. Let μ be a Radon measure in \mathbb{R}^{n+1} . We say that μ has growth of order n (with constant C_0) if

$$\mu(B(x, r)) \leq C_0 r^n \quad \text{for all } x \in \mathbb{R}^{n+1} \text{ and all } r > 0.$$

A measure μ is called n -AD-regular (or just AD-regular or Ahlfors-David regular) if there exists some constant $C > 0$ such that

$$C^{-1} r^n \leq \mu(B(x, r)) \leq C r^n \quad \text{for all } x \in \text{supp}(\mu) \text{ and } 0 < r \leq \text{diam}(\text{supp}(\mu)).$$

Given a signed Radon measure ν in \mathbb{R}^{n+1} we consider the n -dimensional Riesz transform

$$\mathcal{R}\nu(x) = \int \frac{x - y}{|x - y|^{n+1}} d\nu(y),$$

whenever the integral makes sense. For $\varepsilon > 0$, its ε -truncated version is given by

$$\mathcal{R}_\varepsilon \nu(x) = \int_{|x-y|>\varepsilon} \frac{x - y}{|x - y|^{n+1}} d\nu(y).$$

For a positive Radon measure μ and a function $f \in L^1_{loc}(\mu)$, we denote $\mathcal{R}_\mu f \equiv \mathcal{R}(f\mu)$ and $\mathcal{R}_{\mu,\varepsilon} f \equiv \mathcal{R}_\varepsilon(f\mu)$. We say that the Riesz transform \mathcal{R}_μ is bounded in $L^2(\mu)$ if the truncated operators $\mathcal{R}_{\mu,\varepsilon} : L^2(\mu) \rightarrow L^2(\mu)$ are bounded uniformly on $\varepsilon > 0$.

If μ is a measure with no point masses such that \mathcal{R}_μ is bounded in $L^2(\mu)$ and ν is a finite Radon measure, the principal value

$$\text{pv}\mathcal{R}\nu(x) = \lim_{\varepsilon \rightarrow 0} \mathcal{R}_\varepsilon \nu(x)$$

exists μ -a.e. This follows easily from the results of [NToV2], arguing as in [To2, Chapter 8] with the Cauchy transform replaced by the Riesz transform. Abusing notation, we also write $\mathcal{R}\nu(x)$ instead of $\text{pv}\mathcal{R}\nu(x)$.

For $f \in L^1_{loc}(\mu)$ and $A \subset \mathbb{R}^{n+1}$ with $\mu(A) > 0$, we consider the μ -mean

$$m_{\mu,A}(f) = \int_A f d\mu = \frac{1}{\mu(A)} \int_A f d\mu.$$

Also, given a ball $B \subset \mathbb{R}^{n+1}$ and an n -plane L in \mathbb{R}^{n+1} , we denote

$$\beta_{\mu,1}^L(B) = \frac{1}{r(B)^n} \int_B \frac{\text{dist}(x, L)}{r(B)} d\mu(x),$$

where $r(B)$ stands for the radius of B . In a sense, this coefficient measures how close the points from $\text{supp } \mu$ are to the n -plane L in the ball B .

We also set

$$\Theta_\mu(B) = \frac{\mu(B)}{r(B)^n}, \quad P_\mu(B) = \sum_{j \geq 0} 2^{-j} \Theta_\mu(2^j B).$$

So $\Theta_\mu(B)$ is the n -dimensional density of μ on B and $P_\mu(B)$ is some kind of smoothened version of this density.

Our main theorem is the following.

Theorem 1.1. *Let μ be a Radon measure on \mathbb{R}^{n+1} and $B \subset \mathbb{R}^{n+1}$ a ball so that the following conditions hold:*

- (a) *For some constant $C_0 > 0$, $C_0^{-1} r(B)^n \leq \mu(B) \leq C_0 r(B)^n$.*
- (b) *$P_\mu(B) \leq C_0$, and $\mu(B(x, r)) \leq C_0 r^n$ for all $x \in B$ and $0 < r \leq r(B)$.*
- (c) *There is some n -plane L passing through the center of B such that for some $0 < \delta \ll 1$, it holds $\beta_{\mu,1}^L(B) \leq \delta$.*
- (d) *$\mathcal{R}_{\mu|_B}$ is bounded in $L^2(\mu|_B)$ with $\|\mathcal{R}_{\mu|_B}\|_{L^2(\mu|_B) \rightarrow L^2(\mu|_B)} \leq C_1$.*
- (e) *For some constant $0 < \varepsilon \ll 1$,*

$$\int_B |\mathcal{R}\mu(x) - m_{\mu,B}(\mathcal{R}\mu)|^2 d\mu(x) \leq \varepsilon \mu(B).$$

Then there exists some constant $\tau > 0$ such that if δ, ε are small enough (depending on C_0 and C_1), then there is a uniformly n -rectifiable set $\Gamma \subset \mathbb{R}^{n+1}$ such that

$$\mu(B \cap \Gamma) \geq \tau \mu(B).$$

The constant τ and the UR constants of Γ depend on all the constants above.

Some remarks are in order. First, we mention that the notion of uniform n -rectifiability will be introduced in Section 2. For the moment, for the reader's convenience, let us say that this a quantitative (and stronger) version of the notion of n -rectifiability. The UR constants are just the constants involved in the definition of uniform n -rectifiability. We also remark that it is immediate to check that the condition (b) above holds, for example, if μ has growth of order n (with constant $\frac{1}{2}C_0$). The statement in (b) which involves $P_\mu(B)$ is somewhat more general and it is more convenient for applications. Finally, we warn the reader that in the case that μ is not a finite measure, the statement (e) should be understood in the BMO sense. The fact that $P_\mu(B) < \infty$ by the assumption (b) guarantees that $\mathcal{R}\mu(x) - m_{\mu,B}(\mathcal{R}\mu)$ is correctly defined.

Note that, in particular, the theorem above ensures the existence of some piece of positive μ -measure of B where μ and the Hausdorff measure \mathcal{H}^n are mutually absolutely continuous on some subset of Γ . This fact, which at first sight may appear rather surprising, is one of the main difficulties for the proof of this result.

It is worth comparing Theorem 1.1 to Léger's theorem on Menger curvature. Given three points $x, y, z \in \mathbb{R}^2$, their Menger curvature is

$$c(x, y, z) = \frac{1}{R(x, y, z)},$$

where $R(x, y, z)$ is the radius of the circumference passing through x, y, z if they are pairwise different, and $c(x, y, z) = 0$ otherwise. The curvature of μ is defined by

$$c^2(\mu) = \iiint c(x, y, z)^2 d\mu(x) d\mu(y) d\mu(z).$$

This notion was first introduced by Melnikov [Me] when studying analytic capacity and, modulo an “error term”, is comparable to the squared $L^2(\mu)$ norm of the Cauchy transform of μ (see [MV]). One of the main ingredients of the proof of Vitushkin’s conjecture for removable singularities for bounded analytic functions by Guy David [Da] is Léger’s theorem [Lé] (sometimes called also David-Léger theorem). The quantitative version of this theorem asserts that if μ is a Radon measure in \mathbb{R}^2 with growth of order 1 and B a ball such that $\mu(B) \approx r(B)$, and further $c^2(\mu|_B) \leq \varepsilon \mu(B)$ for some $\varepsilon > 0$ small enough, then there exists some (possibly rotated) Lipschitz graph $\Gamma \subset \mathbb{R}^2$ such that $\mu(B \cap \Gamma) \geq \frac{9}{10} \mu(B)$. In particular, as in Theorem 1.1, it follows that a big piece of $\mu|_B$ is mutually absolutely continuous with respect to \mathcal{H}^1 on some subset of Γ . In a sense, one can think that Theorem 1.1 is an analogue for Riesz transforms of the quantitative Léger theorem for Menger curvature. Indeed, the role of the assumption (e) in Theorem 1.1 is played by the condition $c^2(\mu|_B) \leq \varepsilon \mu(B)$. Further, it is not difficult to check that this condition implies that there exists some line L such that $\beta_{\mu,1}^L(B) \leq \delta \mu(B)$, with $\delta = \delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$, analogously to the assumption (c) of Theorem 1.1.

On the other hand, from the theorem of Léger described above, it follows easily that if $\mathcal{H}^1(E) < \infty$ and $c^2(\mathcal{H}^1|_E) < \infty$, then E is 1-rectifiable. The analogous result in the codimension 1 case in \mathbb{R}^{n+1} (proved in [NToV2]) asserts if $E \subset \mathbb{R}^{n+1}$, $\mathcal{H}^n(E) < \infty$, $\mathcal{H}^n|_E$ has growth of order n , and $\|\mathcal{R}(\mathcal{H}^n|_E)\|_{L^2(\mathcal{H}^n|_E)} < \infty$, then E is n -rectifiable. However, as far as we know, this cannot be proved easily using Theorem 1.1.

The proof of Theorem 1.1 is substantially different from the one of Léger’s theorem. When estimating the $L^2(\mu)$ norm of $\mathcal{R}\mu$ we are dealing with a singular integral and there may be cancellations among different scales. So the situation is more delicate than in the case of the curvature $c^2(\mu)$, which is defined by a non-negative integrand (namely, the squared Menger curvature of three points).

To prove Theorem 1.1 we will apply some of the techniques developed in [ENV] and [NToV1]. In particular, by using a variational argument we will estimate from below the $L^2(\mu)$ norm of the Riesz transform of a suitable periodization of a smoothened version of the measure μ restricted to some appropriate cube Q_0 . The assumption that $\beta_{\mu,1}^L(B) \leq \delta$ in (c) is necessary for technical reasons, and we do not know if the theorem holds without this condition.

Finally we are going to describe the aforementioned result on harmonic measure from [AMT] whose proof uses Theorem 1.1 as an essential tool. For simplicity, we will only announce this for domains $\Omega_1, \Omega_2 \subset \mathbb{R}^{n+1}$ satisfying the condition

$$(1.1) \quad \mathcal{H}_\infty^s((\mathbb{R}^{n+1} \setminus \Omega_i) \cap B(x, r)) \approx r^s \quad \text{for all } x \in \partial\Omega_i \text{ and } 0 < r \leq r_0,$$

for some fixed $s \in (n-1, n+1]$, $r_0 > 0$, where \mathcal{H}_∞^s stands for the s -dimensional Hausdorff content. For example, the so-called NTA domains introduced by Jerison and Kenig [JK] satisfy this condition, and also the simply connected domains in the plane.

Theorem 1.2 (Azzam, Mouroglou, Tolsa). *Let $\Omega_1, \Omega_2 \subset \mathbb{R}^{n+1}$, $n \geq 2$, be two disjoint connected domains with $\partial\Omega^1 = \partial\Omega^2$ so that (1.1) holds. For $i = 1, 2$, let $\omega^i = \omega_{\Omega_i}^{x_i}$, $x_i \in \Omega_i$, be the respective harmonic measures with pole at x_i , and let $E \subset \partial\Omega_1$ be a Borel set. If $\omega^1 \ll \omega^2 \ll \omega^1$ on E , then E contains an n -rectifiable subset F with $\omega^1(E \setminus F) = \omega^2(E \setminus F) = 0$ where ω^1 and ω^2 are mutually absolutely continuous with respect to \mathcal{H}^n .*

Up to now this result was known only in the case when Ω_1, Ω_2 are planar domains, by results of Bishop, Carleson, Garnett and Jones [BCGJ] and Bishop [Bi1], and it was an open problem to extend

this to higher dimensions (see Conjecture 8 in [Bi2]). For a partial result in the higher dimensional case, see the nice work [KPT] by Kenig, Preiss, and Toro.

2. PRELIMINARIES

In this paper we will use the letters c, C to denote constants (quite often absolute constants, perhaps depending on n) which may change their values at different occurrences. On the other hand, constants with subscripts, such as C_1 , do not change their values at different occurrences.

We will write $a \lesssim b$ if there is $C > 0$ so that $a \leq Cb$ and $a \lesssim_t b$ if the constant C depends on the parameter t . We write $a \approx b$ to mean $a \lesssim b \lesssim a$ and define $a \approx_t b$ similarly.

We denote the open ball of radius r centered at x by $B(x, r)$. For a ball $B = B(x, r)$ and $a > 0$ we write $r(B)$ for its radius and $aB = B(x, ar)$. The notation $A(x, r_1, r_2)$ stands for an open annulus centered at x with inner radius r_1 and outer radius r_2 .

Given a cube Q , we denote by $\ell(Q)$ its side length. Unless otherwise stated, we assume that its sides are parallel to the coordinate axes of \mathbb{R}^{n+1} . By aQ we denote a cube concentric with Q with side length $a\ell(Q)$. We write

$$\Theta_\mu(Q) = \frac{\mu(Q)}{\ell(Q)^n}, \quad P_\mu(Q) = \sum_{j \geq 0} 2^{-j} \Theta_\mu(2^j Q).$$

We also consider the pointwise densities

$$\Theta_*^n(x, \mu) = \liminf_{r \rightarrow 0+} \frac{\mu(B(x, r))}{(2r)^n}, \quad \Theta^{n,*}(x, \mu) = \limsup_{r \rightarrow 0+} \frac{\mu(B(x, r))}{(2r)^n}.$$

A set $E \subset \mathbb{R}^d$ is called n -rectifiable if there are Lipschitz maps $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^d$, $i = 1, 2, \dots$, such that

$$(2.1) \quad \mathcal{H}^n \left(E \setminus \bigcup_i f_i(\mathbb{R}^n) \right) = 0,$$

where \mathcal{H}^n stands for the n -dimensional Hausdorff measure. Also, one says that a Radon measure μ on \mathbb{R}^d is n -rectifiable if μ vanishes out of an n -rectifiable set $E \subset \mathbb{R}^d$ and moreover μ is absolutely continuous with respect to $\mathcal{H}^n|_E$.

A measure μ in \mathbb{R}^d is uniformly n -rectifiable if it is n -AD-regular and there exist $\theta, M > 0$ such that for all $x \in \text{supp}(\mu)$ and all $r > 0$ there is a Lipschitz mapping g from the ball $B_n(0, r)$ in \mathbb{R}^n to \mathbb{R}^d with $\text{Lip}(g) \leq M$ such that

$$\mu(B(x, r) \cap g(B_n(0, r))) \geq \theta r^n.$$

In the case $n = 1$, it is known that μ is uniformly 1-rectifiable if and only if $\text{supp}(\mu)$ is contained in a rectifiable curve Γ in \mathbb{R}^d such that the arc length measure on Γ is 1-AD-regular.

A set $E \subset \mathbb{R}^d$ is called n -AD-regular if $\mathcal{H}^n|_E$ is n -AD-regular, and it is called uniformly n -rectifiable if $\mathcal{H}^n|_E$ is uniformly n -rectifiable.

3. THE MAIN LEMMA

3.1. Preliminaries and statement of the Main Lemma. Given two Radon measure μ and σ in \mathbb{R}^{n+1} and a cube $Q \subset \mathbb{R}^{n+1}$, we set

$$(3.1) \quad d_Q(\mu, \sigma) = \sup_f \int f d(\mu - \sigma),$$

where the supremum is taken over all the 1-Lipschitz functions supported on Q . Given an n -plane L in \mathbb{R}^{n+1} , then we denote

$$\alpha_\mu^L(Q) = \frac{1}{\ell(Q)^{n+1}} \inf_{c \geq 0} d_Q(\mu, c \mathcal{H}^n|_L).$$

Given $t > 0$, we say that Q has t -thin boundary if

$$\mu(\{x \in 2Q : \text{dist}(x, \partial Q) \leq \lambda \ell(Q)\}) \leq t \lambda \mu(2Q) \quad \text{for all } \lambda > 0.$$

It is well known that for any given cube $Q_0 \subset \mathbb{R}^{n+1}$ and $a > 1$, there exists another cube Q with t -thin boundary such that $Q_0 \subset Q \subset aQ_0$, with t depending just on n and a . For the proof, we refer the reader to Lemma 9.43 of [To2], for example.

Main Lemma 3.1. *Let $n \geq 1$ and let $C_0, C_1 > 0$ be some arbitrary constants. There exists $A = A(C_0, C_1, n) > 10$ big enough and $\varepsilon = \varepsilon(C_0, C_1, n) > 0$ small enough such that if $\delta = \delta(A, C_0, C_1, n) > 0$ is small enough, then the following holds. Let μ be a Radon measure in \mathbb{R}^{n+1} and $Q_0 \subset \mathbb{R}^{n+1}$ a cube centered at the origin satisfying the following properties:*

- (a) $\mu(Q_0) = \ell(Q_0)^n$.
- (b) $P_\mu(AQ_0) \leq C_0$.
- (c) For all $x \in AQ_0$ and $0 < r \leq A\ell(Q_0)$, $\Theta_\mu(B(x, r)) \leq C_0$.
- (d) Q_0 has C_0 -thin boundary.
- (e) $\alpha_\mu^H(3AQ_0) \leq \delta$, where $H = \{x \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$.
- (f) $\mathcal{R}_{\mu|_{2Q_0}}$ is bounded in $L^2(\mu|_{2Q_0})$ with $\|\mathcal{R}_{\mu|_{2Q_0}}\|_{L^2(\mu|_{2Q_0}) \rightarrow L^2(\mu|_{2Q_0})} \leq C_1$.
- (g) We have

$$\int_{Q_0} |\mathcal{R}\mu(x) - m_{\mu, Q_0}(\mathcal{R}\mu)|^2 d\mu(x) \leq \varepsilon \mu(Q_0).$$

Then there exists some constant $\tau > 0$ and a uniformly n -rectifiable set $\Gamma \subset \mathbb{R}^{n+1}$ such that

$$\mu(Q_0 \cap \Gamma) \geq \tau \mu(Q_0).$$

The constant τ and the UR constants of Γ depend on all the constants above.

Note that the condition (c) in the Main Lemma implies that $\mu(2Q_0) \lesssim C_0 \mu(Q_0)$.

3.2. Reduction of Theorem 1.1 to the Main Lemma. Assume that the Main Lemma is proved. Then in order to prove Theorem 1.1 it is enough to show the following.

Lemma 3.2. *Let μ and $B \subset \mathbb{R}^{n+1}$ satisfy the assumptions of Theorem 1.1 with constants C_0, C_1, δ , and ε . For all $A' \geq 10$ and all $\delta', \varepsilon' > 0$, if δ and ε are small enough, there exists a cube Q_0 satisfying:*

- (a) $3A'Q_0 \subset B$ and $\text{dist}(A'Q_0, \partial B) \geq C_0'^{-1} r(B)$, with C_0' depending only on C_0 and n .
- (b) For some constant $\gamma = \gamma(\delta') > 0$, $\gamma r(B) \leq \ell(Q_0) \leq A'^{-1} r(B)$.
- (c) $\mu(Q_0) \geq C_0'^{-1} \ell(Q_0)^n$.
- (d) Q_0 has C_0' -thin boundary.
- (e) $\alpha_\mu^L(3A'Q_0) \leq \delta'$, where L is some n -plane that passes through the center of Q_0 and is parallel to one of the faces.
- (f) $\int_{Q_0} |\mathcal{R}\mu(x) - m_{\mu, Q_0}(\mathcal{R}\mu)|^2 d\mu(x) \leq \varepsilon' \mu(Q_0)$.

Before proving this lemma, we show how this is used to reduce Theorem 1.1 to the Main Lemma.

Proof of Theorem 1.1 using Lemma 3.2 and the Main Lemma 3.1. Let $B \subset \mathbb{R}^{n+1}$ be some ball satisfying the assumptions of Theorem 1.1 with constants C_0, C_1, δ , and ε . Let Q_0 be the cube given by Lemma 3.2, for some constants $A' \geq 10$ and $\delta', \varepsilon' > 0$ to be fixed below.

We just have to check that the assumptions (a)-(g) of the Main Lemma are satisfied by the measure

$$\tilde{\mu} = \frac{\ell(Q_0)^n}{\mu(Q_0)} \mu$$

if A' is big enough and δ', ε' are small enough. Obviously, the assumption (a) from the Main Lemma is satisfied by $\tilde{\mu}$.

To show that (b) holds (with a constant different from C_0), note first that

$$(3.2) \quad \frac{\ell(Q_0)^n}{\mu(Q_0)} \approx 1,$$

with the implicit constant depending on C_0 and C_0' . Indeed, from the assumption (c) in Theorem 1.1, $\mu(Q_0) \lesssim C_0 \ell(Q_0)^n$, and by (c) in Lemma 3.2, $\mu(Q_0) \geq C_0'^{-1} \ell(Q_0)^n$. Then we have

$$P_{\tilde{\mu}}(A'Q_0) \lesssim P_\mu(A'Q_0) = \sum_{j \geq 0: 2^j A'Q_0 \subset B} 2^{-j} \Theta_\mu(2^j A'Q_0) + \sum_{j \geq 0: 2^j A'Q_0 \not\subset B} 2^{-j} \Theta_\mu(2^j A'Q_0).$$

The first sum on the right hand side does not exceed $C C_0'$ because $\Theta_\mu(2^j A'Q_0) \lesssim C_0$ for all cubes $2^j A'Q_0$ contained in B . Also, one can check that the last sum is bounded by $C P_\mu(B)$ because $\ell(2^j A'Q_0) \gtrsim r(B)$ for all the j 's in this sum, taking into account that $\text{dist}(A'Q_0, \partial B) \geq C_0'^{-1} r(B)$.

The assumptions (c)-(e) in the Main Lemma are obviously satisfied too because of (3.2) and the analogous conditions in Lemma 3.2, with somewhat different constants $C_0'', \delta'', \varepsilon''$ replacing C_0, δ, ε . \square

3.3. The proof of Lemma 3.2. Below we identify \mathbb{R}^n with the horizontal n -plane $H = \{x \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$. Then, given a Radon measure σ in \mathbb{R}^n and a cube $Q \subset \mathbb{R}^n$, we denote

$$(3.3) \quad \alpha_\sigma^{\mathbb{R}^n}(Q) = \frac{1}{\ell(Q)^{n+1}} \inf_{c \geq 0} d_Q(\sigma, c \mathcal{H}^n|_{\mathbb{R}^n}),$$

where d_Q is defined as in (3.1) (although now $Q \subset \mathbb{R}^n$ instead of $Q \subset \mathbb{R}^{n+1}$) and the infimum runs over all constants $c > 0$. Note that

$$\alpha_\sigma^{\mathbb{R}^n}(Q) \approx \alpha_\sigma^H(\widehat{Q}),$$

where $\widehat{Q} = Q \times [-\ell(Q)/2, \ell(Q)/2]$. This follows easily from the fact that any 1-Lipschitz function supported in Q can be extended to a C -Lipschitz function supported in \widehat{Q} , with $C \lesssim 1$.

We need a couple of auxiliary results. The first one is the following.

Lemma 3.3. *Suppose that σ is some finite Radon measure supported in \mathbb{R}^n such that $d\sigma(x) = \rho(x) d\mathcal{H}^n|_{\mathbb{R}^n}$, with $\|\rho\|_\infty < \infty$. Then, for every $R \in \mathcal{D}(\mathbb{R}^n)$ we have*

$$\sum_{Q \in \mathcal{D}(\mathbb{R}^n): Q \subset R} \alpha_\sigma^{\mathbb{R}^n}(3Q)^2 \ell(Q)^n \lesssim \|\rho\|_\infty^2 \ell(R)^n.$$

In this lemma, $\mathcal{D}(\mathbb{R}^n)$ stands for the family of the usual dyadic cubes in \mathbb{R}^n .

This lemma can be proved by arguments similar to the ones used in [To1] to show that the analogous estimate holds for Lipschitz graphs. Alternatively, it can be deduced as a corollary of that result for the case where the Lipschitz graph is just a horizontal n -plane, using the auxiliary AD-regular measure $\tilde{\sigma} = 2\|\rho\|_\infty \mathcal{H}^n|_{\mathbb{R}^n} + \sigma$ and taking into account that $\alpha_\sigma^{\mathbb{R}^n}(Q) = \alpha_{\tilde{\sigma}}^{\mathbb{R}^n}(Q)$ for any cube $Q \subset \mathbb{R}^n$.

The second auxiliary result we need is the next one.

Lemma 3.4. *Let σ be some finite Radon measure in \mathbb{R}^n and $R \in \mathcal{D}(\mathbb{R}^n)$ such that*

$$\sigma(Q) \leq C_2 \ell(Q)^n$$

for all the cubes $Q \in \mathcal{D}(\mathbb{R}^n)$ with $\ell(Q) \geq \ell_0$. Then, for every $R \in \mathcal{D}(\mathbb{R}^n)$ we have

$$\sum_{\substack{Q \in \mathcal{D}(\mathbb{R}^n): Q \subset R \\ \ell(Q) \geq \ell_0}} \alpha_\sigma^{\mathbb{R}^n}(3Q)^2 \ell(Q)^n \lesssim C_2^2 \ell(R)^n.$$

Proof. Let $\varphi(x) = m(B(0, \ell_0))^{-1} \chi_{B(0, \ell_0)}(x)$. Consider the function $\rho = \varphi * \sigma$ and the measure $\nu = \rho dx$. We have $\|\rho\|_\infty \lesssim C_2$, since for all $x \in \mathbb{R}^n$

$$\rho(x) = \frac{1}{m(B(0, \ell_0))} \int \varphi(x - y) d\sigma(y) = \frac{\sigma(B(x, \ell_0))}{m(B(x, \ell_0))} \lesssim C_2.$$

Let us see that

$$(3.4) \quad \text{dist}_{3Q}(\nu, \sigma) \lesssim C_2 \ell_0 \ell(Q)^n \quad \text{for any cube } Q \text{ with } \ell(Q) \geq \ell_0.$$

For any function 1-Lipschitz function f supported on $3Q$, we have

$$\left| \int f d\nu - \int f d\sigma \right| = \left| \int f (\varphi * \sigma) dx - \int f d\sigma \right| = \left| \int f * \varphi d\sigma - \int f d\sigma \right|.$$

Since f is 1-Lipschitz we have

$$|f(x) - f * \varphi(x)| = \left| \int_{y \in B(x, \ell_0)} (f(x) - f(y)) \varphi(x - y) dy \right| \leq \int \ell_0 \varphi(x - y) dy = \ell_0.$$

Thus,

$$\left| \int f d\nu - \int f d\sigma \right| \lesssim \ell_0 \sigma(6Q) \lesssim C_2 \ell_0 \ell(Q)^n,$$

since $\text{supp}(f) \cup \text{supp}(f * \varphi) \subset 6Q$, and so (3.4) holds.

From (3.4) we infer that

$$\alpha_\sigma^{\mathbb{R}^n}(3Q) \lesssim \alpha_\nu^{\mathbb{R}^n}(Q) + C_2 \frac{\ell_0}{\ell(Q)},$$

and by Lemma 3.3,

$$\sum_{\substack{Q \in \mathcal{D}(\mathbb{R}^n): Q \subset R \\ \ell(Q) \geq \ell_0}} \alpha_{\sigma}^{\mathbb{R}^n} (3Q)^2 \ell(Q)^n \lesssim \sum_{\substack{Q \in \mathcal{D}(\mathbb{R}^n): Q \subset R \\ \ell(Q) \geq \ell_0}} \left(\alpha_{\nu}^{\mathbb{R}^n} (3Q)^2 + C_2^2 \frac{\ell_0^2}{\ell(Q)^2} \right) \ell(Q)^n \lesssim C_2^2 \ell(R)^n.$$

□

Proof of Lemma 3.2. Let μ and B be as in Theorem 1.1. By a suitable translation and rotation we may assume that the n -plane L from Theorem 1.1 coincides with the horizontal n -plane $H = \{x \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$ and that $B = B(0, r_0)$.

Our first objective consists in finding an auxiliary cube R_0 contained in B , centered in H , and far from ∂B , so that $\mu(R_0) \approx \mu(B)$. The cube Q_0 , to be chosen later, will be an appropriate cube contained in R_0 . To find R_0 , for some constant $0 < d < 1/10$ to be fixed below, we consider a grid \mathcal{Q} of n -dimensional cubes with side length $2dr_0$ in H , so that they cover H and have disjoint interiors. We also consider the family of $(n+1)$ -dimensional cubes

$$\widehat{\mathcal{Q}} = \{Q \times [-dr_0, dr_0] : Q \in \mathcal{Q}\},$$

so that the union of the cubes from $\widehat{\mathcal{Q}}$ equals the strip

$$V = \{x \in \mathbb{R}^{n+1} : \text{dist}(x, H) \leq dr_0\}.$$

For any constant $0 < a < 1$ we have

$$\mu(B \setminus (aB \cap V)) \leq \sum_{\substack{P \in \widehat{\mathcal{Q}}: \\ P \cap (B \setminus aB) \neq \emptyset}} \mu(P \cap B) + \mu\left(B \setminus \bigcup_{P \in \widehat{\mathcal{Q}}} P\right) =: S_1 + S_2.$$

To bound S_1 we use the growth condition of order n of $\mu|_B$:

$$S_1 \lesssim_{C_0} \sum_{\substack{P \in \widehat{\mathcal{Q}}: \\ P \cap (B \setminus aB) \neq \emptyset}} \ell(P)^n \lesssim \mathcal{H}^n(H \cap A(0, (a - n^{1/2}2d)r_0, (1 + n^{1/2}2d)r_0)) \lesssim_{C_0} (d + 1 - a)r_0^n.$$

To estimate S_2 we use the fact that the points $x \in B \setminus \bigcup_{P \in \widehat{\mathcal{Q}}} P$ are at a distance from H larger than dr_0 and apply Chebyshev:

$$S_2 \leq \int_B \frac{\text{dist}(x, H)}{dr_0} d\mu(x) = \frac{1}{d} \beta_{\mu,1}^H(B) r_0^n.$$

So we obtain

$$\mu(B \setminus (aB \cap V)) \leq C(C_0) \left((d + 1 - a) + \frac{1}{d} \beta_{\mu,1}^H(B) \right) \mu(B).$$

We take d and a so that

$$10(n+1)^{1/2}d = (1-a) = \frac{1}{10C(C_0)},$$

and we assume

$$\beta_{\mu,1}^H(B) \leq \delta \leq \frac{d}{10C(C_0)} = \frac{1}{10(n+1)^{1/2}(10C(C_0))^2},$$

so that $\mu(B \setminus (aB \cap V)) \leq \frac{3}{10} \mu(B)$. Now we choose R_0 to be a cube from $\widehat{\mathcal{Q}}$ which intersects $aB \cap V$ and has maximal μ -measure. Obviously,

$$\mu(R_0) \approx_{C_0} \mu(aB \cap V) \approx_{C_0} \mu(B),$$

and since $\text{diam}(R_0) = 2(n+1)^{1/2} d r_0 = \frac{1-a}{5} r_0$, it follows that

$$(3.5) \quad \text{dist}(4R_0, \partial B) \approx_{C_0} (1-a)r_0 \approx_{C_0} r_0.$$

The cube Q_0 we are looking for will be an appropriate cube contained in R_0 . To find this, first we consider the thin strip

$$V_\delta = \{x \in \mathbb{R}^{n+1} : \text{dist}(x, H) \leq \delta^{1/2} r_0\}.$$

Observe that

$$(3.6) \quad \mu(B \setminus V_\delta) \leq \int_B \frac{\text{dist}(x, H)}{\delta^{1/2} r_0} d\mu(x) = \frac{\beta_{\mu,1}^H(B)}{\delta^{1/2}} r_0^n \lesssim_{C_0} \delta^{1/2} \mu(B).$$

Denote by Π the orthogonal projection on H and consider the measure $\sigma = \Pi_\#(\mu|_{V_\delta})$. Since V_δ has width $2\delta^{1/2}$, from the growth condition (b) in Theorem 1.1, it follows that $\sigma(Q) \lesssim_{C_0} \ell(Q)^n$ for any cube Q centered on H with $\ell(Q) \geq \delta^{1/2} r_0$. Assume without loss of generality that R_0 is a dyadic cube. Then, by Lemma 3.4,

$$\sum_{\substack{Q \in \mathcal{D}(\mathbb{R}^n, R_0) \\ \ell(Q) \geq \delta^{1/2} r_0}} \alpha_\sigma^{\mathbb{R}^n} (3Q)^2 \ell(Q)^n \lesssim_{C_0} \ell(R_0)^n,$$

where $\mathcal{D}(\mathbb{R}^n, R_0)$ stands for the family of dyadic cubes in \mathbb{R}^n contained in R_0 . From this inequality, it easily follows that, for any constant $A' > 10$,

$$\sum_{\substack{Q \in \mathcal{D}(\mathbb{R}^n, R_0) : \ell(A'Q) \leq \ell(R_0) \\ \ell(Q) \geq \delta^{1/(4n+1)} r_0}} \alpha_\sigma^{\mathbb{R}^n} (4A'Q)^2 \ell(Q)^n \lesssim_{C_0, A'} \ell(R_0)^n.$$

Note that we have used the fact that $\delta^{1/(4n+1)} > \delta^{1/2}$. Since the number of dyadic generations between the largest cubes $Q \in \mathcal{D}(\mathbb{R}^n)$ with $\ell(A'Q) \leq \ell(R_0)$ and the smallest ones with side length $\ell(Q) \geq \delta^{1/(4n+1)} r_0$ is comparable to

$$\log_2 \frac{C(A')\ell(R_0)}{\delta^{1/(4n+1)} r_0} \approx \log_2 \frac{C(A', C_0)}{\delta^{1/(4n+1)}},$$

we deduce that there exists some intermediate generation j such that

$$\sum_{Q \in \mathcal{D}_j(\mathbb{R}^n, R_0) : \ell(A'Q) \leq \ell(R_0)} \alpha_\sigma^{\mathbb{R}^n} (4A'Q)^2 \ell(Q)^n \lesssim_{C_0, A'} \frac{1}{\log_2 \frac{C(A', C_0)}{\delta^{1/(4n+1)}}} \ell(R_0)^n.$$

Thus, for any $\delta' > 0$, if δ is small enough, we derive

$$\sum_{\substack{Q \in \mathcal{D}_j(\mathbb{R}^n, R_0) : \\ \ell(A'Q) \leq \ell(R_0)}} \alpha_\sigma^{\mathbb{R}^n} (4A'Q)^2 \sigma(Q) \leq C(C_0) \sum_{\substack{Q \in \mathcal{D}_j(\mathbb{R}^n, R_0) : \\ \ell(A'Q) \leq \ell(R_0)}} \alpha_\sigma^{\mathbb{R}^n} (4A'Q)^2 \ell(Q)^n \leq \frac{\delta'^2}{50} \sigma(R_0).$$

Denote by \mathcal{G} the subfamily of cubes from $\mathcal{D}_j(\mathbb{R}^n, R_0)$ such that $\Theta_\sigma(Q) \geq \frac{1}{2} \Theta_\sigma(R_0)$. Observe that

$$\sum_{Q \in \mathcal{D}_j(\mathbb{R}^n, R_0) \setminus \mathcal{G}} \sigma(Q) \leq \frac{1}{2} \Theta_\sigma(R_0) \sum_{Q \in \mathcal{D}_j(\mathbb{R}^n, R_0) \setminus \mathcal{G}} \ell(Q)^n \leq \frac{1}{2} \Theta_\sigma(R_0) \ell(R_0)^n = \frac{1}{2} \sigma(R_0).$$

Hence, $\sum_{Q \in \mathcal{G}} \sigma(Q) \geq \frac{1}{2} \sigma(R_0)$, and so

$$\sum_{Q \in \mathcal{G}} \alpha_{\sigma}^{\mathbb{R}^n} (4A'Q)^2 \sigma(Q) \leq \frac{\delta'^2}{50} \sigma(R_0) \leq \frac{\delta'^2}{25} \sigma\left(\bigcup_{Q \in \mathcal{G}} Q\right).$$

Then we deduce that there exists some cube $Q \in \mathcal{G}$ such that $\alpha_{\sigma}^{\mathbb{R}^n} (4A'Q) \leq \frac{\delta'}{5}$.

Denote $\widehat{Q} = Q \times [-\ell(Q)/2, \ell(Q)/2]$. We wish now to bound $\alpha_{\mu}^H (4A'\widehat{Q})$ in terms of $\alpha_{\sigma}^{\mathbb{R}^n} (4A'Q)$. Let c_H be the constant that minimizes the infimum in the definition of $\alpha_{\sigma}^{\mathbb{R}^n} (4A'Q)$ in (3.3). Given any 1-Lipschitz function f supported on $4A'\widehat{Q}$ we have

$$\begin{aligned} \left| \int f d(\mu - c_H \mathcal{H}^n|_H) \right| &\leq \int_{4A'\widehat{Q} \setminus V_{\delta}} |f| d\mu + \left| \int f d(\mu|_{V_{\delta}} - \sigma) \right| + \left| \int f d(\sigma - c_H \mathcal{H}^n|_H) \right| \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

By (3.6), using also that $\ell(Q) \geq \delta^{1/(4n+4)} r_0$, we have

$$I_1 \leq \|f\|_{\infty} \mu(B \setminus V_{\delta}) \lesssim_{C_0} \delta^{1/2} \ell(4A'Q) \mu(B) \lesssim_{C_0, A'} \delta^{1/2} \ell(Q) r_0^n \lesssim_{C_0, A'} \delta^{1/4} \ell(Q)^{n+1}.$$

Now we deal with I_2 . By the definition of σ and the Lipschitz condition on f , we get:

$$\begin{aligned} I_2 &= \left| \int_{4A'\widehat{Q}} f(x) - f(\Pi(x)) d\mu|_{V_{\delta}}(x) \right| \\ &\leq \int_{4A'\widehat{Q}} \text{dist}(x, H) d\mu|_{V_{\delta}}(x) \leq \beta_{\mu, 1}(B) r_0^{n+1} \leq \delta r_0^{n+1} \leq \delta^{3/4} \ell(Q)^{n+1}. \end{aligned}$$

Finally, concerning I_3 , we have

$$I_3 \leq \alpha_{\sigma}^{\mathbb{R}^n} (4A'Q) \ell(4A'Q)^{n+1} \leq \frac{\delta'}{5} \ell(4A'Q)^{n+1}.$$

Gathering the estimates obtained for I_1, I_2, I_3 and choosing δ small enough we obtain

$$\left| \int f d(\mu - c_H \mathcal{H}^n|_H) \right| \leq \frac{\delta'}{2} \ell(4A'Q)^{n+1},$$

and thus $\alpha_{\mu}^H (4A'\widehat{Q}) \leq \frac{\delta'}{2}$.

Finally, we choose Q_0 to be a cube with thin boundary such that $\widehat{Q} \subset Q_0 \subset 1.1\widehat{Q}$. Since $3A'Q_0 \subset 4A'\widehat{Q} \subset 4R_0$ and $\ell(3A'Q_0) \approx \ell(4A'\widehat{Q})$, we deduce that $3A'Q_0 \subset B$ (by (3.5)) and that $\alpha_{\mu}^H (3A'Q_0) \lesssim \alpha_{\mu}^H (4A'\widehat{Q}) \lesssim \delta'$. Then it is easy to check that Q_0 satisfies all the properties (a)-(e) by construction. Regarding (f), we have

$$\begin{aligned} \int_{Q_0} |\mathcal{R}\mu(x) - m_{\mu, Q_0}(\mathcal{R}\mu)|^2 d\mu(x) &\leq 2 \int_{Q_0} |\mathcal{R}\mu(x) - m_{\mu, B}(\mathcal{R}\mu)|^2 d\mu(x) \\ &\leq 2\varepsilon \mu(B) \approx_{C_0, \delta} \varepsilon \mu(Q_0). \end{aligned}$$

Thus if ε is small enough, (f) holds. \square

4. THE LOCALIZATION LEMMA

This and the remaining sections of this paper are devoted to the proof of the Main Lemma 3.1. We assume that the hypotheses of the Main Lemma 3.1 hold. From now on, we allow all the constants denoted by C and all the implicit constants in the relations “ \lesssim ” and “ \approx ” to depend on the constants C_0 and C_1 in the Main Lemma (but not on A , δ or ε).

Recall that H stands for the horizontal hyperplane $\{x \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$. Also we let c_H be some constant that minimizes the infimum in the definition of $\alpha^H(3AQ_0)$ and we denote $\mathcal{L}_H = c_H \mathcal{H}^n|_H$.

Lemma 4.1. *If δ is small enough (depending on A), then we have $c_H \approx 1$ and $\mu(AQ_0) \lesssim A^n \mu(Q_0)$.*

Proof. Let φ be a non-negative C^1 function supported on $2Q_0$ which equals 1 on Q_0 and satisfies $\|\nabla \varphi\|_\infty \lesssim 1/\ell(Q_0)$. Then we have

$$(4.1) \quad \left| \int \varphi d(\mu - \mathcal{L}_H) \right| \leq \|\nabla \varphi\|_\infty \ell(3AQ_0)^{n+1} \alpha_\mu^H(3AQ_0) \lesssim A^{n+1} \delta \ell(Q_0)^n.$$

Note that the left hand side above equals

$$\left| \int \varphi d\mu - c_H \int_H \varphi d\mathcal{H}^n \right| = |c_1 - c_H| \int_H \varphi d\mathcal{H}^n,$$

with

$$c_1 = \frac{\int \varphi d\mu}{\int_H \varphi d\mathcal{H}^n} \approx 1,$$

taking into account that

$$\mu(Q_0) \leq \int \varphi d\mu \leq \mu(2Q_0) \lesssim C_0 \ell(2Q_0)^n \lesssim C_0 \mu(Q_0).$$

So from (4.1) we deduce that

$$|c_1 - c_H| \lesssim A^{n+1} \delta \frac{\ell(Q_0)^n}{\int_H \varphi d\mathcal{H}^n} \lesssim A^{n+1} \delta.$$

The right hand side is $\ll 1 \approx c_1$ if δ is small enough (depending on A), and so we infer that

$$c_H \approx c_1 \approx 1.$$

The estimate $\mu(AQ_0) \lesssim A^n \ell(Q_0)^n$ is an immediate consequence of the assumptions either (b) or (c) in the Main Lemma 3.1. \square

Lemma 4.2 (Localization Lemma). *If δ is small enough (depending on A), then we have*

$$\int_{Q_0} |\mathcal{R}_\mu \chi_{AQ_0}|^2 d\mu \lesssim \left(\varepsilon + \frac{1}{A^2} + A^{4n+2} \delta^{1/(4n+4)} \right) \mu(Q_0).$$

Proof. Note first that, by standard estimates, for $x, y \in Q_0$,

$$\begin{aligned} |\mathcal{R}_\mu \chi_{(AQ_0)^c}(x) - \mathcal{R}_\mu \chi_{(AQ_0)^c}(y)| &\lesssim \int_{(AQ_0)^c} \frac{|x - y|}{|x - z|^{n+1}} d\mu(z) \\ &\lesssim \frac{|x - y|}{\ell(AQ_0)} P_\mu(AQ_0) \lesssim \frac{1}{A} P_\mu(AQ_0) \lesssim \frac{1}{A}, \end{aligned}$$

taking into account the assumption (b) of the Main Lemma for the last inequality. As a consequence,

$$|\mathcal{R}_\mu \chi_{(AQ_0)^c}(x) - m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{(AQ_0)^c})| \lesssim \frac{1}{A},$$

and so

$$\int_{Q_0} |\mathcal{R}_\mu \chi_{(AQ_0)^c}(x) - m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{(AQ_0)^c})|^2 d\mu(x) \lesssim \frac{1}{A^2} \mu(Q_0).$$

Together with the assumption (g) in the Main Lemma this gives

$$\begin{aligned} \int_{Q_0} |\mathcal{R}_\mu \chi_{AQ_0} - m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{AQ_0})|^2 d\mu &\leq 2 \int_{Q_0} |\mathcal{R}_\mu - m_{\mu, Q_0}(\mathcal{R}_\mu)|^2 d\mu \\ &\quad + 2 \int_{Q_0} |\mathcal{R}_\mu \chi_{(AQ_0)^c} - m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{(AQ_0)^c})|^2 d\mu \\ &\lesssim \varepsilon \mu(Q_0) + \frac{1}{A^2} \mu(Q_0). \end{aligned}$$

Hence, to conclude the proof of the lemma it suffices to show that

$$|m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{AQ_0})| \lesssim A^{2n+1} \delta^{1/(8n+8)} \mu(Q_0).$$

By the antisymmetry of the Riesz kernel we have $m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{Q_0}) = 0$, and so the preceding estimate is equivalent to

$$(4.2) \quad |m_{\mu, Q_0}(\mathcal{R}_\mu \chi_{AQ_0 \setminus Q_0})| \leq A^{2n+1} \delta^{1/(8n+8)} \mu(Q_0).$$

To prove (4.2) first we take some small constant $0 < \kappa < 1/10$ to be fixed below. We let φ be some C^1 function which equals 1 on $(1-\kappa)AQ_0 \setminus (1+\kappa)Q_0$ and vanishes out of $AQ_0 \setminus (1+\frac{\kappa}{2})Q_0$, so that φ is even and further $\|\nabla \varphi\|_\infty \lesssim (\kappa \ell(Q_0))^{-1}$. Then we split

$$(4.3) \quad \left| \int_{Q_0} \mathcal{R}_\mu \chi_{AQ_0 \setminus Q_0} d\mu \right| \leq \int_{Q_0} |\mathcal{R}_\mu (\chi_{AQ_0 \setminus Q_0} - \varphi)| d\mu + \left| \int_{Q_0} \mathcal{R}_\mu \varphi d\mu \right|.$$

To bound the first integral on the right hand side note that $\chi_{AQ_0 \setminus Q_0} - \varphi = \psi_1 + \psi_2$, with

$$|\psi_1| \leq \chi_{AQ_0 \setminus (1-\kappa)AQ_0} \quad \text{and} \quad |\psi_2| \leq \chi_{(1+\kappa)Q_0 \setminus Q_0}.$$

Then we have

$$\begin{aligned} \int_{Q_0} |\mathcal{R}_\mu (\chi_{AQ_0 \setminus Q_0} - \varphi)| d\mu &\leq \int_{Q_0} |\mathcal{R}_\mu \psi_1| d\mu + \int_{Q_0} |\mathcal{R}_\mu \psi_2| d\mu \\ &\leq \|\mathcal{R}_\mu \psi_1\|_{L^\infty(\mu|_{Q_0})} \mu(Q_0) + \|\mathcal{R}_\mu \psi_2\|_{L^2(\mu|_{Q_0})} \mu(Q_0)^{1/2}. \end{aligned}$$

Since $\text{dist}(\text{supp } \psi_1, Q_0) \approx A\ell(Q_0)$, we have

$$\|\mathcal{R}_\mu \psi_1\|_{L^\infty(\mu|_{Q_0})} \lesssim \frac{1}{(A\ell(Q_0))^n} \|\psi_1\|_{L^1(\mu)} \leq \frac{1}{(A\ell(Q_0))^n} \mu(AQ_0 \setminus (1-\kappa)AQ_0).$$

On the other hand, since \mathcal{R}_μ is bounded in $L^2(\mu|_{(1+\kappa)Q_0})$, by the assumption (d) in the Main Lemma, and by the thin boundary property of Q_0 (in combination with the fact that $\mu(2Q_0) \approx \mu(Q_0)$), we get

$$\|\mathcal{R}_\mu \psi_2\|_{L^2(\mu|_{Q_0})} \leq C_1 \|\psi_2\|_{L^2(\mu)} \leq C_1 \mu((1+\kappa)Q_0 \setminus Q_0)^{1/2} \leq C(C_0, C_1) \kappa^{1/2} \mu(Q_0)^{1/2}.$$

Therefore,

$$(4.4) \quad \int_{Q_0} |\mathcal{R}_\mu (\chi_{AQ_0 \setminus Q_0} - \varphi)| d\mu \lesssim \frac{1}{A^n} \mu(AQ_0 \setminus (1-\kappa)AQ_0) + \kappa^{1/2} \mu(Q_0).$$

To estimate $\mu(AQ_0 \setminus (1 - \kappa)AQ_0)$ we will use the fact that $\alpha_\mu^H(3AQ_0) \leq \delta$. To this end, first we consider a function $\tilde{\varphi}$ supported on $A(1 + \kappa)Q_0 \setminus (1 - 2\kappa)AQ_0$ which equals 1 on $AQ_0 \setminus (1 - \kappa)AQ_0$, with $\|\nabla \tilde{\varphi}\|_\infty \lesssim 1/(A\kappa\ell(Q_0))$. Then we have

(4.5)

$$\begin{aligned} \mu(AQ_0 \setminus (1 - \kappa)AQ_0) &\leq \int \tilde{\varphi} d\mu \\ &\leq \left| \int \tilde{\varphi} d(\mu - \mathcal{L}_H) \right| + \int \tilde{\varphi} d\mathcal{L}_H \\ &\leq \|\nabla \tilde{\varphi}\|_\infty \ell(3AQ_0)^{n+1} \alpha_\mu^H(3AQ_0) + \mathcal{L}_H((1 + \kappa)AQ_0 \setminus (1 - 2\kappa)AQ_0) \\ &\lesssim \left(\frac{A^n}{\kappa} \delta + \kappa A^n \right) \ell(Q_0)^n, \end{aligned}$$

where we used the estimate for c_H in Lemma 4.1 for the last inequality. Hence, plugging this estimate into (4.4) we obtain

$$(4.6) \quad \int_{Q_0} |\mathcal{R}_\mu(\chi_{AQ_0 \setminus Q_0} - \varphi)| d\mu \lesssim \left(\frac{1}{\kappa} \delta + \kappa + \kappa^{1/2} \right) \mu(Q_0) \lesssim \left(\frac{\delta}{\kappa} + \kappa^{1/2} \right) \mu(Q_0).$$

It remains to estimate the last summand in the inequality (4.3). To this end we write

$$\begin{aligned} (4.7) \quad \left| \int_{Q_0} \mathcal{R}_\mu \varphi d\mu \right| &\leq \left| \int_{Q_0} \mathcal{R}_\mu \varphi d(\mu - \mathcal{L}_H) \right| + \left| \int_{Q_0} \mathcal{R}(\varphi\mu - \varphi\mathcal{L}_H) d\mathcal{L}_H \right| \\ &\quad + \left| \int_{Q_0} \mathcal{R}(\varphi\mathcal{L}_H) d\mathcal{L}_H \right| \\ &= T_1 + T_2 + T_3. \end{aligned}$$

Since φ is even, by the antisymmetry of the Riesz kernel it follows easily that $T_3 = 0$.

To deal with T_1 , consider another auxiliary function $\hat{\varphi}$ supported on Q_0 which equals 1 on $(1 - \hat{\kappa})Q_0$, for some small constant $0 < \hat{\kappa} < \kappa$, so that $\|\nabla \hat{\varphi}\|_\infty \lesssim 1/(\hat{\kappa}\ell(Q_0))$. Then we write

$$T_1 \leq \left| \int \hat{\varphi} \mathcal{R}_\mu \varphi d(\mu - \mathcal{L}_H) \right| + \left| \int (\chi_{Q_0} - \hat{\varphi}) \mathcal{R}_\mu \varphi d(\mu - \mathcal{L}_H) \right| = T_{1,a} + T_{1,b}.$$

To estimate $T_{1,a}$ we set

$$T_{1,a} \leq \|\nabla(\hat{\varphi} \mathcal{R}_\mu \varphi)\|_\infty \ell(3AQ_0)^{n+1} \alpha^H(3AQ_0).$$

We write

$$\|\nabla(\hat{\varphi} \mathcal{R}_\mu \varphi)\|_\infty \leq \|\nabla(\mathcal{R}_\mu \varphi)\|_{\infty, Q_0} + \|\nabla \hat{\varphi}\|_\infty \|\mathcal{R}_\mu \varphi\|_{\infty, Q_0}.$$

Since $\text{dist}(\text{supp } \varphi, Q_0) \geq \frac{\kappa}{2}\ell(Q_0)$ and $\mu(AQ_0) \lesssim \ell(AQ_0)^n$ (by Lemma 4.1), we have

$$\|\mathcal{R}_\mu \varphi\|_{\infty, Q_0} \lesssim \frac{\mu(AQ_0)}{(\kappa\ell(Q_0))^n} \lesssim \frac{A^n}{\kappa^n},$$

and, anagously,

$$\|\nabla(\mathcal{R}_\mu \varphi)\|_{\infty, Q_0} \lesssim \frac{\mu(AQ_0)}{(\kappa\ell(Q_0))^{n+1}} \lesssim \frac{A^n}{\kappa^{n+1}\ell(Q_0)}.$$

Hence,

$$\|\nabla(\hat{\varphi} \mathcal{R}_\mu \varphi)\|_\infty \lesssim \frac{A^n}{\kappa^{n+1}\ell(Q_0)} + \frac{A^n}{\hat{\kappa}\kappa^n\ell(Q_0)} \lesssim \frac{A^n}{\hat{\kappa}\kappa^n\ell(Q_0)}.$$

So we have

$$T_{1,a} \lesssim \frac{A^{2n+1}}{\widehat{\kappa} \kappa^n} \delta \mu(Q_0).$$

Now we consider the term $T_{1,b}$. We write

$$T_{1,b} \leq \|\chi_{Q_0} - \widehat{\varphi}\|_{L^1(\mu + \mathcal{L}_H)} \|\mathcal{R}_\mu \varphi\|_{\infty, Q_0}.$$

Recall that $\|\mathcal{R}_\mu \varphi\|_{\infty, Q_0} \lesssim \frac{A^n}{\kappa^n}$. Also, by the construction of $\widehat{\varphi}$ and the thin boundary of Q_0 ,

$$\|\chi_{Q_0} - \widehat{\varphi}\|_{L^1(\mu + \mathcal{L}_H)} \lesssim \mu(Q_0 \setminus (1 - \widehat{\kappa})Q_0) + \mathcal{L}_H(Q_0 \setminus (1 - \widehat{\kappa})Q_0) \lesssim \widehat{\kappa} \mu(Q_0).$$

So we obtain

$$T_{1,b} \leq \frac{A^n}{\kappa^n} \widehat{\kappa} \mu(Q_0).$$

Thus,

$$T_1 \lesssim \left(\frac{A^{2n+1}}{\kappa^n \widehat{\kappa}} \delta + \frac{A^n}{\kappa^n} \widehat{\kappa} \right) \mu(Q_0).$$

Choosing $\widehat{\kappa} = \delta^{1/2}$, say, we get

$$T_1 \lesssim \left(\frac{A^{2n+1}}{\kappa^n} + \frac{A^n}{\kappa^n} \right) \delta^{1/2} \mu(Q_0) \leq \frac{A^{2n+1}}{\kappa^n} \delta^{1/2} \mu(Q_0).$$

Finally, we turn our attention to T_2 . By Fubini we have

$$T_2 = \left| \int \mathcal{R}(\chi_{Q_0} \mathcal{L}_H) \varphi d(\mu - \mathcal{L}_H) \right| \leq \|\nabla(\mathcal{R}(\chi_{Q_0} \mathcal{L}_H) \varphi)\|_{\infty} \ell(3AQ_0)^{n+1} \alpha^H(3AQ_0).$$

Observe that

$$\|\nabla(\mathcal{R}(\chi_{Q_0} \mathcal{L}_H) \varphi)\|_{\infty} \leq \|\nabla(\mathcal{R}(\chi_{Q_0} \mathcal{L}_H))\|_{\infty, \text{supp } \varphi} + \|\mathcal{R}(\chi_{Q_0} \mathcal{L}_H)\|_{\infty, \text{supp } \varphi} \|\nabla \varphi\|_{\infty}.$$

Using that $\text{dist}(\text{supp } \varphi, Q_0) \geq \frac{\kappa}{2} \ell(Q_0)$ and that $\mathcal{L}_H(Q_0) = c_H \mathcal{H}^n(Q_0 \cap H) \approx \ell(Q_0)^n$, we derive

$$\|\mathcal{R}(\chi_{Q_0} \mathcal{L}_H)\|_{\infty, \text{supp } \varphi} \lesssim \frac{\mathcal{L}_H(Q_0)}{(\kappa \ell(Q_0))^n} \lesssim \frac{1}{\kappa^n}$$

and

$$\|\nabla(\mathcal{R}(\chi_{Q_0} \mathcal{L}_H))\|_{\infty, \text{supp } \varphi} \lesssim \frac{\mathcal{L}_H(Q_0)}{(\kappa \ell(Q_0))^{n+1}} \lesssim \frac{1}{\kappa^{n+1} \ell(Q_0)}.$$

So we obtain

$$T_2 \lesssim \frac{A^{n+1}}{\kappa^{n+1}} \delta \mu(Q_0).$$

Gathering the estimates for T_1 and T_2 , by (4.7) we deduce

$$\left| \int_{Q_0} \mathcal{R}_\mu \varphi d\mu \right| \lesssim \frac{A^{2n+1}}{\kappa^n} \delta^{1/2} \mu(Q_0) + \frac{A^{n+1}}{\kappa^{n+1}} \delta \mu(Q_0) \lesssim \frac{A^{2n+1}}{\kappa^{n+1}} \delta^{1/2} \mu(Q_0).$$

Plugging this estimate and (4.6) into (4.3), we obtain

$$\begin{aligned} \left| \int_{Q_0} \mathcal{R}_\mu \chi_{AQ_0 \setminus Q_0} d\mu \right| &\lesssim \left(\frac{\delta}{\kappa} + \kappa^{1/2} \right) \mu(Q_0) + \frac{A^{2n+1}}{\kappa^{n+1}} \delta^{1/2} \mu(Q_0) \\ &\lesssim \left(\frac{A^{2n+1}}{\kappa^{n+1}} \delta^{1/2} + \kappa^{1/2} \right) \mu(Q_0). \end{aligned}$$

So if we choose $\kappa = \delta^{1/(4n+4)}$, we get

$$\left| \int_{Q_0} \mathcal{R}_\mu \chi_{AQ_0 \setminus Q_0} d\mu \right| \lesssim \left(A^{2n+1} \delta^{1/4} + \delta^{1/(8n+8)} \right) \mu(Q_0) \lesssim A^{2n+1} \delta^{1/(8n+8)} \mu(Q_0),$$

which yields (4.2) and finishes the proof of the lemma. \square

From now on we will assume that δ is small enough, depending on A , so that the conclusion in the preceding lemma holds.

5. THE DYADIC LATTICE OF DAVID AND MATTILA

Now we will consider the dyadic lattice of cells with small boundaries of David-Mattila associated with a Radon measure σ [DM, Theorem 3.2]. Its properties are summarized in the next lemma.

Lemma 5.1 (David, Mattila). *Let σ be a compactly supported Radon measure in \mathbb{R}^{n+1} . Consider two constants $K_0 > 1$ and $A_0 > 5000 K_0$ and denote $W = \text{supp } \sigma$. Then there exists a sequence of partitions of W into Borel subsets Q , $Q \in \mathcal{D}_{\sigma,k}$, which we will refer to as cells, with the following properties:*

- For each integer $k \geq 0$, W is the disjoint union of the cells Q , $Q \in \mathcal{D}_{\sigma,k}$, and if $k < l$, $Q \in \mathcal{D}_{\sigma,l}$, and $R \in \mathcal{D}_{\sigma,k}$, then either $Q \cap R = \emptyset$ or else $Q \subset R$.
- The general position of the cells Q can be described as follows. For each $k \geq 0$ and each cell $Q \in \mathcal{D}_{\sigma,k}$, there is a ball $B(Q) = B(z_Q, r(Q))$ such that

$$z_Q \in W, \quad A_0^{-k} \leq r(Q) \leq K_0 A_0^{-k},$$

$$W \cap B(Q) \subset Q \subset W \cap 28B(Q) = W \cap B(z_Q, 28r(Q)),$$

and

the balls $5B(Q)$, $Q \in \mathcal{D}_{\sigma,k}$, are disjoint.

- The cells $Q \in \mathcal{D}_{\sigma,k}$ have small boundaries. That is, for each $Q \in \mathcal{D}_{\sigma,k}$ and each integer $l \geq 0$, set

$$N_l^{\text{ext}}(Q) = \{x \in W \setminus Q : \text{dist}(x, Q) < A_0^{-k-l}\},$$

$$N_l^{\text{int}}(Q) = \{x \in Q : \text{dist}(x, W \setminus Q) < A_0^{-k-l}\},$$

and

$$N_l(Q) = N_l^{\text{ext}}(Q) \cup N_l^{\text{int}}(Q).$$

Then

$$(5.1) \quad \sigma(N_l(Q)) \leq (C^{-1} K_0^{-3(n+1)-1} A_0)^{-l} \sigma(90B(Q)).$$

- Denote by $\mathcal{D}_{\sigma,k}^{\text{db}}$ the family of cells $Q \in \mathcal{D}_{\sigma,k}$ for which

$$(5.2) \quad \sigma(100B(Q)) \leq K_0 \sigma(B(Q)).$$

We have that $r(Q) = A_0^{-k}$ when $Q \in \mathcal{D}_{\sigma,k} \setminus \mathcal{D}_{\sigma,k}^{\text{db}}$ and

$$(5.3) \quad \sigma(100B(Q)) \leq K_0^{-l} \sigma(100^{l+1}B(Q)) \quad \text{for all } l \geq 1 \text{ with } 100^l \leq K_0 \text{ and } Q \in \mathcal{D}_{\sigma,k} \setminus \mathcal{D}_{\sigma,k}^{\text{db}}.$$

We use the notation $\mathcal{D}_\sigma = \bigcup_{k \geq 0} \mathcal{D}_{\sigma,k}$. Observe that the families $\mathcal{D}_{\sigma,k}$ are only defined for $k \geq 0$. So the diameters of the cells from \mathcal{D} are uniformly bounded from above.

Remark 5.2. Any two disjoint cells $Q, Q' \in \mathcal{D}_\sigma$ satisfy $\frac{1}{2}B(Q) \cap \frac{1}{2}B(Q') = \emptyset$. This holds with $\frac{1}{2}$ replaced by 5 by the statements in the lemma above in the case that Q, Q' are of the same generation $\mathcal{D}_{\sigma,k}$. If $Q \in \mathcal{D}_{\sigma,j}$ and $Q' \in \mathcal{D}_{\sigma,k}$ with $j \neq k$, this follows easily too. Indeed, assume $j < k$, and suppose that $\frac{1}{2}B(Q) \cap \frac{1}{2}B(Q') \neq \emptyset$. Since $r(Q') \ll r(Q)$ (by choosing A_0 big enough), this implies that $B(Q') \subset B(Q)$, and so

$$B(Q') \cap W \subset B(Q) \cap W \subset Q,$$

which implies that $Q' \cap Q \neq \emptyset$ and gives a contradiction.

Given $Q \in \mathcal{D}_{\sigma,k}$, we denote $J(Q) = k$. We set $\ell(Q) = 56 K_0 A_0^{-k}$ and we call it the side length of Q . Note that

$$\frac{1}{28} K_0^{-1} \ell(Q) \leq \text{diam}(28B(Q)) \leq \ell(Q).$$

Observe that $r(Q) \approx \text{diam}(Q) \approx \ell(Q)$. Also we call z_Q the center of Q , and the cell $Q' \in \mathcal{D}_{\sigma,k-1}$ such that $Q' \supset Q$ the parent of Q . We set $B_Q = 28B(Q) = B(z_Q, 28r(Q))$, so that

$$W \cap \frac{1}{28}B_Q \subset Q \subset B_Q.$$

We assume A_0 to be big enough so that the constant $C^{-1} K_0^{-3(n+1)-1} A_0$ in (5.1) satisfies

$$C^{-1} K_0^{-3(n+1)-1} A_0 > A_0^{1/2} > 10.$$

Then we deduce that, for all $0 < \lambda \leq 1$,

$$\begin{aligned} & \sigma(\{x \in Q : \text{dist}(x, W \setminus Q) \leq \lambda \ell(Q)\}) + \sigma(\{x \in 3.5B_Q : \text{dist}(x, Q) \leq \lambda \ell(Q)\}) \\ (5.4) \quad & \leq c \lambda^{1/2} \sigma(3.5B_Q). \end{aligned}$$

We denote $\mathcal{D}_\sigma^{db} = \bigcup_{k \geq 0} \mathcal{D}_{\sigma,k}^{db}$. Note that, in particular, from (5.2) it follows that

$$(5.5) \quad \sigma(3.5B_Q) \leq \sigma(100B(Q)) \leq K_0 \sigma(Q) \quad \text{if } Q \in \mathcal{D}_\sigma^{db}.$$

For this reason we will call the cells from \mathcal{D}_σ^{db} doubling. Given $Q \in \mathcal{D}_\sigma$, we denote by $\mathcal{D}_\sigma(Q)$ the family of cells from \mathcal{D}_σ which are contained in Q . Analogously, we write $\mathcal{D}_\sigma^{db}(Q) = \mathcal{D}_\sigma^{db} \cap \mathcal{D}(Q)$.

As shown in [DM, Lemma 5.28], every cell $R \in \mathcal{D}_\sigma$ can be covered σ -a.e. by a family of doubling cells:

Lemma 5.3. *Let $R \in \mathcal{D}_\sigma$. Suppose that the constants A_0 and K_0 in Lemma 5.1 are chosen suitably. Then there exists a family of doubling cells $\{Q_i\}_{i \in I} \subset \mathcal{D}_\sigma^{db}$, with $Q_i \subset R$ for all i , such that their union covers σ -almost all R .*

The following result is proved in [DM, Lemma 5.31].

Lemma 5.4. *Let $R \in \mathcal{D}_\sigma$ and let $Q \subset R$ be a cell such that all the intermediate cells $S, Q \subsetneq S \subsetneq R$ are non-doubling (i.e. belong to $\mathcal{D}_\sigma \setminus \mathcal{D}_\sigma^{db}$). Then*

$$(5.6) \quad \sigma(100B(Q)) \leq A_0^{-10n(J(Q)-J(R)-1)} \sigma(100B(R)).$$

From the preceding lemma we deduce:

Lemma 5.5. *Let $Q, R \in \mathcal{D}_\sigma$ be as in Lemma 5.4. Then*

$$\Theta_\sigma(100B(Q)) \leq K_0 A_0^{-9n(J(Q)-J(R)-1)} \Theta_\sigma(100B(R))$$

and

$$\sum_{S \in \mathcal{D}_\sigma: Q \subset S \subset R} \Theta_\sigma(100B(S)) \leq C \Theta_\sigma(100B(R)),$$

with C depending on K_0 and A_0 .

For the easy proof, see [To3, Lemma 4.4], for example.

6. THE LOW DENSITY CELLS AND THE STOPPING CELLS

We consider the measure $\sigma = \mu|_{Q_0}$ and the associated dyadic lattice \mathcal{D}_σ introduced in Section 5, re-scaled appropriately, so that we can assume that Q_0 is a cell from \mathcal{D}_σ . Below we allow all the constants denoted by C and all the implicit constants in the relations “ \lesssim ” and “ \approx ” to depend on the constants A_0 and K_0 from the construction of the lattice \mathcal{D}_σ .

Let $0 < \theta_0 \ll 1$ be a very small constant to be fixed later. We denote by LD the family of those cells from \mathcal{D}_σ such that $\Theta_\sigma(3.5B_Q) \leq \theta_0$ and have maximal side length.

The main difficulty for the proof of the Main Lemma 3.1 consists in showing that the following holds.

Key Lemma 6.1. *There exists some constant ε_0 such that if A is big enough and $\theta_0, \delta, \varepsilon$ are small enough (with δ possibly depending on A), then*

$$\mu\left(\bigcup_{Q \in \text{LD}} Q\right) \leq (1 - \varepsilon_0) \mu(Q_0).$$

The proof of this result will be carried out along the next sections of this paper. In what follows we will assume that

$$(6.1) \quad \mu\left(\bigcup_{Q \in \text{LD}} Q\right) > (1 - \varepsilon_0) \mu(Q_0)$$

and we will get a contradiction for ε_0 small enough. To this end, first we need to construct another family of stopping cells which we will denote by Stop. This is defined as follows. For each $Q \in \text{LD}$ we consider the family of maximal cells contained in Q from \mathcal{D}_σ^{db} (so they are doubling) with side length at most $t\ell(Q)$, where $0 < t < 1$ is some small parameter which will be fixed below. We denote this family by $\text{Stop}(Q)$. Then we define

$$\text{Stop} = \bigcup_{Q \in \text{LD}} \text{Stop}(Q).$$

Note that, by Lemma 5.3, it is immediate that, for each $Q \in \text{LD}$, the cells from $\text{Stop}(Q)$ cover μ -almost all Q . So proving the assumption (6.1) is equivalent to showing

$$\mu\left(\bigcup_{Q \in \text{Stop}} Q\right) > (1 - \varepsilon_0) \mu(Q_0)$$

We need the following auxiliary result.

Lemma 6.2. *If we choose $t = \theta_0^{1/(n+1)}$, then we have:*

$$\Theta_\mu(2B_Q) \leq P_\mu(2B_Q) \lesssim \theta_0^{1/(n+1)} \quad \text{for all } Q \in \text{Stop}.$$

Proof. Let $Q \in \text{Stop}$ and $R \in \text{LD}$ such that $Q \subset R$. The first inequality in the lemma is trivial and so we only have to prove the second one. Let $R' \in \mathcal{D}_\sigma$ the maximal cell such that $Q \subset R' \subset R$ with $\ell(R') \leq t\ell(R)$, so that $\ell(R') \approx t\ell(R)$. Then we write

$$\begin{aligned} P_\mu(2B_Q) &\lesssim \sum_{P \in \mathcal{D}_\sigma: Q \subset P \subset R'} \Theta_\mu(2B_P) \frac{\ell(Q)}{\ell(P)} + \sum_{P \in \mathcal{D}_\sigma: R' \subset P \subset R} \Theta_\mu(2B_P) \frac{\ell(Q)}{\ell(P)} \\ &\quad + \sum_{P \in \mathcal{D}_\sigma: R \subset P \subset Q_0} \Theta_\mu(2B_P) \frac{\ell(Q)}{\ell(P)} + \sum_{k \geq 1} \Theta_\mu(2^k Q_0) \frac{\ell(Q)}{\ell(2^k Q_0)} \\ &= S_1 + S_2 + S_3 + S_4. \end{aligned}$$

To deal with the sums S_1 and S_2 , note that for all $P \subset R$, since $2B_P \subset 2B_R$ (assuming A_0 to be big enough), we have

$$\Theta_\mu(2B_P) = \frac{\mu(2B_P)}{r(2B_P)^n} \leq \frac{\mu(2B_R)}{r(2B_P)^n} = \Theta_\mu(2B_R) \frac{r(B_R)^n}{r(B_P)^n} \approx \Theta_\mu(2B_R) \frac{\ell(R)^n}{\ell(P)^n}.$$

Therefore, since $\Theta_\mu(2B_R) \lesssim \theta_0$ and all the cells P appearing in S_2 satisfy $\ell(P) \geq t\ell(R)$, we deduce that all such cells satisfy $\Theta_\mu(2B_P) \lesssim \frac{1}{t^n} \Theta_\mu(2B_R) \lesssim \frac{\theta_0}{t^n}$, and thus

$$S_2 \lesssim \frac{\theta_0}{t^n} \sum_{P \in \mathcal{D}_\sigma: R' \subset P \subset R} \frac{\ell(Q)}{\ell(P)} \lesssim \frac{\theta_0}{t^n}.$$

Also, since there are no μ -doubling cells between R' and Q , from Lemma 5.5 we deduce that the cells P in the sum S_1 satisfy

$$\Theta_\mu(2B_P) \lesssim \Theta_\mu(2B_{R'}) \lesssim \frac{1}{t^n} \Theta_\mu(2B_R) \lesssim \frac{\theta_0}{t^n},$$

and therefore we also get

$$S_1 \lesssim \frac{\theta_0}{t^n} \sum_{P \in \mathcal{D}_\sigma: Q \subset P \subset R'} \frac{\ell(Q)}{\ell(P)} \lesssim \frac{\theta_0}{t^n}.$$

For the cells P in the sum S_3 we just take into account that $\Theta_\mu(2B_P) \lesssim 1$, and thus

$$S_3 \lesssim \sum_{P \in \mathcal{D}_\sigma: R \subset P \subset Q_0} \frac{\ell(Q)}{\ell(P)} \lesssim \frac{\ell(Q)}{\ell(R)} \lesssim t.$$

Regarding the sum S_4 , note that

$$S_4 = \sum_{1 \leq k < \log_2 A} \Theta_\mu(2^k Q_0) \frac{\ell(Q)}{\ell(2^k Q_0)} + \sum_{k: 2^k \geq A} \Theta_\mu(2^k Q_0) \frac{\ell(Q)}{\ell(2^k Q_0)} = S_{4,a} + S_{4,b}.$$

For the indices k in $S_{4,a}$ we use the fact that $\Theta_\mu(2^k Q_0) \lesssim C_0$, and so we get

$$S_{4,a} \lesssim \sum_{1 \leq k < \log_2 A} \frac{\ell(Q)}{\ell(2^k Q_0)} \lesssim \frac{\ell(Q)}{\ell(Q_0)} \leq t.$$

For $S_{4,b}$ we write

$$S_{4,b} \lesssim \frac{\ell(Q)}{\ell(AQ_0)} \sum_{j \geq 0} \Theta_\mu(2^j AQ_0) \frac{\ell(AQ_0)}{\ell(2^j AQ_0)} \lesssim \frac{\ell(Q)}{\ell(AQ_0)} P_\mu(AQ_0) \lesssim \frac{\ell(Q)}{\ell(AQ_0)} \leq t.$$

Hence,

$$P_\mu(2B_Q) \lesssim \frac{\theta_0}{t^n} + t \approx \theta_0^{1/(n+1)},$$

recalling that $t = \theta_0^{1/(n+1)}$. □

From now on we assume that we have chosen $t = \theta_0^{1/(n+1)}$, so that the conclusion of the preceding lemma holds.

The family Stop may consist of an infinite number of cells. For technical reasons, it is convenient to consider a finite subfamily of Stop which contains a very big proportion of the μ measure of Stop . So we let Stop_0 be a *finite* subfamily of Stop such that

$$(6.2) \quad \mu\left(\bigcup_{Q \in \text{Stop}_0} Q\right) > (1 - 2\varepsilon_0) \mu(Q_0).$$

We denote by Bad the family of the cells $P \in \text{Stop}$ such that $1.1B_P \cap \partial Q_0 \neq \emptyset$.

Lemma 6.3. *We have*

$$\mu\left(\bigcup_{Q \in \text{Bad}} Q\right) \lesssim \theta_0^{1/(n+1)} \mu(Q_0).$$

Proof. Let $I \subset \text{Bad}$ an arbitrary finite family of bad cells. We apply the covering theorem of triple balls of Vitali to the family $\{1.15B_Q\}_{Q \in I}$, so that we get a subfamily $J \subset I$ satisfying

- $1.15B_P \cap 1.15B_Q = \emptyset$ for different cells $P, Q \in J$, and
- $\bigcup_{P \in I} 1.15B_P \subset \bigcup_{Q \in J} 3.45B_Q$.

Then, using that

$$\mu(3.45B_Q) \leq \mu(3.5B_Q) \lesssim \mu(B_Q) \lesssim \theta_0^{1/(n+1)} r(B_Q)^n \quad \text{for all } Q \in J,$$

we get

$$\mu\left(\bigcup_{P \in I} B_P\right) \leq \sum_{Q \in J} \mu(3.45B_Q) \lesssim \theta_0^{1/(n+1)} \sum_{Q \in J} r(B_Q)^n.$$

For each $Q \in J$ we have $1.1B_Q \cap \partial Q_0 \neq \emptyset$ and so we deduce that

$$\mathcal{H}^n(1.15B_Q \cap \partial Q_0) \gtrsim r(B_Q)^n.$$

Thus, using also that the balls $1.15B_Q$, $Q \in J$, are pairwise disjoint,

$$\mu\left(\bigcup_{P \in I} P\right) \lesssim \theta_0^{1/(n+1)} \sum_{Q \in J} \mathcal{H}^n(1.15B_Q \cap \partial Q_0) \leq \theta_0^{1/(n+1)} \mathcal{H}^n(\partial Q_0) \approx \theta_0^{1/(n+1)} \mu(Q_0),$$

and the lemma follows. □

We will now define an auxiliary measure μ_0 . First, given a small constant $0 < \kappa_0 \ll 1$ (to be fixed below) and $Q \in \mathcal{D}_\sigma$, we denote

$$(6.3) \quad I_{\kappa_0}(Q) = \{x \in Q : \text{dist}(x, \text{supp } \sigma \setminus Q) \geq \kappa_0 \ell(Q)\}.$$

So $I_{\kappa_0}(Q)$ is some kind of inner subset of Q . We set

$$\mu_0 = \mu|_{Q_0^c} + \sum_{Q \in \text{Stop}_0 \setminus \text{Bad}} \mu|_{I_{\kappa_0}(Q)}.$$

Observe that, by the doubling and small boundary condition (5.4) of $Q \in \text{Stop}_0$, we have

$$\mu(Q \setminus I_{\kappa_0}(Q)) \lesssim \kappa_0^{1/2} \mu(3.5B_Q) \lesssim \kappa_0^{1/2} \mu(Q).$$

Combining this estimate with (6.2) and Lemma 6.3, we get

$$(6.4) \quad \begin{aligned} \|\mu - \mu_0\| &= \mu(Q_0) - \mu_0(Q_0) \\ &= \mu(Q_0) - \sum_{Q \in \text{Stop}_0 \setminus \text{Bad}} \mu(I_{\kappa_0}(Q)) \\ &= \mu\left(Q_0 \setminus \bigcup_{Q \in \text{Stop}_0} Q\right) + \sum_{Q \in \text{Bad}} \mu(Q) + \sum_{Q \in \text{Stop}_0 \setminus \text{Bad}} \mu(Q \setminus I_{\kappa_0}(Q)) \\ &\leq 2\varepsilon_0 \mu(Q_0) + C\theta_0^{1/(n+1)} \mu(Q_0) + C\kappa_0^{1/2} \mu(Q_0). \end{aligned}$$

Together with Lemma 4.2 this yields the following.

Lemma 6.4. *If δ is small enough (depending on A), then we have*

$$\int_{Q_0} |\mathcal{R}(\chi_{AQ_0}\mu_0)|^2 d\mu_0 \lesssim \left(\varepsilon + \frac{1}{A^2} + \delta^{1/(8n+8)} + \varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2}\right) \mu(Q_0).$$

Proof. We have

$$\begin{aligned} \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}\mu_0)|^2 d\mu_0 &\leq 2 \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}\mu)|^2 d\mu + 2 \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}(\mu - \mu_0))|^2 d\mu \\ &\lesssim \left(\varepsilon + \frac{1}{A^2} + A^{4n+2} \delta^{1/(4n+4)} + \varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2}\right) \mu(Q_0), \end{aligned}$$

by Lemma 4.2, the $L^2(\mu|_{Q_0})$ boundedness of $\mathcal{R}_{\mu|_{Q_0}}$, and (6.4). \square

7. THE PERIODIC MEASURE $\tilde{\mu}$

Let \mathcal{M} be the lattice of cubes in \mathbb{R}^{n+1} obtained by translating Q_0 in directions parallel to H , so that H coincides with the union of the n -dimensional cubes from the family $\{P \cap H\}_{P \in \mathcal{M}}$ and the cubes have disjoint interiors. For each $P \in \mathcal{M}$, denote by z_P the center of P and consider the translation $T_P : x \rightarrow x + z_P$, so that $P = T_P(Q_0)$. Note that $\{z_P : P \in \mathcal{M}\}$ coincides with the set $(\ell(Q_0)\mathbb{Z}^n) \times \{0\}$. We define

$$\tilde{\mu} = \sum_{P \in \mathcal{M}} (T_P)_\#(\mu_0|_{Q_0}).$$

That is,

$$\tilde{\mu}(E) = \sum_{P \in \mathcal{M}} \mu_0(Q_0 \cap T_P^{-1}(E)) = \sum_{P \in \mathcal{M}} \mu_0(Q_0 \cap (E - z_P)).$$

It is easy to check that:

- (i) $\tilde{\mu}$ is periodic with respect to \mathcal{M} , that is, for all $P \in \mathcal{M}$ and all $E \subset \mathbb{R}^{n+1}$, $\tilde{\mu}(E + z_P) = \tilde{\mu}(E)$.
- (ii) $\chi_{Q_0} \tilde{\mu} = \mu_0$.

The latter property holds because $\mu_0(\partial Q_0) = 0$.

For simplicity, from now on we will assume that A is a big enough odd natural number.

Lemma 7.1. *We have*

$$\alpha_{\tilde{\mu}}^H(3AQ_0) \leq C_3 A^{n+1} \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \delta^{1/2} \right).$$

In fact,

$$\text{dist}_{3AQ_0}(\tilde{\mu}, \mathcal{L}_H) \leq C_3 A^{n+1} \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \delta^{1/2} \right) \ell(3AQ_0)^{n+1},$$

where \mathcal{L}_H is the same minimizing measure as the one for $\alpha_{\mu}^H(3AQ_0)$.

Proof. Let f be a Lipschitz function supported on $3AQ_0$ with Lipschitz constant at most 1. Denote by \mathcal{M}_0 the family of cubes from \mathcal{M} which are contained in $3AQ_0$. Let $\kappa > 0$ be some small parameter to be fixed below. Consider a C^1 function φ supported on Q_0 which equals 1 on $(1-\kappa)Q_0$, with $\|\nabla \varphi\|_{\infty} \lesssim 1/(\kappa \ell(Q_0))$ and denote $\varphi_P(x) = \varphi(x - z_P)$. Then we write

$$(7.1) \quad \left| \int f d(\tilde{\mu} - \mathcal{L}_H) \right| \leq \sum_{P \in \mathcal{M}_0} \left| \int_P f d(\tilde{\mu} - \mathcal{L}_H) \right| \\ \leq \sum_{P \in \mathcal{M}_0} \left| \int \varphi_P f d(\tilde{\mu} - \mathcal{L}_H) \right| + \sum_{P \in \mathcal{M}_0} \int |(\chi_P - \varphi_P) f| d(\tilde{\mu} + \mathcal{L}_H).$$

Let us estimate the first sum on the right hand side. Since $\tilde{\mu}|_P = (T_P)_{\#} \mu_0|_{Q_0}$ and $\mathcal{L}_H = (T_P)_{\#} \mathcal{L}_H$, we have

$$\left| \int \varphi_P f d(\tilde{\mu} - \mathcal{L}_H) \right| = \left| \int \varphi(x) f(x + z_P) d(\mu_0 - \mathcal{L}_H) \right| \\ \leq \left| \int \varphi(x) f(x + z_P) d(\mu_0 - \mu) \right| + \left| \int \varphi(x) f(x + z_P) d(\mu - \mathcal{L}_H) \right| \\ = I_1 + I_2.$$

To estimate I_1 we use (6.4) and the fact that, by the mean value theorem, $\|\varphi f(\cdot + z_P)\|_{\infty} \lesssim \ell(3AQ_0)$:

$$I_1 \leq \left| \int \varphi(x) f(x + z_P) d(\mu_0 - \mu) \right| \lesssim (\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2}) \ell(3AQ_0)^{n+1}.$$

Concerning I_2 , we write

$$I_2 \lesssim \|\nabla(\varphi f(\cdot + z_P))\|_{\infty} \ell(3AQ_0)^{n+1} \alpha_{\mu}^H(3AQ_0).$$

Note that

$$\|\nabla(\varphi_P f)\|_{\infty} \leq \|\nabla f\|_{\infty} + \|f\|_{\infty} \|\nabla \varphi_P\|_{\infty} \lesssim 1 + C A \ell(Q_0) \frac{1}{\kappa \ell(Q_0)} \lesssim \frac{A}{\kappa}.$$

Thus,

$$I_2 \lesssim A \frac{\delta}{\kappa} \ell(3AQ_0)^{n+1}.$$

Hence,

$$\left| \int \varphi_P f d(\tilde{\mu} - \mathcal{L}_H) \right| \lesssim A \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \frac{\delta}{\kappa} \right) \ell(3AQ_0)^{n+1}.$$

To deal with the last sum on the right hand side of (7.1) we write

$$\begin{aligned} \int |(\chi_P - \varphi_P) f| d(\tilde{\mu} + \mathcal{L}_H) &\leq \|\chi_P - \varphi_P\|_{L^1(\tilde{\mu} + \mathcal{L}_H)} \|f\|_\infty \\ &\lesssim (\mu + \mathcal{L}_H)(Q_0 \setminus (1 - \kappa)Q_0) \ell(3AQ_0). \end{aligned}$$

By the thin boundary condition on Q_0 ,

$$\mu(Q_0 \setminus (1 - \kappa)Q_0) \lesssim \kappa \mu(Q_0) = \kappa \ell(Q_0)^n.$$

Clearly, the same estimate holds replacing μ by \mathcal{L}_H . So we deduce

$$\int |(\chi_P - \varphi_P) f| d(\tilde{\mu} + \mathcal{L}_H) \lesssim \kappa \ell(Q_0)^{n+1}.$$

Taking into account that the number of cubes $P \in \mathcal{M}_0$ is comparable to A^n , we get

$$\left| \int f d(\tilde{\mu} - \mathcal{L}_H) \right| \lesssim A^{n+1} \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \frac{\delta}{\kappa} + \kappa \right) \ell(3AQ_0)^{n+1}.$$

Choosing $\kappa = \delta^{1/2}$, the lemma follows. \square

From now on, to simplify notation we will denote

$$(7.2) \quad \tilde{\delta} = C_3 A^{n+1} \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \delta^{1/2} \right).$$

So the preceding lemma ensures that $\alpha_\mu^H(3AQ_0) \leq \tilde{\delta}$. We assume that the parameters ε_0 , θ_0 , κ_0 , and δ are small enough so that $\tilde{\delta} \ll 1$.

Lemma 7.2. *We have*

$$(7.3) \quad \int_{Q_0} |\mathcal{R}(\chi_{AQ_0} \tilde{\mu})|^2 d\tilde{\mu} \leq C_4 \left(\varepsilon + \frac{1}{A^2} + A^{4n+2} \delta^{\frac{1}{4n+4}} + \varepsilon_0 + \theta_0^{\frac{1}{n+1}} + \kappa_0^{\frac{1}{2}} + A^{2n+2} \tilde{\delta}^{\frac{2}{4n+5}} \right) \tilde{\mu}(Q_0).$$

Proof. Since $\tilde{\mu}|_{Q_0} = \mu_0|_{Q_0}$, we have

$$(7.4) \quad \int_{Q_0} |\mathcal{R}(\chi_{AQ_0} \tilde{\mu})|^2 d\tilde{\mu} \leq 2 \int_{Q_0} |\mathcal{R}(\chi_{AQ_0} \mu_0)|^2 d\mu_0 + 2 \int_{Q_0} |\mathcal{R}(\chi_{AQ_0} (\tilde{\mu} - \mu_0))|^2 d\mu_0.$$

The first integral on the right hand side has been estimated in Lemma 6.4. So we only have to deal with the second one. The arguments that we will use will be similar to some of the ones in Lemma 4.2.

First, note that, using again that $\tilde{\mu}|_{Q_0} = (\mu_0)|_{Q_0}$ and that $(\mu_0)|_{Q_0^c} = \mu|_{Q_0^c}$, we have

$$\mathcal{R}(\chi_{AQ_0} (\tilde{\mu} - \mu_0)) = \mathcal{R}(\chi_{AQ_0 \setminus Q_0} (\tilde{\mu} - \mu)).$$

Let $0 < \kappa < 1/10$ be some small constant to be fixed below. Let φ be a C^1 function which equals 1 on $(1 - \kappa)AQ_0 \setminus (1 + \kappa)Q_0$ and vanishes out of $AQ_0 \setminus (1 + \frac{\kappa}{2})Q_0$, with $\|\nabla \varphi\|_\infty \lesssim (\kappa \ell(Q_0))^{-1}$.

We split

(7.5)

$$\int_{Q_0} |\mathcal{R}(\chi_{AQ_0}(\tilde{\mu} - \mu_0))|^2 d\mu_0 \leq 2 \int_{Q_0} |\mathcal{R}((\chi_{AQ_0 \setminus Q_0} - \varphi)(\tilde{\mu} - \mu))|^2 d\mu_0 + 2 \int_{Q_0} |\mathcal{R}(\varphi(\tilde{\mu} - \mu))|^2 d\mu_0.$$

Concerning the first integral on the right hand side note that $\chi_{AQ_0 \setminus Q_0} - \varphi = \psi_1 + \psi_2$, with

$$|\psi_1| \leq \chi_{AQ_0 \setminus (1-\kappa)AQ_0} \quad \text{and} \quad |\psi_2| \leq \chi_{(1+\kappa)Q_0 \setminus Q_0}.$$

So we have

$$\begin{aligned} \int_{Q_0} |\mathcal{R}((\chi_{AQ_0} - \varphi)(\tilde{\mu} - \mu))|^2 d\mu_0 &\lesssim \int_{Q_0} |\mathcal{R}(\psi_1(\tilde{\mu} - \mu))|^2 d\tilde{\mu} + \int_{Q_0} |\mathcal{R}(\psi_2(\tilde{\mu} - \mu))|^2 d\tilde{\mu} \\ &\leq \|\mathcal{R}(\psi_1(\tilde{\mu} - \mu))\|_{L^\infty(\tilde{\mu}|_{Q_0})}^2 \mu(Q_0) \\ &\quad + \|\mathcal{R}(\psi_2(\tilde{\mu} - \mu))\|_{L^4(\tilde{\mu}|_{Q_0})}^2 \mu(Q_0)^{1/2}. \end{aligned}$$

Since $\text{dist}(\text{supp } \psi_1, Q_0) \approx A\ell(Q_0)$, we get

$$\|\mathcal{R}(\psi_1(\tilde{\mu} - \mu))\|_{L^\infty(\tilde{\mu}|_{Q_0})} \lesssim \frac{1}{(A\ell(Q_0))^n} \|\psi_1\|_{L^1(\tilde{\mu} + \mu)} \leq \frac{1}{(A\ell(Q_0))^n} (\tilde{\mu} + \mu)(AQ_0 \setminus (1-\kappa)AQ_0).$$

Recall that in (4.5) it has been shown that

$$\mu(AQ_0 \setminus (1-\kappa)AQ_0) \lesssim \left(\frac{A^n}{\kappa} \delta + \kappa A^n \right) \ell(Q_0)^n.$$

To prove this we used the fact that $\alpha_\mu^H(3AQ_0) \leq \delta$, or more precisely, that $\text{dist}_{3AQ_0}(\mu, \mathcal{L}_H) \leq \delta$.

The same inequality holds replacing μ by $\tilde{\mu}$ and δ by $\tilde{\delta}$, as shown in Lemma 7.1. So by arguments analogous to the ones in (4.5) it follows that

$$\tilde{\mu}(AQ_0 \setminus (1-\kappa)AQ_0) \lesssim \left(\frac{A^n}{\kappa} \tilde{\delta} + \kappa A^n \right) \ell(Q_0)^n.$$

So we deduce that

$$\|\mathcal{R}(\psi_1(\tilde{\mu} - \mu))\|_{L^\infty(\tilde{\mu}|_{Q_0})} \lesssim \frac{\delta + \tilde{\delta}}{\kappa} + \kappa \lesssim \frac{\tilde{\delta}}{\kappa} + \kappa,$$

taking into account that $\delta \leq \tilde{\delta}$ for the last inequality.

Next we will estimate $\|\mathcal{R}(\psi_2(\tilde{\mu} - \mu))\|_{L^4(\tilde{\mu}|_{Q_0})}$. By the triangle inequality, we have

$$\|\mathcal{R}(\psi_2(\tilde{\mu} - \mu))\|_{L^4(\tilde{\mu}|_{Q_0})} \leq \|\mathcal{R}_\mu \psi_2\|_{L^4(\mu|_{Q_0})} + \|\mathcal{R}_{\tilde{\mu}} \psi_2\|_{L^4(\tilde{\mu}|_{Q_0})}.$$

Recall that \mathcal{R}_μ is bounded in $L^2(\mu|_{2Q_0})$, and so in $L^4(\mu|_{2Q_0})$, and that $\text{supp } \psi_2 \subset (1+\kappa)Q_0 \setminus Q_0 \subset 2Q_0$. Hence, using also the thin boundary property of Q_0 , we obtain

$$\|\mathcal{R}_\mu \psi_2\|_{L^4(\mu|_{Q_0})}^4 \lesssim \|\psi_2\|_{L^4(\mu|_{2Q_0})}^4 \lesssim \mu((1+\kappa)Q_0 \setminus Q_0) \lesssim \kappa \mu(Q_0).$$

We can apply the same argument to estimate $\|\mathcal{R}_{\tilde{\mu}} \psi_2\|_{L^4(\tilde{\mu}|_{Q_0})}$. This is due to the fact that $\mathcal{R}_{\tilde{\mu}}$ is bounded in $L^2(\tilde{\mu}|_{2Q_0})$. This is an easy consequence of the fact that, given two measures μ_1 and μ_2 with growth of order n such that, for $i = 1, 2$, \mathcal{R}_{μ_i} is bounded in $L^2(\mu_i)$, then $\mathcal{R}_{\mu_1 + \mu_2}$ is bounded in $L^2(\mu_1 + \mu_2)$. For the proof, see Proposition 2.25 of [To2], for example. So applying this result to a finite number of translated copies of $\mu|_{Q_0}$, we deduce that $\mathcal{R}_{\tilde{\mu}}$ is bounded in $L^2(\tilde{\mu}|_{2Q_0})$ and so in $L^4(\tilde{\mu}|_{2Q_0})$. Thus we also have

$$\|\mathcal{R}_{\tilde{\mu}} \psi_2\|_{L^4(\tilde{\mu}|_{Q_0})}^4 \lesssim \kappa \tilde{\mu}(Q_0) \leq \kappa \mu(Q_0).$$

Gathering the estimates above, it turns out that the first integral on the right side of (7.5) satisfies the following:

$$(7.6) \quad \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}(\tilde{\mu} - \mu_0))|^2 d\mu_0 \lesssim \left(\frac{\tilde{\delta}}{\kappa} + \kappa \right)^2 \mu(Q_0) + \kappa^{1/2} \mu(Q_0) \lesssim \left(\frac{\tilde{\delta}}{\kappa} + \kappa^{1/4} \right)^2 \mu(Q_0).$$

It remains to estimate the second integral on the right hand side of (7.5). To this end for any $x \in Q_0$ we set

$$\begin{aligned} |\mathcal{R}(\varphi(\tilde{\mu} - \mu_0)(x))| &= \left| \int K(x-y) \varphi(y) d(\tilde{\mu} - \mu)(y) \right| \\ &\leq \left| \int K(x-y) \varphi(y) d(\tilde{\mu} - \mathcal{L}_H)(y) \right| \\ &\quad + \left| \int K(x-y) \varphi(y) d(\mu - \mathcal{L}_H)(y) \right| \\ &\leq \|\nabla(K(x-\cdot)\varphi)\|_\infty [d_{3AQ_0}(\tilde{\mu}, \mathcal{L}_H) + d_{3AQ_0}(\mu, \mathcal{L}_H)], \end{aligned}$$

where in the first identity we used the fact that μ_0 coincides with μ on the support of φ . Taking into account that $\text{dist}(x, \text{supp } \varphi) \gtrsim \kappa \ell(Q_0)$, we obtain

$$\|\nabla(K(x-\cdot)\varphi)\|_\infty \leq \|\nabla K(x-\cdot)\|_{\infty, \text{supp } \varphi} + \|K(x-\cdot)\|_{\infty, \text{supp } \varphi} \|\nabla \varphi\|_\infty \lesssim \frac{1}{(\kappa \ell(Q_0))^{n+1}}.$$

By Lemma 7.1, $\text{dist}_{3AQ_0}(\tilde{\mu}, \mathcal{L}_H) \leq \tilde{\delta} \ell(3AQ_0)^{n+1}$ and, by the assumption (e) in the Main Lemma, $\text{dist}_{3AQ_0}(\mu, \mathcal{L}_H) \leq \delta \ell(3AQ_0)^{n+1}$. Therefore,

$$|\mathcal{R}(\varphi(\tilde{\mu} - \mu_0)(x))| \lesssim \frac{1}{(\kappa \ell(Q_0))^{n+1}} (\delta + \tilde{\delta}) \ell(3AQ_0)^{n+1} \lesssim \frac{A^{n+1}}{\kappa^{n+1}} \tilde{\delta}.$$

So the last integral on the right hand side of (7.5) satisfies

$$(7.7) \quad \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}(\tilde{\mu} - \mu_0))|^2 d\tilde{\mu} \lesssim \left(\frac{A^{n+1}}{\kappa^{n+1}} \tilde{\delta} \right)^2 \mu(Q_0).$$

From (7.5), (7.6) and (7.7) we deduce that

$$\begin{aligned} \int_{Q_0} |\mathcal{R}(\chi_{AQ_0}(\tilde{\mu} - \mu_0))|^2 d\mu_0 &\lesssim \left(\frac{\tilde{\delta}}{\kappa} + \kappa^{1/4} + \frac{A^{n+1}}{\kappa^{n+1}} \tilde{\delta} \right)^2 \mu(Q_0) \\ &\lesssim \left(\kappa^{1/4} + \frac{A^{n+1}}{\kappa^{n+1}} \tilde{\delta} \right)^2 \mu(Q_0). \end{aligned}$$

Choosing $\kappa = \tilde{\delta}^{\frac{4}{4n+5}}$, the right hand side above equals

$$\left(\tilde{\delta}^{\frac{1}{4n+5}} + A^{n+1} \tilde{\delta}^{\frac{1}{4n+5}} \right)^2 \mu(Q_0) \lesssim A^{2n+2} \tilde{\delta}^{\frac{2}{4n+5}} \mu(Q_0).$$

Together with (7.4) and Lemma 7.2, this yields (7.3). \square

To simplify notation we will write

$$(7.8) \quad \tilde{\varepsilon} = C_4 \left(\varepsilon + \frac{1}{A^2} + A^{4n+2} \delta^{\frac{1}{4n+4}} + \varepsilon_0 + \theta_0^{\frac{1}{n+1}} + \kappa_0^{\frac{1}{2}} + A^{2n+2} \tilde{\delta}^{\frac{2}{4n+5}} \right),$$

so that the preceding lemma guarantees that

$$\int_{Q_0} |\mathcal{R}(\chi_{AQ_0} \tilde{\mu})|^2 d\tilde{\mu} \leq \tilde{\varepsilon} \tilde{\mu}(Q_0).$$

We will also need the following auxiliary result below.

Lemma 7.3. *For all $Q \in \text{Stop}_0 \setminus \text{Bad}$, we have*

$$\int_{1.1B_Q \setminus Q} \int_Q \frac{1}{|x-y|^n} d\tilde{\mu}(x) d\tilde{\mu}(y) \lesssim \theta_0^{\frac{1}{2(n+1)^2}} \tilde{\mu}(Q).$$

Proof. Since for any $Q \in \text{Stop}_0 \setminus \text{Bad}$ the ball $1.1B_Q$ is contained in Q_0 , we have that $\tilde{\mu}$ coincides with μ_0 in the above domain of integration.

Let $0 < \kappa < 1$ be some small constant to be fixed below. Then we split

$$(7.9) \quad \begin{aligned} \int_{1.1B_Q \setminus Q} \int_Q \frac{1}{|x-y|^n} d\tilde{\mu}(y) d\tilde{\mu}(x) &= \int_{x \in 1.1B_Q \setminus Q} \int_{y \in Q: |x-y| > \kappa \ell(Q)} \frac{1}{|x-y|^n} d\tilde{\mu}(y) d\tilde{\mu}(x) \\ &\quad + \int_{x \in 1.1B_Q \setminus Q} \int_{y \in Q: |x-y| \leq \kappa \ell(Q)} \frac{1}{|x-y|^n} d\tilde{\mu}(y) d\tilde{\mu}(x). \end{aligned}$$

First we deal with the first integral on the right hand side:

$$\begin{aligned} \int_{x \in 1.1B_Q \setminus Q} \int_{y \in Q: |x-y| > \kappa \ell(Q)} \frac{1}{|x-y|^n} d\tilde{\mu}(y) d\tilde{\mu}(x) &\leq \frac{1}{\kappa^n \ell(Q)^n} \tilde{\mu}(1.1B_Q) \tilde{\mu}(Q) \\ &\lesssim \frac{1}{\kappa^n} \Theta_{\tilde{\mu}}(1.1B_Q) \tilde{\mu}(Q) \lesssim \frac{\theta_0^{\frac{1}{n+1}}}{\kappa^n} \mu(Q), \end{aligned}$$

by Lemma 6.2.

Let us turn our attention to the last integral in (7.9). To estimate this we take into account the fact that given $x \in 1.1B_Q \setminus Q$, if $y \in Q$, then $|x-y| \geq \text{dist}(x, Q)$. Then by the growth of order n of $\mu|_{Q_0}$ and standard estimates, we get

$$\int_{y \in Q: |x-y| \leq \kappa \ell(Q)} \frac{1}{|x-y|^n} d\tilde{\mu}(y) \lesssim \log \left(2 + \frac{\kappa \ell(Q)}{\text{dist}(x, Q)} \right) \quad \text{for all } x \in 1.1B_Q \setminus Q.$$

For each $j \geq 0$, denote

$$U_j = \{x \in 1.1B_Q \setminus Q : \text{dist}(x, Q) \leq 2^{-j} \kappa \ell(Q)\}.$$

By the small boundary property of Q and the fact that Q is doubling,

$$\mu(U_j) \lesssim (2^{-j} \kappa)^{1/2} \mu(3.5B_Q) \lesssim (2^{-j} \kappa)^{1/2} \mu(Q).$$

Then we obtain

$$\begin{aligned}
\int_{x \in 1.1B_Q \setminus Q} \int_{y \in Q: |x-y| \leq \kappa \ell(Q)} \frac{1}{|x-y|^n} d\tilde{\mu}(y) d\tilde{\mu}(x) &\leq \sum_{j \geq 0} \int_{U_j \setminus U_{j+1}} \log \left(2 + \frac{\kappa \ell(Q)}{\text{dist}(x, Q)} \right) d\mu(x) \\
&\lesssim \sum_{j \geq 0} \log \left(2 + \frac{\kappa \ell(Q)}{2^{-j-1} \kappa \ell(Q)} \right) \mu(U_j) \\
&\lesssim \sum_{j \geq 0} (j+1) (2^{-j} \kappa)^{1/2} \mu(Q) \\
&\lesssim \kappa^{1/2} \mu(Q).
\end{aligned}$$

Thus we have

$$\int_{1.1B_Q \setminus Q} \int_Q \frac{1}{|x-y|^n} d\tilde{\mu}(x) d\tilde{\mu}(y) \lesssim (\theta_0^{1/(n+1)} \kappa^{-n} + \kappa^{1/2}) \mu(Q).$$

Choosing $\kappa = \theta_0^{\frac{1}{(n+1)^2}}$, the lemma follows. \square

It is easy to check that

$$(7.10) \quad \tilde{\mu}(B(x, r)) \lesssim r^n \quad \text{for all } x \in \mathbb{R}^{n+1} \text{ and all } r > 0.$$

This follows from the analogous estimate for $\mu|_{Q_0}$ and the periodicity of $\tilde{\mu}$, and is left for the reader. On the other hand, in general, we cannot guarantee that the estimates for the coefficients $P_\mu(2B_Q)$ in Lemma 6.2 also hold with μ replaced by $\tilde{\mu}$. However, we have following substitute.

Lemma 7.4. *The function*

$$p_{\tilde{\mu}}(x) = \sum_{Q \in \text{Stop}_0 \setminus \text{Bad}} \chi_Q P_{\tilde{\mu}}(2B_Q)$$

satisfies

$$\int_{Q_0} p_{\tilde{\mu}}^2 d\tilde{\mu} \lesssim \theta_0^{\frac{2}{3(n+1)}} \tilde{\mu}(Q_0).$$

Proof. Once more, let $0 < \kappa < 1$ be some small constant to be fixed below. We split

$$(7.11) \quad \int_{Q_0} p_{\tilde{\mu}}^2 d\tilde{\mu} = \int_{x \in Q_0: \text{dist}(x, \partial Q_0) \leq \kappa \ell(Q_0)} p_{\tilde{\mu}}(x)^2 d\tilde{\mu}(x) + \int_{x \in Q_0: \text{dist}(x, \partial Q_0) > \kappa \ell(Q_0)} p_{\tilde{\mu}}(x)^2 d\tilde{\mu}(x).$$

For the first integral on the right hand side we just take into account that $p_{\tilde{\mu}}(x) \lesssim 1$ by (7.10), and thus

$$\begin{aligned}
\int_{x \in Q_0: \text{dist}(x, \partial Q_0) \leq \kappa \ell(Q_0)} p_{\tilde{\mu}}(x)^2 d\tilde{\mu}(x) &\lesssim \mu(\{x \in Q_0 : \text{dist}(x, \partial Q_0) \leq \kappa \ell(Q_0)\}) \\
&\lesssim \kappa \mu(Q_0) \approx \kappa \tilde{\mu}(Q_0).
\end{aligned}$$

Let us deal with the the last integral on the right hand side of (7.11). Consider $x \in Q \in \text{Stop}_0$ such that $\text{dist}(x, \partial Q_0) > \kappa \ell(Q_0)$. We assume that $\kappa \gg t = \theta_0^{\frac{1}{(n+1)}}$. Since $\ell(Q) \leq t \ell(Q_0)$,

$$\text{dist}(x, \partial Q_0) \approx \text{dist}(2B_Q, \partial Q_0) \gtrsim \kappa \ell(Q_0).$$

Then, we can write

$$p_{\tilde{\mu}}(x) \lesssim P_{\mu}(2B_Q) + \sum_{j \geq 1: 2^j B_Q \cap \partial Q_0 \neq \emptyset} 2^{-j} \Theta_{\tilde{\mu}}(2^j B_Q) \lesssim \theta_0^{\frac{1}{n+1}} + \sum_{j \geq 1: 2^j B_Q \cap \partial Q_0 \neq \emptyset} 2^{-j},$$

by Lemma 6.2. For the last sum we have

$$\sum_{j \geq 1: 2^j B_Q \cap \partial Q_0 \neq \emptyset} 2^{-j} \approx \frac{\ell(Q)}{\text{dist}(x, \partial Q_0)} \lesssim \frac{t \ell(Q_0)}{\kappa \ell(Q_0)} = \frac{\theta_0^{\frac{1}{n+1}}}{\kappa},$$

and so we obtain

$$p_{\tilde{\mu}}(x) \lesssim \theta_0^{\frac{1}{n+1}} + \frac{\theta_0^{\frac{1}{n+1}}}{\kappa} \approx \frac{\theta_0^{\frac{1}{n+1}}}{\kappa}.$$

Therefore,

$$\int_{x \in Q_0: \text{dist}(x, \partial Q_0) \geq \kappa \ell(Q_0)} p_{\tilde{\mu}}(x)^2 d\tilde{\mu}(x) \lesssim \frac{\theta_0^{\frac{2}{n+1}}}{\kappa^2} \tilde{\mu}(Q_0).$$

Gathering the estimates above, we obtain

$$\int_{Q_0} p_{\tilde{\mu}}^2 d\tilde{\mu} \lesssim \left(\kappa + \frac{\theta_0^{\frac{2}{n+1}}}{\kappa^2} \right) \tilde{\mu}(Q_0).$$

Choosing $\kappa = \theta_0^{\frac{2}{3(n+1)}}$, the lemma follows. \square

8. THE APPROXIMATING MEASURE η

We consider the measure

$$\eta_0 = \sum_{Q \in \text{Stop}_0 \setminus \text{Bad}} \mu_0(Q) \frac{\mathcal{H}^{n+1}|_{\frac{1}{4}B(Q)}}{\mathcal{H}^{n+1}(\frac{1}{4}B(Q))}.$$

So, in a sense, η_0 can be considered as an approximation of $\mu_0|_{Q_0}$ which is absolutely continuous with respect to \mathcal{H}^{n+1} . Further, since the family Stop_0 is finite, the density of η with respect to \mathcal{H}^{n+1} is bounded.

Recall that, by Remark 5.2, the balls $\frac{1}{2}B(Q)$, $Q \in \mathcal{D}_\sigma$, are pairwise disjoint. So the balls $\frac{1}{4}B(Q)$ in the sum above satisfy

$$\text{dist}(\tfrac{1}{4}B(Q), \tfrac{1}{4}B(Q')) \geq \tfrac{1}{4} [r(B(Q)) + r(B(Q'))] \quad \text{if } Q \neq Q'.$$

Now we define the following periodic version of η_0 . Let \mathcal{M} be the lattice of cubes from \mathbb{R}^{n+1} introduced in Section 7. Recall that, for $P \in \mathcal{M}$, z_P stands for the center of P and T_P is the translation $T_P(x) = x + z_P$. We define

$$\eta = \sum_{P \in \mathcal{M}} (T_P)_\# \eta_0.$$

In this way, η can be considered as a kind of approximation of $\tilde{\mu}$.

The following result should be compared to Lemma 7.2.

Lemma 8.1. *We have*

$$\int_{Q_0} |\mathcal{R}(\chi_{AQ_0}\eta)|^2 d\eta \lesssim \varepsilon' \eta(Q_0),$$

where $\varepsilon' = \tilde{\varepsilon} + A^n \kappa_0^{-2n-2} \theta_0^{\frac{1}{(n+1)^2}}$.

Proof. To simplify notation, we denote $\mathcal{S} = \text{Stop}_0 \setminus \text{Bad}$. We consider the function

$$f = \sum_{Q \in \mathcal{S}} m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu})) \chi_Q.$$

It is clear that

$$(8.1) \quad \|f\|_{L^2(\tilde{\mu})}^2 \leq \|\mathcal{R}(\chi_{AQ_0}\tilde{\mu})\|_{L^2(\tilde{\mu}|_{Q_0})}^2 \leq \tilde{\varepsilon} \tilde{\mu}(Q_0) = \tilde{\varepsilon} \eta(Q_0).$$

For all $x \in \frac{1}{4}B(Q)$, $Q \in \mathcal{S}$, we write

$$(8.2) \quad \begin{aligned} |\mathcal{R}(\chi_{AQ_0}\eta)(x)| &\leq |\mathcal{R}(\chi_{\frac{1}{4}B(Q)}\eta)(x)| + |\mathcal{R}(\chi_{AQ_0 \setminus \frac{1}{4}B(Q)}\eta)(x) - \mathcal{R}(\chi_{AQ_0 \setminus Q}\tilde{\mu})(x)| \\ &\quad + |\mathcal{R}(\chi_{AQ_0 \setminus Q}\tilde{\mu})(x) - m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))| + |m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))| \\ &=: T_1 + T_2 + T_3 + |m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))|. \end{aligned}$$

Using that

$$\eta|_{\frac{1}{4}B(Q)} = \tilde{\mu}(Q) \frac{\mathcal{H}^{n+1}|_{\frac{1}{4}B(Q)}}{\mathcal{H}^{n+1}(\frac{1}{4}B(Q))},$$

it follows easily that

$$T_1 = |\mathcal{R}(\chi_{\frac{1}{4}B(Q)}\eta)(x)| \lesssim \frac{\tilde{\mu}(Q)}{r(B(Q))^n} \lesssim \theta_0^{1/(n+1)}.$$

Next we will deal with the term T_3 in (8.2). To this end, for $x \in \frac{1}{4}B(Q)$ we set

$$(8.3) \quad \begin{aligned} &|\mathcal{R}(\chi_{AQ_0 \setminus Q}\tilde{\mu})(x) - m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))| \\ &\leq |\mathcal{R}(\chi_{1.1B_Q \setminus Q}\tilde{\mu})(x)| + |\mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu})(x) - m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu}))| \\ &\quad + |m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{1.1B_Q \setminus Q}\tilde{\mu}))|, \end{aligned}$$

taking into account that $m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_Q\tilde{\mu})) = 0$, by the antisymmetry of the Riesz kernel. The first term on the right hand side satisfies

$$|\mathcal{R}(\chi_{1.1B_Q \setminus Q}\tilde{\mu})(x)| \leq \int_{1.1B_Q \setminus Q} \frac{1}{|x-y|^n} d\tilde{\mu}(y) \lesssim \frac{\tilde{\mu}(1.1B_Q)}{r(B(Q))^n} \lesssim \theta_0^{1/(n+1)},$$

recalling that $x \in \frac{1}{4}B(Q)$ and that $\Theta_\mu(1.1B_Q) \lesssim \theta_0^{1/(n+1)}$ for the last estimate.

Now we turn our attention to the second term on the right hand side of (8.3). For $x' \in Q \in \mathcal{S}$, we have

$$\begin{aligned} |\mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu})(x) - \mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu})(x')| &\leq \int_{AQ_0 \setminus 1.1B_Q} |K(x-y) - K(x'-y)| d\tilde{\mu}(y) \\ &\lesssim P_{\tilde{\mu}}(2B_Q). \end{aligned}$$

taking into account that the distance both from x and x' to $(1.1B_Q)^c$ is larger than $cr(B_Q)$. Averaging on $x' \in Q$ with respect to $\tilde{\mu}$ we get

$$|\mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu})(x) - m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0 \setminus 1.1B_Q}\tilde{\mu}))| \lesssim P_{\tilde{\mu}}(2B_Q).$$

To estimate the last term in (8.3) we just apply Lemma 7.3:

$$|m_{\tilde{\mu},Q}(\mathcal{R}(\chi_{1.1B_Q \setminus Q} \tilde{\mu}))| \leq \frac{1}{\tilde{\mu}(Q)} \int_{1.1B_Q \setminus Q} \int_Q \frac{1}{|x-y|^n} d\tilde{\mu}(x) d\tilde{\mu}(y) \lesssim \theta_0^{\frac{1}{2(n+1)^2}}.$$

Thus, we obtain

(8.4)

$$T_3 = |\mathcal{R}(\chi_{AQ_0 \setminus Q} \tilde{\mu})(x) - m_{\tilde{\mu},Q}(\mathcal{R}(\chi_{AQ_0} \tilde{\mu}))| \lesssim \theta_0^{\frac{1}{n+1}} + P_{\tilde{\mu}}(2B_Q) + \theta_0^{\frac{1}{2(n+1)^2}} \lesssim \theta_0^{\frac{1}{2(n+1)^2}} + P_{\tilde{\mu}}(2B_Q).$$

To deal with the term T_2 in (8.2) we need to introduce some additional notation. We set

$$J = \bigcup_{P \in \mathcal{M}} \{T_P(R') : R' \in \text{Stop}_0 \setminus \text{Bad}\}.$$

For $R \in J$ such that $R = T_P(R')$, $R' \in \text{Stop}_0 \setminus \text{Bad}$, we set $B(R) = T_P(B(R'))$ and $B_R = T_P(B_{R'})$. Also, we denote by J_A the family of cells $R \in J$ which are contained in AQ_0 . In this way, we have

$$\chi_{AQ_0} \tilde{\mu} = \sum_{R \in J_A} \tilde{\mu}|_R, \quad \text{and} \quad \chi_{AQ_0} \eta = \sum_{R \in J_A} \tilde{\mu}(R) \frac{\mathcal{H}^{n+1}|_{\frac{1}{4}B(R)}}{\mathcal{H}^{n+1}(\frac{1}{4}B(R))}$$

(recall that we assume A to be a big odd natural number). Note that the cells $R \in J$ are pairwise disjoint. Further, by the definition of the family Bad , if $R \in J$ is contained in some cube $T_P(Q_0)$, then the ball $1.1B_R$ is also contained in $T_P(Q_0)$. Together with the doubling property of the cells from Stop_0 this guarantees that, for all $R \in J$,

$$\tilde{\mu}(1.1B_R) \lesssim C_0 \tilde{\mu}(R).$$

Now for $x \in \frac{1}{4}B(Q)$ we write

$$\begin{aligned} (8.5) \quad T_2 &= |\mathcal{R}(\chi_{AQ_0 \setminus \frac{1}{4}B(Q)} \eta)(x) - \mathcal{R}(\chi_{AQ_0 \setminus Q} \tilde{\mu})(x)| \\ &\leq \sum_{R \in J_A : R \neq Q} \left| \int K(x-y) d(\eta_{\frac{1}{4}B(R)} - \tilde{\mu}|_R) \right| \\ &\leq \sum_{R \in J_A : R \neq Q} \int |K(x-y) - K(x-z_R)| d(\eta_{\frac{1}{4}B(R)} + \tilde{\mu}|_R), \end{aligned}$$

using that $\eta(\frac{1}{4}B(R)) = \tilde{\mu}(R)$ for the last inequality.

We claim that, for $x \in \frac{1}{4}B(Q)$ and $y \in \frac{1}{4}B(R) \cup \text{supp}(\tilde{\mu}|_R)$,

$$(8.6) \quad |K(x-y) - K(x-z_R)| \lesssim \frac{\ell(R)}{\kappa_0^{n+1} D(Q, R)^{n+1}},$$

where

$$D(Q, R) = \ell(Q) + \ell(R) + \text{dist}(Q, R).$$

To show (8.6) note first that

$$(8.7) \quad x \in \frac{1}{4}B(Q), \quad z_R \in \frac{1}{4}B(R) \quad \Rightarrow \quad |x - z_R| \gtrsim D(Q, R),$$

since $\frac{1}{2}B(Q) \cap \frac{1}{2}B(R) = \emptyset$. Analogously, because of the same reason,

$$(8.8) \quad x \in \frac{1}{4}B(Q), \quad y \in \frac{1}{4}B(R) \quad \Rightarrow \quad |x - y| \gtrsim D(Q, R).$$

Also,

$$(8.9) \quad x \in \frac{1}{4}B(Q), \quad y \in \text{supp}(\tilde{\mu}|_R) \quad \Rightarrow \quad |x - y| \gtrsim \kappa_0 D(Q, R),$$

To prove this, note that

$$(8.10) \quad y \in \text{supp}(\tilde{\mu}|_R) = I_{\kappa_0}(R) \subset R,$$

which implies that $y \notin B(Q)$ and thus $|x - y| \geq \frac{3}{4}r(B(Q)) \approx \ell(Q)$. In the case $r(B(Q)) \geq 2\kappa_0\ell(R)$ this implies that

$$|x - y| \gtrsim \ell(Q) + \kappa_0\ell(R).$$

Otherwise, from (8.10), since $z_Q \in \text{supp} \tilde{\mu}$ and $y \in I_{\kappa_0}(R)$, by the definition of $I_{\kappa_0}(R)$,

$$|z_Q - y| \geq \kappa_0\ell(R),$$

and then, as $|z_Q - x| \leq \frac{1}{4}r(B(Q)) \leq \frac{1}{2}\kappa_0\ell(R)$, we infer that

$$|x - y| \geq |z_Q - y| - |z_Q - x| \geq \frac{\kappa_0}{2}\ell(R).$$

So in any case we have $|x - y| \gtrsim \kappa_0(\ell(Q) + \ell(R))$. It is easy to deduce (8.9) from this estimate. We leave the details for the reader.

From (8.7), (8.8), and (8.9), and the fact that $K(\cdot)$ is a standard Calderón-Zygmund kernel, we get (8.6). Plugging this estimate into (8.5), we obtain

$$T_2 \lesssim \frac{1}{\kappa_0^{n+1}} \sum_{R \in J_A} \frac{\ell(R) \tilde{\mu}(R)}{D(Q, R)^{n+1}}.$$

So from (8.2) and the estimates for the terms T_1, T_2 and T_3 , we infer that for all $x \in \frac{1}{4}B(Q)$ with $Q \in \mathcal{S}$

$$(8.11) \quad |\mathcal{R}(\chi_{AQ_0}\eta)(x)| \lesssim |m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))| + \theta_0^{\frac{1}{2(n+1)^2}} + P_{\tilde{\mu}}(2B_Q) + \frac{1}{\kappa_0^{n+1}} \sum_{R \in J_A} \frac{\ell(R) \tilde{\mu}(R)}{D(Q, R)^{n+1}}.$$

Denote

$$\tilde{p}_{\tilde{\mu}}(x) = \sum_{Q \in J} \chi_{\frac{1}{4}B(Q)} P_{\tilde{\mu}}(2B_Q) \quad \text{and} \quad \tilde{g}(x) = \sum_{Q \in \mathcal{S}} \sum_{R \in J_A} \frac{\ell(R)}{D(Q, R)^{n+1}} \tilde{\mu}(R) \chi_{\frac{1}{4}B(Q)}(x).$$

Squaring and integrating (8.11) with respect to η on Q_0 , we get

$$(8.12) \quad \begin{aligned} \|\mathcal{R}(\chi_{AQ_0}\eta)\|_{L^2(\eta|_{Q_0})}^2 &\lesssim \sum_{Q \in \mathcal{S}} |m_{\tilde{\mu}, Q}(\mathcal{R}(\chi_{AQ_0}\tilde{\mu}))|^2 \eta(\tfrac{1}{4}B(Q)) \\ &\quad + \theta_0^{\frac{1}{(n+1)^2}} \eta(Q_0) + \|\tilde{p}_{\tilde{\mu}}\|_{L^2(\eta|_{Q_0})}^2 + \frac{1}{\kappa_0^{2n+2}} \|\tilde{g}\|_{L^2(\eta|_{Q_0})}^2, \end{aligned}$$

Note that, since $\eta(\frac{1}{4}B(Q)) = \tilde{\mu}(Q)$, the first sum on the right hand side of (8.12) equals $\|f\|_{L^2(\tilde{\mu})}^2$, which does not exceed $\tilde{\varepsilon}\eta(Q_0)$, by (8.1). By an analogous argument we deduce that $\|\tilde{p}_{\tilde{\mu}}\|_{L^2(\eta|_{Q_0})}^2 = \|p_{\tilde{\mu}}\|_{L^2(\tilde{\mu}|_{Q_0})}^2$ and $\|\tilde{g}\|_{L^2(\eta|_{Q_0})}^2 = \|g\|_{L^2(\tilde{\mu}|_{Q_0})}^2$, where

$$p_{\tilde{\mu}}(x) = \sum_{Q \in J} \chi_Q P_{\tilde{\mu}}(2B_Q) \quad \text{and} \quad g(x) = \sum_{Q \in \mathcal{S}} \sum_{R \in J_A} \frac{\ell(R)}{D(Q, R)^{n+1}} \tilde{\mu}(R) \chi_Q(x).$$

We will estimate $\|g\|_{L^2(\tilde{\mu}|_{Q_0})}$ by duality: for any non-negative function $h \in L^2(\tilde{\mu}|_{Q_0})$, we set

$$(8.13) \quad \int g h d\tilde{\mu} = \sum_{Q \in \mathcal{S}} \sum_{R \in J_A} \frac{\ell(R)}{D(Q, R)^{n+1}} \tilde{\mu}(R) \int_Q h d\tilde{\mu} = \sum_{R \in J_A} \tilde{\mu}(R) \sum_{Q \in \mathcal{S}} \frac{\ell(R)}{D(Q, R)^{n+1}} \int_Q h d\tilde{\mu}.$$

For each $z \in R \in J_A$ we have

$$\begin{aligned} \sum_{Q \in \mathcal{S}} \frac{\ell(R)}{D(Q, R)^{n+1}} \int_Q h d\tilde{\mu} &\lesssim \int \frac{\ell(R) h(y)}{(\ell(R) + |z - y|)^{n+1}} d\tilde{\mu}(y) \\ &= \int_{|z-y| \leq \ell(R)} \cdots + \sum_{j \geq 1} \int_{2^{j-1}\ell(R) < |z-y| \leq 2^j\ell(R)} \cdots \\ &\lesssim \sum_{j \geq 0} \int_{B(z, 2^j\ell(R))} h d\tilde{\mu} \frac{2^{-j} \tilde{\mu}(B(z, 2^j\ell(R)))}{(2^j\ell(R))^n} \\ &\lesssim M_{\tilde{\mu}} h(z) P_{\tilde{\mu}}(B(z, \ell(R))), \end{aligned}$$

where $M_{\tilde{\mu}}$ stands for the centered maximal Hardy-Littlewood operator with respect to $\tilde{\mu}$. Then, by (8.13),

$$\begin{aligned} \int g h d\tilde{\mu} &\lesssim \sum_{R \in J_A} \inf_{z \in R} [M_{\tilde{\mu}} h(z) P_{\tilde{\mu}}(B(z, \ell(R)))] \tilde{\mu}(R) \leq \int_{AQ_0} M_{\tilde{\mu}} h p_{\tilde{\mu}} d\tilde{\mu} \\ &\lesssim \|M_{\tilde{\mu}} h\|_{L^2(\tilde{\mu})} \|p_{\tilde{\mu}}\|_{L^2(\tilde{\mu}|_{AQ_0})} \lesssim \|h\|_{L^2(\tilde{\mu})} \|p_{\tilde{\mu}}\|_{L^2(\tilde{\mu}|_{AQ_0})}. \end{aligned}$$

Thus, by Lemma 7.4 and recalling that $\tilde{\mu}$ is \mathcal{M} -periodic,

$$\|g\|_{L^2(\tilde{\mu}|_{Q_0})}^2 \lesssim \|p_{\tilde{\mu}}\|_{L^2(\tilde{\mu}|_{AQ_0})}^2 = A^n \|p_{\tilde{\mu}}\|_{L^2(\tilde{\mu}|_{Q_0})}^2 \lesssim A^n \theta_0^{\frac{2}{3(n+1)}} \tilde{\mu}(Q_0).$$

Plugging this estimate into (8.12) and recalling that $\|g\|_{L^2(\eta|_{Q_0})} = \|g\|_{L^2(\tilde{\mu})}$, we obtain

$$\|\mathcal{R}(\chi_{AQ_0}\eta)\|_{L^2(\eta|_{Q_0})}^2 \lesssim \left(\tilde{\varepsilon} + \theta_0^{\frac{1}{(n+1)^2}} + \frac{A^n}{\kappa_0^{2n+2}} \theta_0^{\frac{2}{3(n+1)}} \right) \eta(Q_0) \lesssim \left(\tilde{\varepsilon} + \frac{A^n}{\kappa_0^{2n+2}} \theta_0^{\frac{1}{(n+1)^2}} \right) \eta(Q_0),$$

as wished. \square

Note that the Riesz kernel is locally integrable with respect to η (recall that the number of cells from Stop_0 is finite). So for any bounded function f with compact support the integral $\int K(x - y) f(y) d\eta(y)$ is absolutely convergent for all $x \in \mathbb{R}^{n+1}$.

Now we wish to extend the definition of $\mathcal{R}_\eta f(x)$ to \mathcal{M} -periodic functions $f \in L^\infty(\eta)$ in a pointwise way (not only in a BMO sense, say). We consider a non-negative radial C^1 function ϕ supported on $B(0, 2)$ which equals 1 on $B(0, 1)$, and we set $\phi_r(x) = \phi(\frac{x}{r})$ for $r > 0$. We denote $\tilde{K}_r(x - y) = K(x - y)\phi_r(x - y)$ and we define

$$\tilde{\mathcal{R}}_{\eta,r} f(x) = \tilde{\mathcal{R}}_r(f\eta)(x) = \int \tilde{K}_r(x - y) f(y) d\eta(y),$$

and

$$(8.14) \quad \text{pv}\mathcal{R}_\eta f(x) = \text{pv}\mathcal{R}(f\eta)(x) = \lim_{r \rightarrow \infty} \tilde{\mathcal{R}}_{\eta,r} f(x),$$

whenever the limit exists. Let us remark that one may also define the principal value in a more typical way by

$$(8.15) \quad \lim_{r \rightarrow \infty} \int_{|x-y| < r} K(x-y) f(y) d\eta(y).$$

However, the definition (8.14) has some technical advantages and simplifies the exposition below. Nevertheless, one can show that both definitions (8.14) and (8.15) coincide, at least for the \mathcal{M} -periodic functions $f \in L^\infty(\eta)$ (we will not prove this fact because this will be not needed below).

Lemma 8.2. *Let $f \in L^\infty(\eta)$ be \mathcal{M} -periodic, that is, $f(x + z_P) = f(x)$ for all $x \in \mathbb{R}^{n+1}$ and all $P \in \mathcal{M}$. Then:*

- $\text{pv}\mathcal{R}_\eta f(x)$ exists for all $x \in \mathbb{R}^{n+1}$ and $\tilde{\mathcal{R}}_{\eta,r} f \rightarrow \text{pv}\mathcal{R}_\eta f$ as $r \rightarrow \infty$ uniformly in compact subsets of \mathbb{R}^{n+1} . The convergence is also uniform on $\text{supp } \eta$. Further, given any compact set $F \subset \mathbb{R}^{n+1}$, there is $r_0 = r_0(F) > 0$ such that for $s > r \geq r_0$,

$$(8.16) \quad \|\tilde{\mathcal{R}}_s(f\eta) - \tilde{\mathcal{R}}_r(f\eta)\|_{\infty, F} \lesssim \frac{c_F}{r} \|f\|_\infty,$$

where c_F is some constant depending on F .

- The function $\text{pv}\mathcal{R}_\eta f$ is \mathcal{M} -periodic and continuous in \mathbb{R}^{n+1} , and harmonic in $\mathbb{R}^{n+1} \setminus \text{supp}(f\eta)$.

The arguments to prove the lemma are standard. However, for the reader's convenience we will show the details.

Proof. By the \mathcal{M} -periodicity of the measure $\nu := f\eta$, it is immediate that the functions $\tilde{\mathcal{R}}_r(f\eta)$, $r > 0$, are \mathcal{M} -periodic too. On the other hand, using that η is absolutely continuous with respect to Lebesgue measure with a uniformly bounded density, it is straightforward to check that each $\tilde{\mathcal{R}}_r(f\eta)$ is also continuous and bounded in \mathbb{R}^{n+1} . Then, except harmonicity in $\mathbb{R}^{n+1} \setminus \text{supp}(f\eta)$, all the statements in the lemma follow if we show that the family of functions $\{\tilde{\mathcal{R}}_r(f\eta)\}_{r>0}$ satisfies (8.16) for any compact subset $F \subset \mathbb{R}^{n+1}$. Indeed this clearly implies the uniform convergence on compact subsets and on $\text{supp } \eta$, since $\text{supp } \eta$ is \mathcal{M} -periodic,

Let $s > r \geq r_0$ and, denote $\tilde{K}_{r,s}(x-y) = \tilde{K}_s(x-y) - \tilde{K}_r(x-y)$. Notice that $\tilde{K}_{r,s}$ is a standard Calderón-Zygmund kernel (with constants independent of r and s). We write

$$\nu = \sum_{P \in \mathcal{M}} (T_P)_\# (\chi_{Q_0} \nu),$$

so that

$$\tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) = \int \tilde{K}_{r,s}(x-y) d\left(\sum_{P \in \mathcal{M}} (T_P)_\# (\chi_{Q_0} \nu)\right)(y).$$

Since the support of $\tilde{K}_{r,s}(x-y)$ is compact, the last sum only has a finite number of non-zero terms, and so we can change the order of summation and integration:

$$(8.17) \quad \begin{aligned} \tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) &= \sum_{P \in \mathcal{M}} \int \tilde{K}_{r,s}(x-y) d[(T_P)_\# (\chi_{Q_0} \nu)](y) \\ &= \sum_{P \in \mathcal{M}} \int_{Q_0} \tilde{K}_{r,s}(x-y-z_P) d\nu(y). \end{aligned}$$

By the antisymmetry of the kernel $\tilde{K}_{r,s}$, from the last equation we derive

$$\tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) = - \sum_{P \in \mathcal{M}} \int_{Q_0} \tilde{K}_{r,s}(z_P - (x - y)) d\nu(y).$$

Also, by the definition of \mathcal{M} , it is clear that $P \in \mathcal{M}$ if and only if $-P \in \mathcal{M}$. So replacing z_P by $-z_P$ does not change the last sum in (8.17). Hence we have

$$\tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) = \sum_{P \in \mathcal{M}} \int_{Q_0} \tilde{K}_{r,s}(z_P + (x - y)) d\nu(y).$$

Averaging the last two equations we get

$$(8.18) \quad \tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) = \frac{1}{2} \sum_{P \in \mathcal{M}} \int_{Q_0} [\tilde{K}_{r,s}(z_P + (x - y)) - \tilde{K}_{r,s}(z_P - (x - y))] d\nu(y).$$

Note that if x belongs to a compact set $F \subset \mathbb{R}^{n+1}$ and $y \in Q_0$, then both $(x - y)$ and $-(x - y)$ lie in some compact set \tilde{F} . Observe also that $\tilde{K}_{r,s}$ vanishes in $B(0, r)$. So if we assume $r_0 \geq 2 \operatorname{diam}(\tilde{F})$, say, then both $\tilde{K}_{r,s}(z_P + (x - y))$ and $\tilde{K}_{r,s}(z_P - (x - y))$ vanish unless $|z_P| \geq r$. For such x, y we have $|x - y| \leq \operatorname{diam}(\tilde{F}) \leq \frac{1}{2}r \leq \frac{1}{2}|z_P|$ and so

$$|z_P + (x - y)| \approx |z_P - (x - y)| \approx |z_P| \geq r.$$

Then we obtain

$$|\tilde{K}_{r,s}(z_P + (x - y)) - \tilde{K}_{r,s}(z_P - (x - y))| \lesssim \frac{|x - y|}{|z_P|^{n+1}} \lesssim \frac{\operatorname{diam}(\tilde{F})}{|z_P|^{n+1}}.$$

Plugging this estimate into (8.18) we obtain

$$|\tilde{\mathcal{R}}_s(f\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x)| \lesssim \sum_{P \in \mathcal{M}: |z_P| \geq r} \frac{\operatorname{diam}(\tilde{F})}{|z_P|^{n+1}} |\nu|(Q_0) \leq \sum_{P \in \mathcal{M}: |z_P| \geq r} \frac{\operatorname{diam}(\tilde{F})}{|z_P|^{n+1}} \ell(P)^n \|f\|_\infty.$$

It is easy to check that

$$\sum_{P \in \mathcal{M}: |z_P| \geq r} \frac{\ell(P)^n}{|z_P|^{n+1}} \lesssim \frac{1}{r}.$$

So we deduce

$$\|\tilde{\mathcal{R}}_s(f\eta) - \tilde{\mathcal{R}}_r(f\eta)\|_{\infty, F} \lesssim \frac{\operatorname{diam}(\tilde{F})}{r} \|f\|_\infty \rightarrow 0 \quad \text{as } r \rightarrow \infty,$$

as wished.

It remains to prove that $\operatorname{pv}\mathcal{R}_\nu f$ is harmonic in $\mathbb{R}^{n+1} \setminus \operatorname{supp}(f\eta)$. Consider a closed ball $\overline{B}(0, r_1)$ and $x \in \overline{B}(0, r_1)$. Then we have

$$\mathcal{R}(f\phi_r\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x) = \int K(x - y) (\phi_r(y) - \phi_r(x - y)) f(y) d\eta(y).$$

We write

$$|\phi_r(y) - \phi_r(x - y)| \lesssim \|\nabla \phi_r\|_\infty |x| \lesssim \frac{|x|}{r}.$$

For $r \geq 4r_1$, it is easy to check that $\phi_r(y) - \phi_r(x - y) = 0$ unless $|x - y| \approx |y| \approx r$. Thus

$$\begin{aligned} |\mathcal{R}(f\phi_r\eta)(x) - \tilde{\mathcal{R}}_r(f\eta)(x)| &\lesssim \int_{\substack{|y| \leq Cr \\ C^{-1}r \leq |x-y| \leq Cr}} \frac{1}{|x-y|^n} \frac{|x|}{r} |f(y)| d\eta(y) \\ &\lesssim \frac{|x|}{r^{n+1}} \|f\|_\infty \eta(B(0, Cr)) \lesssim \frac{r_1}{r} \|f\|_\infty. \end{aligned}$$

That is to say,

$$\|\mathcal{R}(f\phi_r\eta) - \tilde{\mathcal{R}}_r(f\eta)\|_{\infty, \overline{B}(0, r_1)} \lesssim \frac{r_1}{r} \|f\|_\infty \rightarrow 0 \quad \text{as } r \rightarrow \infty.$$

Since $\tilde{\mathcal{R}}_r(f\eta)$ converges uniformly to $\text{pv}\mathcal{R}_\eta f$ in F as $r \rightarrow \infty$, it follows that $\mathcal{R}(f\phi_r\eta)$ also converges uniformly to $\text{pv}\mathcal{R}_\eta f$ in $\overline{B}(0, r_1)$.

Note now that $\mathcal{R}(f\phi_r\eta)$ is harmonic out of $\text{supp}(f\eta)$ because $f\phi_r\eta$ has compact support, and so by its local uniform convergence to $\text{pv}\mathcal{R}_\eta f$, we deduce that $\text{pv}\mathcal{R}_\eta f$ is harmonic out of $\text{supp}(f\eta)$ too. \square

From now on, to simplify notation we will denote $\text{pv}\mathcal{R}_\eta f$ just by $\mathcal{R}_\eta f$.

Lemma 8.3. *Let $L_{\mathcal{M}}^\infty(\eta)$ denote the Banach space of the \mathcal{M} -periodic functions which belong to $L^\infty(\eta)$ equipped with the norm $\|\cdot\|_{L^\infty(\eta)}$. The map $\mathcal{R}_\eta : L_{\mathcal{M}}^\infty(\eta) \rightarrow L_{\mathcal{M}}^\infty(\eta)$ is bounded. Further, for all $f \in L_{\mathcal{M}}^\infty(\eta)$ and $r > 0$ big enough we have*

$$(8.19) \quad \|\mathcal{R}(f\eta) - \tilde{\mathcal{R}}_r(f\eta)\|_{L^\infty(\eta)} \lesssim \frac{\|f\|_{L^\infty(\eta)}}{r}.$$

We remark that the bound on the norm of \mathcal{R}_η from $L_{\mathcal{M}}^\infty(\eta)$ to $L_{\mathcal{M}}^\infty(\eta)$ depends strongly on the construction of η . This is finite due to the fact that the number of cells from Stop_0 is finite, but it may explode as this number grows. The precise value of the norm will not play any role in the estimates below, we just need to know that this is finite.

Proof. Since f is \mathcal{M} -periodic, from (8.16) we infer that for $s > r \geq r_0 = r_0(Q_0)$,

$$\|\tilde{\mathcal{R}}_s(f\eta) - \tilde{\mathcal{R}}_r(f\eta)\|_{\infty, F} \lesssim \frac{c_F}{r} \|f\|_\infty,$$

Letting $s \rightarrow \infty$, $\tilde{\mathcal{R}}_s(f\eta)$ converges uniformly to $\mathcal{R}_\eta f$ and so we get (8.19).

To prove the boundedness of $\mathcal{R}_\eta : L_{\mathcal{M}}^\infty(\eta) \rightarrow L_{\mathcal{M}}^\infty(\eta)$, note first that \tilde{K}_{r_0} is compactly supported and η is absolutely continuous with respect to Lebesgue measure on a compact set with a uniformly bounded density. Hence we deduce that $\tilde{\mathcal{R}}_{\eta, r_0} : L_{\mathcal{M}}^\infty(\eta) \rightarrow L_{\mathcal{M}}^\infty(\eta)$ is bounded, which together with (8.19) applied to $\tilde{\mathcal{R}}_{r_0}$ implies that $\mathcal{R}_\eta : L_{\mathcal{M}}^\infty(\eta) \rightarrow L_{\mathcal{M}}^\infty(\eta)$ is bounded. \square

From now on, given $x \in \mathbb{R}^{n+1}$, we denote

$$x_H = (x_1, \dots, x_n),$$

so that $x = (x_H, x_{n+1})$. Also, we write

$$\mathcal{R}^H = (\mathcal{R}_1, \dots, \mathcal{R}_n),$$

where \mathcal{R}_j stands for the j -th component of \mathcal{R} , so that $\mathcal{R} = (\mathcal{R}^H, \mathcal{R}_{n+1})$.

For simplicity, in the arguments below we will assume that the function ϕ defined slightly above (8.14) is of the form $\phi(x) = \tilde{\phi}(|x|^2)$, for some even C^1 function $\tilde{\phi}$ which equals 1 on $[0, 1]$ and vanishes out of $[0, 2^{1/2}]$.

Lemma 8.4. *Let $f \in L^1_{loc}(\eta)$ be \mathcal{M} -periodic. Then,*

(a) *Let $\tilde{A} \geq 5$ be some odd natural number. For all $x \in 2Q_0$,*

$$|\mathcal{R}(\chi_{(\tilde{A}Q_0)^c} f \eta)(x)| \lesssim \frac{1}{\tilde{A} \ell(Q_0)^n} \int_{Q_0} |f| d\eta.$$

(b) *For all $x \in \mathbb{R}^{n+1}$ such that $\text{dist}(x, H) \geq \ell(Q_0)$,*

$$(8.20) \quad |\mathcal{R}(f \eta)(x)| \lesssim \frac{1}{\ell(Q_0)^n} \int_{Q_0} |f| d\eta$$

and

$$(8.21) \quad |\mathcal{R}^H(f \eta)(x)| \lesssim \frac{1}{\text{dist}(x, H) \ell(Q_0)^{n-1}} \int_{Q_0} |f| d\eta$$

Proof. We denote $\nu = f \eta$. The arguments to prove the estimate in (a) are quite similar to the ones used in the proof of Lemma 8.2. Since we are assuming that \tilde{A} is some odd number, there is a subset $\mathcal{M}_{\tilde{A}} \subset \mathcal{M}$ such that

$$\chi_{(\tilde{A}Q_0)^c} \nu = \sum_{P \in \mathcal{M}_{\tilde{A}}} (T_P)_\# (\chi_{Q_0} \nu).$$

Further the cubes $P \in \mathcal{M}_{\tilde{A}}$ satisfy $|z_P| \gtrsim \tilde{A} \ell(Q_0)$. So for any $x \in Q_0$ and all $r > 0$ we have

$$\tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c} \nu)(x) = \int \tilde{K}_r(x - y) d\left(\sum_{P \in \mathcal{M}_{\tilde{A}}} (T_P)_\# (\chi_{Q_0} \nu) \right)(y).$$

Since the support of $\tilde{K}_r(x - y)$ is compact, the last sum only has a finite number of non-zero terms, and so we can change the order of summation and integration, and thus

$$(8.22) \quad \begin{aligned} \tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c} \nu)(x) &= \sum_{P \in \mathcal{M}_{\tilde{A}}} \int \tilde{K}_r(x - y) d[(T_P)_\# (\chi_{Q_0} \nu)](y) \\ &= \sum_{P \in \mathcal{M}_{\tilde{A}}} \int_{Q_0} \tilde{K}_r(x - y - z_P) d\nu(y). \end{aligned}$$

By the antisymmetry of the kernel \tilde{K}_r , from the last equation we get

$$\tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c} \nu)(x) = - \sum_{P \in \mathcal{M}_{\tilde{A}}} \int_{Q_0} \tilde{K}_r(z_P - (x - y)) d\nu(y).$$

Also, by the definition of $\mathcal{M}_{\tilde{A}}$, it follows that $P \in \mathcal{M}_{\tilde{A}}$ if and only if $-P \in \mathcal{M}_{\tilde{A}}$. So replacing z_P by $-z_P$ does not change the last sum in (8.22), and then we have

$$\tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c} \nu)(x) = \sum_{P \in \mathcal{M}_{\tilde{A}}} \int_{Q_0} \tilde{K}_r(z_P + (x - y)) d\nu(y).$$

Averaging the last two equations we get

$$(8.23) \quad \tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c}\nu)(x) = \frac{1}{2} \sum_{P \in \mathcal{M}_{\tilde{A}}} \int_{Q_0} [\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))] d\nu(y).$$

Note now that $x \in 2Q_0$, $y \in Q_0$ and, recalling that $|z_P| \gtrsim \tilde{A}\ell(Q_0)$ for $P \in \mathcal{M}_{\tilde{A}}$, we have

$$|z_P + (x - y)| \approx |z_P - (x - y)| \approx |z_P|.$$

Thus,

$$|\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{|x - y|}{|z_P|^{n+1}} \lesssim \frac{\ell(Q_0)}{|z_P|^{n+1}}.$$

Then, from this estimate and (8.23) we deduce that

$$|\tilde{\mathcal{R}}_r(\chi_{(\tilde{A}Q_0)^c}\nu)(x)| \lesssim \sum_{P \in \mathcal{M}: |z_P| \geq C^{-1}\tilde{A}\ell(Q_0)} \frac{\ell(Q_0)}{|z_P|^{n+1}} |\nu|(Q_0) \lesssim \frac{|\nu|(Q_0)}{\tilde{A}\ell(Q_0)^n}.$$

as wished.

To prove the first estimate in (b), let $x \in \mathbb{R}^{n+1}$ be such that $\text{dist}(x, H) \geq \ell(Q_0)$. Since $\mathcal{R}\nu$ is \mathcal{M} -periodic, we may assume that $x_H \in Q_0 \cap H$. As in (8.23), for any $r > 0$ we have

$$(8.24) \quad \tilde{\mathcal{R}}_r\nu(x) = \frac{1}{2} \sum_{P \in \mathcal{M}} \int_{Q_0} [\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))] d\nu(y).$$

We claim that for x as above and $y \in Q_0$,

$$(8.25) \quad |\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{\text{dist}(x, H)}{(\text{dist}(x, H) + |z_P|)^{n+1}}.$$

Indeed, if $|z_P| \geq 2|x - y|$, then $\text{dist}(x, H) + |x - z_P| \approx |x - y| + |z_P| \approx |z_P|$, and thus

$$|\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{|x - y|}{|z_P|^{n+1}} \approx \frac{\text{dist}(x, H)}{|z_P|^{n+1}} \approx \frac{\text{dist}(x, H)}{(\text{dist}(x, H) + |z_P|)^{n+1}},$$

which shows that (8.25) holds in this case.

On the other hand, if $|z_P| < 2|x - y|$, then

$$|\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{1}{|(z_P - y) + x|^n} + \frac{1}{|(z_P + y) - x|^n}.$$

It is immediate to check that $\text{dist}(x, y - z_P) \approx \text{dist}(x, z_P + y) \approx \text{dist}(x, H) \approx |x - y|$, and so we deduce that

$$|\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{1}{|x - y|^n}.$$

Further from the condition $|z_P| < 2|x - y|$ we infer that

$$|x - y| \approx |x - y| + |z_P| \approx \text{dist}(x, H) + |z_P|,$$

and thus

$$|\tilde{K}_r(z_P + (x - y)) - \tilde{K}_r(z_P - (x - y))| \lesssim \frac{|x - y|}{(|x - y| + |z_P|)^{n+1}} \approx \frac{\text{dist}(x, H)}{(\text{dist}(x, H) + |z_P|)^{n+1}},$$

which completes the proof of (8.25).

From (8.24) and (8.25) we deduce

$$\begin{aligned} |\tilde{\mathcal{R}}_r \nu(x)| &\lesssim \sum_{P \in \mathcal{M}} \int_{Q_0} \frac{\text{dist}(x, H)}{(\text{dist}(x, H) + |z_P|)^{n+1}} d|\nu|(y) \\ &= |\nu|(Q_0) \frac{\text{dist}(x, H)}{\ell(P)^n} \sum_{P \in \mathcal{M}} \frac{\ell(P)^n}{(\text{dist}(x, H) + |z_P|)^{n+1}} \end{aligned}$$

It is straightforward to check that

$$\sum_{P \in \mathcal{M}} \frac{\ell(P)^n}{(\text{dist}(x, H) + |z_P|)^{n+1}} \lesssim \frac{1}{\text{dist}(x, H)},$$

and thus (8.20) follows.

We turn now our attention to the last estimate in (b). Again let $x \in \mathbb{R}^{n+1}$ be such that $\text{dist}(x, H) \geq \ell(Q_0)$ and $x_H \in Q_0 \cap H$, so that the identity (8.24) is still valid. We claim that for $y \in Q_0$ and r big enough,

$$(8.26) \quad |\tilde{K}_r^H(z_P + (x - y)) - \tilde{K}_r^H(z_P - (x - y))| \lesssim \frac{\ell(Q_0)}{(\text{dist}(x, H) + |z_P|)^{n+1}},$$

where \tilde{K}_r^H is the kernel of $\tilde{\mathcal{R}}_r^H$. To prove this observe that

$$\tilde{K}_r^H(z) = z_H \psi_r(|z|^2), \quad \text{with } \psi_r(t) = \frac{\tilde{\phi}_r(t)}{t^{\frac{n+1}{2}}}.$$

Then we have

$$\begin{aligned} &|\tilde{K}_r^H(z_P + (x - y)) - \tilde{K}_r^H(z_P - (x - y))| \\ &= |((z_{P,H} + (x_H - y_H)) \psi_r(|z_P + (x - y)|^2) - (z_{P,H} - (x_H - y_H)) \psi_r(|z_P - (x - y)|^2))| \\ &\leq 2|x_H - y_H| \psi_r(|z_P + (x - y)|^2) \\ &\quad + |z_{P,H} - (x_H - y_H)| |\psi_r(|z_P - (x - y)|^2) - \psi_r(|z_P + (x - y)|^2)| \\ &=: T_1 + T_2. \end{aligned}$$

To deal with T_1 we write

$$T_1 \leq \frac{2|x_H - y_H|}{|z_P + (x - y)|^{n+1}}.$$

Note then that $|x_H - y_H| \leq \ell(Q_0)$, while $|x - y| \approx \text{dist}(x, H)$. Further, it is easy to check that

$$(8.27) \quad |z_P + (x - y)| \approx |z_P - (x - y)| \approx |z_P| + \text{dist}(x, H),$$

which implies that

$$T_1 \lesssim \frac{\ell(Q_0)}{(\text{dist}(x, H) + |z_P|)^{n+1}}.$$

Now we will estimate T_2 . To this end we intend to apply the mean value theorem. It is easy to check that for all $r, t > 0$,

$$|\psi_r'(t)| \lesssim \frac{1}{t^{\frac{n+3}{2}}},$$

and then, by (8.27),

$$|\psi_r(|z_P - (x - y)|^2) - \psi_r(|z_P + (x - y)|^2)| \lesssim \frac{||z_P - (x - y)|^2 - |z_P + (x - y)|^2|}{(\text{dist}(x, H) + |z_P|)^{n+3}}.$$

Now we have

$$\begin{aligned} ||z_P - (x - y)|^2 - |z_P + (x - y)|^2| &= |[z_{P,H} - (x_H - y_H)]^2 + (x_{n+1} - y_{n+1})^2] \\ &\quad - [(z_{P,H} + (x_H - y_H))^2 + (x_{n+1} - y_{n+1})^2]| \\ &= 4 |z_{P,H} \cdot (x_H - y_H)| \leq 4 |z_P| \ell(Q_0). \end{aligned}$$

Thus we infer that

$$T_2 \lesssim \frac{|z_{P,H} - (x_H - y_H)| |z_P| \ell(Q_0)}{(\text{dist}(x, H) + |z_P|)^{n+3}} \lesssim \frac{\ell(Q_0)}{(\text{dist}(x, H) + |z_P|)^{n+1}}.$$

Together with the estimate above for T_1 this yields (8.26).

From (8.24) and (8.26) we obtain

$$|\tilde{\mathcal{R}}_r \nu(x)| \lesssim |\nu|(Q_0) \sum_{P \in \mathcal{M}} \frac{\ell(Q_0)}{(\text{dist}(x, H) + |z_P|)^{n+1}}.$$

It is easy to check that

$$\sum_{P \in \mathcal{M}} \frac{\ell(Q_0)^{n+1}}{(\text{dist}(x, H) + |z_P|)^{n+1}} \lesssim \frac{\ell(Q_0)}{\text{dist}(x, H)},$$

and then (8.21) follows. \square

Lemma 8.5. *We have*

$$\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \lesssim \left(\varepsilon' + \frac{1}{A^2} \right) \eta(Q_0).$$

Proof. By Lemma 8.1 it is enough to show that

$$\int_{Q_0} |\mathcal{R}(\chi_{(AQ_0)^c} \eta)|^2 d\eta \lesssim \frac{1}{A^2} \eta(Q_0),$$

which is an immediate consequence of Lemma 8.4 (a). \square

Remark 8.6. By taking A big enough and δ, ε small enough in the assumptions of the Main Lemma 3.1, and then choosing appropriately the parameters $\varepsilon_0, \kappa_0, \theta_0$, it follows that

$$(8.28) \quad \int_{Q_0} |\mathcal{R}\eta|^2 d\eta \ll \eta(Q_0).$$

Indeed, the preceding lemma asserts that

$$\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \lesssim \left(\varepsilon' + \frac{1}{A^2} \right) \eta(Q_0),$$

with ε' given in Lemma 8.1 by

$$\varepsilon' = \tilde{\varepsilon} + A^n \kappa_0^{-2n-2} \theta_0^{\frac{1}{(n+1)^2}},$$

where $\tilde{\varepsilon}$ is defined in (7.8) by

$$\tilde{\varepsilon} = C_4 \left(\varepsilon + \frac{1}{A^2} + A^{4n+2} \delta^{\frac{1}{4n+4}} + \varepsilon_0 + \theta_0^{\frac{1}{n+1}} + \kappa_0^{\frac{1}{2}} + A^{2n+2} \tilde{\delta}^{\frac{2}{4n+5}} \right),$$

and $\tilde{\delta}$ in (7.2) by

$$\tilde{\delta} = C_3 A^{n+1} \left(\varepsilon_0 + \theta_0^{1/(n+1)} + \kappa_0^{1/2} + \delta^{1/2} \right).$$

Hence if we take first A big enough and then $\varepsilon_0, \kappa_0, \delta, \theta_0$ small enough (depending on A), so that moreover $\theta_0 \ll \kappa_0$ (to ensure that $A^n \kappa_0^{-2n-2} \theta_0^{\frac{1}{(n+1)^2}} \ll 1$), then (8.28) follows.

9. PROOF OF THE KEY LEMMA BY CONTRADICTION

9.1. A variational argument and an almost everywhere inequality.

Lemma 9.1. *Suppose that, for some $0 < \lambda \leq 1$, the inequality*

$$\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \leq \lambda \eta(Q_0)$$

holds. Then, there is a function $b \in L^\infty(\eta)$ such that

- (i) $0 \leq b \leq 2$,
- (ii) b is \mathcal{M} -periodic,
- (iii) $\int_{Q_0} b d\eta = \eta(Q_0)$,

and such that the measure $\nu = b\eta$ satisfies

$$(9.1) \quad \int_{Q_0} |\mathcal{R}\nu|^2 d\nu \leq \lambda \nu(Q_0)$$

and

$$(9.2) \quad |\mathcal{R}\nu(x)|^2 + 2\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \leq 6\lambda \quad \text{for } \nu\text{-a.e. } x \in \mathbb{R}^{n+1}.$$

Proof. In order to find such a function b , we consider the following class of admissible functions

$$(9.3) \quad \mathcal{A} = \left\{ a \in L^\infty(\eta) : a \geq 0, \text{ } a \text{ is } \mathcal{M}\text{-periodic, and } \int_{Q_0} a d\eta = \eta(Q_0) \right\}$$

and we define a functional J on \mathcal{A} by

$$(9.4) \quad J(a) = \lambda \|a\|_{L^\infty(\eta)} \eta(Q_0) + \int_{Q_0} |\mathcal{R}(a\eta)|^2 a d\eta.$$

Observe that $1 \in \mathcal{A}$ and

$$J(1) = \lambda \eta(Q_0) + \int_{Q_0} |\mathcal{R}\eta|^2 d\eta \leq 2\lambda \eta(Q_0),$$

Thus

$$\inf_{a \in \mathcal{A}} J(a) \leq 2\lambda \eta(Q_0).$$

Since $J(a) \geq \lambda \|a\|_{L^\infty(\eta)} \eta(Q_0)$, it is clear that

$$\inf_{a \in \mathcal{A}} J(a) = \inf_{a \in \mathcal{A} : \|a\|_{L^\infty(\eta)} \leq 2} J(a).$$

We claim that J attains a global minimum on \mathcal{A} , i.e., there is a function $b \in \mathcal{A}$ such that $J(b) \leq J(a)$ for all $a \in \mathcal{A}$. Indeed, by the Banach-Alaoglu theorem there exists a sequence $\{a_k\}_k \subset \mathcal{A}$, with $J(a_k) \rightarrow \inf_{a \in \mathcal{A}} J(a)$, $\|a_k\|_{L^\infty(\eta)} \leq 2$, so that a_k converges weakly $*$ in $L^\infty(\eta)$ to some function $b \in \mathcal{A}$. It is clear that b satisfies (i), (ii) and (iii). Also, since $y \mapsto \frac{x-y}{|x-y|^{n+1}}$ belongs to $L^1_{loc}(\eta)$ (recall that η has bounded density with respect to Lebesgue measure), it follows that for all $x \in Q_0$ and all $r > 0$, $\tilde{\mathcal{R}}_r(a_k\eta)(x) \rightarrow \tilde{\mathcal{R}}_r(b\eta)(x)$ as $k \rightarrow \infty$. To see that for all $x \in Q_0$, $\mathcal{R}(a_k\eta)(x) \rightarrow \mathcal{R}(b\eta)(x)$, we write for any $k, r > 0$ big enough

$$\begin{aligned} |\mathcal{R}(a_k\eta)(x) - \mathcal{R}(b\eta)(x)| &\leq |\mathcal{R}(a_k\eta)(x) - \tilde{\mathcal{R}}_r(a_k\eta)(x)| + |\tilde{\mathcal{R}}_r(a_k\eta)(x) - \tilde{\mathcal{R}}_r(b\eta)(x)| \\ &\quad + |\tilde{\mathcal{R}}_r(b\eta)(x) - \mathcal{R}(b\eta)(x)| \\ &\leq C \frac{\|a_k\|_{L^\infty(\eta)} + \|b\|_{L^\infty(\eta)}}{r} + |\tilde{\mathcal{R}}_r(a_k\eta)(x) - \tilde{\mathcal{R}}_r(b\eta)(x)|, \end{aligned}$$

applying (8.19) for the last inequality. Taking limits in k in both sides we obtain

$$\limsup_{k \rightarrow \infty} |\mathcal{R}(a_k\eta)(x) - \mathcal{R}(b\eta)(x)| \leq \frac{C}{r} + \limsup_{k \rightarrow \infty} |\tilde{\mathcal{R}}_r(a_k\eta)(x) - \tilde{\mathcal{R}}_r(b\eta)(x)| = \frac{C}{r}.$$

Since this holds for all $r > 0$ big enough, we infer that $\mathcal{R}(a_k\eta)(x) \rightarrow \mathcal{R}(b\eta)(x)$ as $k \rightarrow \infty$, as wished. Taking into account that

$$|\mathcal{R}(a_k\eta)(x)| \leq \frac{C}{r} + |\tilde{\mathcal{R}}_r(a_k\eta)(x)| \leq \frac{C}{r} + 2 \int_{|x-y| \leq r} \frac{1}{|x-y|^n} d\eta(y) \leq C(r)$$

for all $r > 0$ big enough, by the dominated convergence theorem we infer that

$$\int_{Q_0} |\mathcal{R}(a_k\eta)|^2 d\eta \rightarrow \int_{Q_0} |\mathcal{R}(b\eta)|^2 d\eta \quad \text{as } k \rightarrow \infty.$$

Using also that $\|b\|_{L^\infty(\eta)} \leq \limsup_k \|a_k\|_{L^\infty(\eta)}$, it follows that $J(b) \leq \limsup_k J(a_k)$, which proves the claim that $J(\cdot)$ attains a minimum at b .

The estimate (9.1) for $\nu = b\eta$ follows from the fact that $J(b) \leq J(1)$, because the property (iii) implies that $\|b\|_{L^\infty(\eta)} \geq 1$.

To prove (9.2) we perform a blow-up argument taking advantage of the fact that b is a minimizer for J . Let B be any ball contained in Q_0 and centered on $\text{supp } \nu \cap Q_0$. Let

$$(9.5) \quad P_{\mathcal{M}}(B) = \bigcup_{R \in \mathcal{M}} (B + z_R)$$

be the “periodic extension” of B with respect to \mathcal{M} . Now, for every $0 \leq t < 1$, define

$$(9.6) \quad b_t = (1 - t\chi_{P_{\mathcal{M}}(B)})b + t \frac{\nu(B)}{\nu(Q_0)} b.$$

It is clear that $b_t \in \mathcal{A}$ for all $0 \leq t < 1$ and $b_0 = b$. Therefore,

$$\begin{aligned} (9.7) \quad J(b) &\leq J(b_t) = \lambda \|b_t\|_\infty \eta(Q_0) + \int_{Q_0} |\mathcal{R}(b_t\eta)|^2 b_t d\eta \\ &\leq \lambda \left(1 + t \frac{\nu(B)}{\nu(Q_0)}\right) \|b\|_\infty \eta(Q_0) + \int_{Q_0} |\mathcal{R}(b_t\eta)|^2 b_t d\eta := h(t). \end{aligned}$$

Since $h(0) = J(b)$, we have that $h(0) \leq h(t)$ for $0 \leq t < 1$ and, thus $h'(0+) \geq 0$ (assuming that $h'(0+)$ exists). Notice that

$$\left. \frac{db_t}{dt} \right|_{t=0} = -\chi_{P_{\mathcal{M}}(B)} b + \frac{\nu(B)}{\nu(Q_0)} b,$$

Therefore,

$$\begin{aligned} 0 \leq h'(0+) &= \lambda \frac{\nu(B)}{\nu(Q_0)} \|b\|_{\infty} \eta(Q_0) + \left. \frac{d}{dt} \right|_{t=0} \int_{Q_0} |\mathcal{R}(b_t \eta)|^2 b_t d\eta \\ &= \lambda \frac{\nu(B)}{\nu(Q_0)} \|b\|_{\infty} \eta(Q_0) + 2 \int_{Q_0} \mathcal{R} \left(\left. \frac{db_t}{dt} \right|_{t=0} \eta \right) \cdot \mathcal{R} \nu b d\eta + \int_{Q_0} |\mathcal{R} \nu|^2 \left. \frac{db_t}{dt} \right|_{t=0} d\eta \\ &= \lambda \frac{\nu(B)}{\nu(Q_0)} \|b\|_{\infty} \eta(Q_0) + 2 \int_{Q_0} \mathcal{R} \left(\left(-\chi_{P_{\mathcal{M}}(B)} b + \frac{\nu(B)}{\nu(Q_0)} b \right) \eta \right) \cdot \mathcal{R} \nu b d\eta \\ &\quad + \int_{Q_0} |\mathcal{R} \nu|^2 \left(-\chi_{P_{\mathcal{M}}(B)} b + \frac{\nu(B)}{\nu(Q_0)} b \right) d\eta \\ &= \lambda \frac{\nu(B)}{\nu(Q_0)} \|b\|_{\infty} \eta(Q_0) - 2 \int_{Q_0} \mathcal{R}(\chi_{P_{\mathcal{M}}(B)} \nu) \cdot \mathcal{R} \nu d\nu + 2 \frac{\nu(B)}{\nu(Q_0)} \int_{Q_0} |\mathcal{R} \nu|^2 d\nu \\ &\quad - \int_B |\mathcal{R} \nu|^2 d\nu + \frac{\nu(B)}{\nu(Q_0)} \int_{Q_0} |\mathcal{R} \nu|^2 d\nu, \end{aligned}$$

where we used that $P_{\mathcal{M}}(B) \cap Q_0 = B$ in the last identity. The fact that the derivatives above commute with the integral sign and with the operator \mathcal{R} is guaranteed by the fact that b_t is an affine function of t and then one can expand the integrand $|\mathcal{R}(b_t \eta)|^2 b_t$ and obtain a polynomial expression on t . Using also that $\lambda \leq 1$ and that $J(b) \leq 2\lambda \nu(Q_0)$, we get

$$\begin{aligned} (9.8) \quad \int_B |\mathcal{R} \nu|^2 d\nu + 2 \int_{Q_0} \mathcal{R}(\chi_{P_{\mathcal{M}}(B)} \nu) \cdot \mathcal{R} \nu d\nu &\leq \frac{\nu(B)}{\nu(Q_0)} \left[\lambda \|b\|_{\infty} \eta(Q_0) + 3 \int_{Q_0} |\mathcal{R} \nu|^2 d\nu \right] \\ &\leq 3 J(b) \nu(B) \leq 6\lambda \nu(B). \end{aligned}$$

We claim now that

$$(9.9) \quad \int_{Q_0} \mathcal{R}(\chi_{P_{\mathcal{M}}(B)} \nu) \cdot \mathcal{R} \nu d\nu = \int_B \mathcal{R}^*((\mathcal{R} \nu) \nu) d\nu.$$

Assuming this for the moment, from (9.8) and (9.9), dividing by $\nu(B)$, we obtain

$$\frac{1}{\nu(B)} \int_B |\mathcal{R} \nu|^2 d\nu + \frac{2}{\nu(B)} \int_B \mathcal{R}^*((\mathcal{R} \nu) \nu) d\nu \leq 6\lambda,$$

and so, letting $\nu(B) \rightarrow 0$ and applying Lebesgue's differentiation theorem, we obtain

$$|\mathcal{R} \nu(x)|^2 + 2\mathcal{R}^*((\mathcal{R} \nu) \nu)(x) \leq 6\lambda \quad \text{for } \nu\text{-a.e. } x \in \mathbb{R}^{n+1},$$

as desired.

It remains to prove the claim (9.9). By the uniform convergence of $\tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)} \nu)$ and $\tilde{\mathcal{R}}_r \nu$ to $\mathcal{R}(\chi_{P_{\mathcal{M}}(B)} \nu)$ and $\mathcal{R} \nu$, respectively, we have

$$(9.10) \quad \int_{Q_0} \mathcal{R}(\chi_{P_{\mathcal{M}}(B)} \nu) \cdot \mathcal{R} \nu d\nu = \lim_{r \rightarrow \infty} \int_{Q_0} \tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)} \nu) \cdot \tilde{\mathcal{R}}_r \nu d\nu.$$

Since $\tilde{K}_r(x - \cdot)$ has compact support, for all $x \in Q_0$,

$$\tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)}\nu)(x) = \int_{P_{\mathcal{M}}(B)} \tilde{K}_r(x - y) d\nu(y) = \sum_{P \in \mathcal{M}} \int \tilde{K}_r(x - y) d((T_P)_\#(\chi_B\nu)(y)).$$

For the last identity we have used the fact that the sum above runs only over a finite number of $P \in \mathcal{M}$ because there is only a finite number of non-zero terms (in fact, we may assume these $P \in \mathcal{M}$ to be independent of $x \in Q_0$). Thus we have

$$\tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)}\nu)(x) = \sum_{P \in \mathcal{M}} \int_B \tilde{K}_r(x - y - z_P) d\nu(y) = \sum_{P \in \mathcal{M}} \tilde{\mathcal{R}}_r(\chi_B\nu)(x - z_P),$$

and so

$$\begin{aligned} \int_{Q_0} \tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)}\nu)(x) \cdot \tilde{\mathcal{R}}_r\nu d\nu(x) &= \sum_{P \in \mathcal{M}} \int_{Q_0} \tilde{\mathcal{R}}_r(\chi_B\nu)(x - z_P) \cdot \tilde{\mathcal{R}}_r\nu(x) d\nu(x) \\ &= \sum_{P \in \mathcal{M}} \int_{Q_0 - z_P} \tilde{\mathcal{R}}_r(\chi_B\nu)(x) \cdot \tilde{\mathcal{R}}_r\nu(x + z_P) d((T_P)^{-1}_\#\nu)(x) \end{aligned}$$

Since $\tilde{\mathcal{R}}_r\nu$ is \mathcal{M} -periodic, $\tilde{\mathcal{R}}_r\nu(x + z_P) = \tilde{\mathcal{R}}_r\nu(x)$ and $(T_P)^{-1}_\#\nu = \nu$, and then by Fubini,

$$\begin{aligned} (9.11) \quad \int_{Q_0} \tilde{\mathcal{R}}_r(\chi_{P_{\mathcal{M}}(B)}\nu)(x) \cdot \tilde{\mathcal{R}}_r\nu(x) d\nu(x) &= \sum_{P \in \mathcal{M}} \int_{Q_0 - z_P} \tilde{\mathcal{R}}_r(\chi_B\nu)(x) \cdot \tilde{\mathcal{R}}_r\nu(x) d\nu(x) \\ &= \int \tilde{\mathcal{R}}_r(\chi_B\nu)(x) \cdot \tilde{\mathcal{R}}_r\nu(x) d\nu(x) \\ &= \int_B \tilde{\mathcal{R}}_r^*((\tilde{\mathcal{R}}_r\nu)\nu)(y) d\nu(y). \end{aligned}$$

Since $\tilde{\mathcal{R}}_r\nu$ converges uniformly to $\mathcal{R}\nu$ as $r \rightarrow \infty$ and $\tilde{\mathcal{R}}_r^*$ tends to \mathcal{R}^* in operator norm in $L^\infty_{\mathcal{M}}(\eta) \rightarrow L^\infty_{\mathcal{M}}(\eta)$, we deduce that

$$\lim_{r \rightarrow \infty} \int_B \tilde{\mathcal{R}}_r^*((\tilde{\mathcal{R}}_r\nu)\nu)(y) d\nu(y) = \int_B \mathcal{R}^*((\mathcal{R}\nu)\nu) d\nu.$$

Together with (9.10) and (9.11) this yields (9.9). \square

9.2. A maximum principle.

Lemma 9.2. Assume that $\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \leq \lambda\eta(Q_0)$ for some $0 < \lambda \leq 1$, and let b and ν be as in Lemma 9.1. Let $K_S > 10$ be some (big) constant and let S be the horizontal strip

$$S = \{x \in \mathbb{R}^{n+1} : |x_{n+1}| \leq K_S \ell(Q_0)\}.$$

Also, set

$$f(x) = c_S x_{n+1} e_{n+1} = c_S(0, \dots, 0, x_{n+1}), \quad \text{with } c_S = \int \frac{1}{(|y_H|^2 + (K_S \ell(Q_0))^2)^{\frac{n+1}{2}}} d\nu(y).$$

Then, we have

$$(9.12) \quad |\mathcal{R}\nu(x) - f(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \lesssim \lambda^{1/2} + \frac{1}{K_S^2} \quad \text{for all } x \in S.$$

Further,

$$(9.13) \quad c_S \lesssim \frac{1}{K_S \ell(Q_0)}.$$

Proof. The inequality (9.13) is very easy. Indeed, we just have to use that $\nu(B(x, r)) \lesssim r^n$ for all $x \in \mathbb{R}^{n+1}$ and $r > 0$, and use standard estimates which we leave for the reader.

To prove (9.12), we denote

$$F(x) = |\mathcal{R}\nu(x) - f(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x).$$

It is clear that F is subharmonic in $\mathbb{R}^{n+1} \setminus \text{supp}(\nu)$ and continuous in the whole space \mathbb{R}^{n+1} , by Lemma 8.2. So if we show that the estimate in (9.12) holds for all $x \in \text{supp } \nu \cup \partial S$, then this will be also satisfied in the whole S . Indeed, since F is \mathcal{M} -periodic and continuous in S , the maximum of F in S is attained, and since F is subharmonic in $\mathring{S} \setminus \text{supp } \nu$, this must be attained in $\text{supp } \nu \cup \partial S$.

First we check that the inequality in (9.12) holds for all $x \in \text{supp } \nu$. To this end, recall that by Lemma 9.1

$$|\mathcal{R}\nu(x)|^2 + 2\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \leq 6\lambda \quad \nu\text{-almost everywhere in } \text{supp}(\nu),$$

and this inequality extends to the whole $\text{supp}(\nu)$ by continuity. Therefore we have, for all $x \in \text{supp}(\nu)$,

$$\begin{aligned} F(x) &= |\mathcal{R}\nu(x) - f(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \leq 2|\mathcal{R}\nu(x)|^2 + 2|f(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \\ &\leq 12\lambda + 2|f(x)|^2 \leq 12\lambda + (c_S \ell(Q_0))^2 \lesssim \lambda + \frac{1}{K_S^2}, \end{aligned}$$

where we took into account that $|x_{n+1}| \leq \frac{1}{2}\ell(Q_0)$ for $x \in \text{supp } \nu$ and we used (9.13).

Our next objective consists in getting an upper bound for F in ∂S . By applying Lemma 8.4 to the function $\mathcal{R}\nu$ (which is \mathcal{M} -periodic), with \mathcal{R}^* instead of \mathcal{R} (since \mathcal{R} is antisymmetric we are allowed to do this) we deduce that, for all $x \in \partial S$,

$$|\mathcal{R}^*((\mathcal{R}\nu)\nu)(x)| \lesssim \frac{1}{\ell(Q_0)^n} \int_{Q_0} |\mathcal{R}\nu| \, d\nu \lesssim \frac{1}{\ell(Q_0)^n} \left(\int_{Q_0} |\mathcal{R}\nu|^2 \, d\nu \right)^{1/2} \nu(Q_0)^{1/2} \lesssim \lambda^{1/2}.$$

It suffices to show now that $|\mathcal{R}\nu(x) - f(x)| \lesssim \frac{1}{K_S}$ for all $x \in \partial S$. We write $\mathcal{R}\nu(x) = (\mathcal{R}^H \nu(x), \mathcal{R}_{n+1} \nu(x))$. From (8.21) we infer that

$$|\mathcal{R}^H \nu(x)| \lesssim \frac{1}{K_S \ell(Q_0)^n} \nu(Q_0) \lesssim \frac{1}{K_S}.$$

Hence it just remains to prove that

$$(9.14) \quad |\mathcal{R}_{n+1} \nu(x) e_{n+1} - f(x)| \lesssim \frac{1}{K_S} \quad \text{for all } x \in \partial S.$$

To prove this estimate we can assume without loss of generality that $x_{n+1} = K_S \ell(Q_0)$ and that $x_H \in Q_0 \cap H$, by the \mathcal{M} -periodicity of $\mathcal{R}_{n+1} \nu$. Since $f(x) = c_S K_S \ell(Q_0) e_{n+1}$ for this point x , (9.14) is equivalent to

$$(9.15) \quad |\mathcal{R}_{n+1} \nu(x) - c_S K_S \ell(Q_0)| \lesssim \frac{1}{K_S}.$$

Note first that

$$\mathcal{R}_{n+1}\nu(x) = \lim_{r \rightarrow 0} \int \phi_r(x-y) \frac{x_{n+1} - y_{n+1}}{|x-y|^{n+1}} d\nu(y) = \int \frac{x_{n+1} - y_{n+1}}{|x-y|^{n+1}} d\nu(y),$$

by an easy application of the dominated convergence theorem (using that $|x_{n+1} - y_{n+1}| \leq \text{dist}(x, H) + \ell(Q_0)$). Consider the point $x_0 = (0, K_S \ell(Q_0))$. Since for all $y \in \text{supp } \nu$,

$$|x - x_0| + |y - y_H| \leq \ell(Q_0) \leq \frac{1}{2}|x - y|,$$

and since the $(n+1)$ -th component of $K(\cdot)$, which we denote by $K_{n+1}(\cdot)$, is a standard Calderón-Zygmund kernel,

$$|K_{n+1}(x-y) - K_{n+1}(x_0 - y_H)| \lesssim \frac{|x - x_0| + |y - y_H|}{|x - y|^{n+1}} \lesssim \frac{\ell(Q_0)}{|x - y|^{n+1}}.$$

Therefore, integrating with respect to ν , we derive

$$|\mathcal{R}_{n+1}\nu(x) - c_S K_S \ell(Q_0)| = \left| \int (K_{n+1}(x-y) - K_{n+1}(x_0 - y_H)) d\nu(y) \right| \lesssim \int \frac{\ell(Q_0)}{|x - y|^{n+1}} d\nu(y).$$

Since $\text{dist}(x, \text{supp } \nu) \gtrsim K_S \ell(Q_0)$ and ν is a measure with growth of order n , by standard estimates it follows that

$$\int \frac{\ell(Q_0)}{|x - y|^{n+1}} d\nu(y) \lesssim \frac{1}{K_S},$$

which proves (9.15) and finishes the proof of the lemma. \square

The next result is an immediate consequence of Lemma 9.2.

Lemma 9.3. *Assume that, for some $0 < \lambda \leq 1$, the inequality*

$$\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \leq \lambda \eta(Q_0)$$

is satisfied, and let b and ν be as in Lemma 9.1. Then, we have

$$(9.16) \quad |\mathcal{R}\nu(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \lesssim \lambda^{1/2} \quad \text{for all } x \in \mathbb{R}^{n+1}.$$

Proof. This follows by letting $K_S \rightarrow \infty$ in the inequality (9.12), taking into account that $c_S \rightarrow 0$, by (9.13). \square

9.3. The contradiction.

Lemma 9.4. *Suppose that, for some $0 < \lambda \leq 1$, the inequality*

$$(9.17) \quad \int_{Q_0} |\mathcal{R}\eta|^2 d\eta \leq \lambda \eta(Q_0)$$

is satisfied, and let b and ν be as in Lemma 9.1. Then, there exists some constant $c_3 > 0$ depending only¹ on n, C_0, C_1 such that

$$\lambda \geq c_3.$$

¹In fact, keeping track of the dependencies, one can check that c_3 depends only on n and C_0 , and not on C_1 . However, this is not necessary for the proof of the Key Lemma.

Proof. By Lemma 9.3, we have

$$(9.18) \quad |\mathcal{R}\nu(x)|^2 + 4\mathcal{R}^*((\mathcal{R}\nu)\nu)(x) \lesssim \lambda^{1/2} \quad \text{everywhere in } \mathbb{R}^{n+1}.$$

Now pick a smooth function φ with $\chi_{Q_0} \leq \varphi \leq \chi_{2Q_0}$ and $\|\nabla\varphi\|_\infty \lesssim \frac{1}{\ell(Q_0)}$. Set $\psi = C_5\nabla\varphi$, so that $\mathcal{R}^*(\psi\mathcal{H}^{n+1}) = \varphi$. Then, we have

$$\begin{aligned} \eta(Q_0) = \nu(Q_0) &\leq \int \varphi \, d\nu = \int \mathcal{R}^*(\psi\mathcal{H}^{n+1}) \, d\nu \\ &= \int \mathcal{R}\nu \, \psi \, d\mathcal{H}^{n+1} \leq \left(\int |\mathcal{R}\nu|^2 |\psi| \, d\mathcal{H}^{n+1} \right)^{1/2} \left(\int |\psi| \, d\mathcal{H}^{n+1} \right)^{1/2}. \end{aligned}$$

First of all, observe that

$$\|\psi\|_\infty \lesssim \frac{1}{\ell(Q_0)} \quad \text{and} \quad \int |\psi| \, d\mathcal{H}^{n+1} \lesssim \ell(Q_0)^n$$

and so

$$(9.19) \quad \eta(Q_0) \lesssim \left(\int |\mathcal{R}\nu|^2 |\psi| \, d\mathcal{H}^{n+1} \right)^{1/2} \ell(Q_0)^{n/2}.$$

Furthermore, by (9.18) we have

$$\begin{aligned} (9.20) \quad \int |\mathcal{R}\nu|^2 |\psi| \, d\mathcal{H}^{n+1} &\leq C \lambda^{1/2} \int |\psi| \, d\mathcal{H}^{n+1} + 4 \left| \int \mathcal{R}^*((\mathcal{R}\nu)\nu) |\psi| \, d\mathcal{H}^{n+1} \right| \\ &\lesssim \lambda^{1/2} \ell(Q_0)^n + \left| \int \mathcal{R}^*(\chi_{(5Q_0)^c}(\mathcal{R}\nu)\nu) |\psi| \, d\mathcal{H}^{n+1} \right| + \left| \int \mathcal{R}^*(\chi_{5Q_0}(\mathcal{R}\nu)\nu) |\psi| \, d\mathcal{H}^{n+1} \right|. \end{aligned}$$

To estimate the first integral on the right hand side we apply Lemma 8.4 (a) with $\tilde{A} = 5$ and $f = \mathcal{R}\nu b$ (where b is such that $b\eta = \nu$), and then we deduce that for all $x \in 2Q_0$,

$$\begin{aligned} |\mathcal{R}^*(\chi_{(5Q_0)^c}(\mathcal{R}\nu)\nu)(x)| &\lesssim \frac{1}{\ell(Q_0)^n} \int_{Q_0} |\mathcal{R}\nu b| \, d\eta \\ &= \frac{1}{\ell(Q_0)^n} \int_{Q_0} |\mathcal{R}\nu| \, d\nu \lesssim \left(\int_{Q_0} |\mathcal{R}\nu|^2 \, d\nu \right)^{1/2} \lesssim \lambda^{1/2}. \end{aligned}$$

Thus, recalling that ψ is supported in $2Q_0$,

$$\left| \int \mathcal{R}^*(\chi_{(5Q_0)^c}(\mathcal{R}\nu)\nu) |\psi| \, d\mathcal{H}^{n+1} \right| \lesssim \lambda^{1/2} \|\psi\|_1 \lesssim \lambda^{1/2} \nu(Q_0).$$

Concerning the last integral on the right hand side of (9.20), we have

$$\begin{aligned} \left| \int \mathcal{R}^*(\chi_{5Q_0}(\mathcal{R}\nu)\nu) |\psi| \, d\mathcal{H}^{n+1} \right| &= \left| \int_{5Q_0} \mathcal{R}\nu \cdot \mathcal{R}(|\psi| \, d\mathcal{H}^{n+1}) \, d\nu \right| \\ &\leq \left(\int_{5Q_0} |\mathcal{R}\nu|^2 \, d\nu \right)^{1/2} \left(\int_{5Q_0} |\mathcal{R}(|\psi| \, d\mathcal{H}^{n+1})|^2 \, d\nu \right)^{1/2}. \end{aligned}$$

The first integral on the right hand side does not exceed $c\lambda\nu(Q_0)$ (by (9.1) and the periodicity of $\mathcal{R}\nu$). For the second one, using that $|\psi| \lesssim \frac{1}{\ell(Q_0)}\chi_{2Q_0}$, it follows easily that $\|\mathcal{R}(|\psi| \mathcal{H}^{n+1})\|_\infty \lesssim 1$. So we get

$$\left| \int \mathcal{R}^*(\chi_{5Q_0}(\mathcal{R}\nu)\nu)|\psi| d\mathcal{H}^{n+1} \right| \lesssim \lambda^{1/2}\nu(Q_0).$$

So from (9.20) and the last estimates we deduce that

$$\int |\mathcal{R}\nu|^2 |\psi| d\mathcal{H}^{n+1} \lesssim \lambda^{1/2}\nu(Q_0).$$

Thus, by (9.19),

$$\nu(Q_0) \lesssim \lambda^{1/4}\nu(Q_0).$$

That is, $\lambda \gtrsim 1$. □

Now to conclude the proof of the Key Lemma 6.1 we only have to recall that, by Remark 8.6, $\int_{Q_0} |\mathcal{R}\eta|^2 d\eta \ll \eta(Q_0)$ if A is big enough and $\delta, \varepsilon, \kappa_0, \theta_0$ small enough and chosen suitably, under the assumption that ε_0 is small enough too. This contradicts the conclusion of Lemma 9.4. Hence, (6.1) cannot hold and thus we are done.

10. CONSTRUCTION OF THE AD-REGULAR MEASURE ζ AND THE UNIFORMLY RECTIFIABLE SET Γ IN THE MAIN LEMMA

Denote

$$(10.1) \quad F = Q_0 \cap \text{supp } \mu \setminus \bigcup_{Q \in \text{LD}} Q.$$

It is easy to check that $0 < \Theta_*^n(x, \mu) \leq \Theta^{n,*}(x, \mu) < \infty$ for μ -a.e. $x \in F$. Since \mathcal{R}_μ is bounded on $L^2(\mu|_F)$ it follows that $\mu|_F$ is n -rectifiable, by the Nazarov-Tolsa-Volberg theorem [NToV2]. However, to get a big piece of a set contained in a uniformly n -rectifiable set Γ like the one required in the Main Lemma and in Theorem 1.1 we have to argue more carefully. To this end, first we will construct an auxiliary AD-regular measure ζ such that $\zeta(F) \gtrsim \mu(F)$, and then we will apply the Nazarov-Tolsa-Volberg theorem [NToV1] for AD-regular measures.

Next we are going to construct the aforementioned auxiliary measure ζ . The arguments for this construction can be considered as quantitative version of the ones from [NToV2], which rely on a covering theorem of Pajot (see [Pa]).

Recall the notation $\sigma = \mu|_{Q_0}$. Consider the maximal dyadic operator

$$\mathcal{M}_{\mathcal{D}_\sigma} f(x) = \sup_{Q \in \mathcal{D}_\sigma : x \in Q} \frac{1}{\sigma(Q)} \int_Q |f| d\sigma,$$

where \mathcal{D}_σ is the David-Mattila lattice associated σ . Let F be as in (10.1) and set

$$\tilde{F} = \left\{ x \in F : \mathcal{M}_{\mathcal{D}_\sigma}(\chi_{F^c})(x) \leq 1 - \frac{\varepsilon_0}{2} \right\}.$$

We wish to show that

$$(10.2) \quad \sigma(\tilde{F}) \geq \frac{1}{2} \sigma(F).$$

To this end, note that

$$F \setminus \tilde{F} = \left\{ x \in F : \mathcal{M}_{\mathcal{D}_\sigma}(\chi_{F^c})(x) > 1 - \frac{\varepsilon_0}{2} \right\}$$

and consider a collection of maximal (and thus disjoint) cells $\{Q_i\}_{i \in J} \subset \mathcal{D}_\sigma$ such that $\sigma(Q_i \setminus F) > (1 - \frac{\varepsilon_0}{2})\sigma(Q_i)$. Observe that

$$F \setminus \tilde{F} = \bigcup_{i \in J} Q_i \cap F.$$

Clearly, the cells Q_i satisfy $\sigma(Q_i \cap F) \leq \frac{\varepsilon_0}{2}\sigma(Q_i)$ and so we have

$$\sigma(F \setminus \tilde{F}) \leq \sum_{i \in J} \sigma(Q_i \cap F) \leq \sum_{i \in J} \frac{\varepsilon_0}{2} \sigma(Q_i) \leq \frac{\varepsilon_0}{2} \sigma(Q_0) \leq \frac{1}{2} \sigma(F),$$

which proves (10.2).

For each $i \in J$ we consider the family \mathcal{A}_i of maximal doubling cells from \mathcal{D}_σ^{db} which cover Q_i , and we define

$$\mathcal{A} = \bigcup_{i \in J} \mathcal{A}_i.$$

Finally, we denote by \mathcal{A}_0 the subfamily of the cells $P \in \mathcal{A}$ such that $\sigma(P \cap F) > 0$. Now, for each $Q \in \mathcal{A}_0$ we consider an n -dimensional sphere $S(Q)$ concentric with $B(Q)$ and with radius $\frac{1}{4}r(B(Q))$. We define

$$\zeta = \sigma|_{\tilde{F}} + \sum_{Q \in \mathcal{A}_0} \mathcal{H}^n|_{S(Q)}.$$

Remark 10.1. If $P \in \mathcal{A}_0$ and $P \subset Q_i$ for some $i \in J$, then

$$\ell(P) \approx_{\theta_0, C_0} \ell(Q_i).$$

Indeed, since P is a maximal doubling cell contained in Q_i , by Lemma 5.5 and the fact that $3.5B_P \subset 100B(P)$,

$$\Theta_\sigma(3.5B_P) \lesssim \Theta_\sigma(100B(P)) \lesssim A_0^{-9n(J(P)-J(Q_i))} \Theta_\sigma(100B(Q_i)) \lesssim_{C_0} A_0^{-9n(J(P)-J(Q_i))}.$$

Since $\sigma(P \cap F) > 0$, it turns out that P is not contained in any cell from LD, and so $\Theta_\sigma(3.5B_P) > \theta_0$. So we have

$$\theta_0 \lesssim_{C_0} A_0^{-9n(J(P)-J(Q_i))},$$

which implies that $|J(P) - J(Q_i)| \lesssim_{\theta_0, C_0} 1$.

A very similar argument shows that if $P \in \mathcal{D}_\sigma$ satisfies $P \cap F \neq \emptyset$ (and so it is not contained in any cell from LD), then there exists some $Q \in \mathcal{D}_\sigma^{db}$ which contains Q and such that

$$\ell(P) \approx_{\theta_0, C_0} \ell(Q).$$

The details are left for the reader.

From the two statements above, it follows that for any cell $P \in \mathcal{D}_\sigma$ which is not strictly contained in any cell from \mathcal{A}_0 there exists some cell $\hat{P} \in \mathcal{D}_\sigma^{db}$ which is not contained in any cell Q_i , $i \in J$, so that $P \subset \hat{P}$ and $\ell(P) \approx_{\theta_0, C_0} \ell(Q)$.

Lemma 10.2. *The measure ζ is AD regular, with the AD-regularity constant depending on C_0 , θ_0 , and ε_0 .*

Proof. First we will show the upper AD-regularity of ζ . That is, we will prove that $\zeta(B(x, r)) \leq C(C_0, \theta_0) r^n$ for all x, r . By the upper AD-regularity of σ , it is enough to show that the measure

$$\nu = \sum_{Q \in \mathcal{A}_0} \mathcal{H}^n|_{S(Q)}$$

is also upper AD-regular. So we have to prove that

$$(10.3) \quad \nu(B(x, r)) \leq C(C_0, \theta_0) r^n \quad \text{for all } x \in \bigcup_{Q \in \mathcal{A}_0} S(Q) \text{ and all } r > 0.$$

Take $x \in S(Q)$, for some $Q \in \mathcal{A}_0$. Clearly the estimate above holds if the only sphere $S(P)$, $P \in \mathcal{A}_0$, that intersects $B(x, r)$ is just $S(Q)$ itself. So assume that $B(x, r)$ intersects a sphere $S(P)$, $P \in \mathcal{A}_0$, with $P \neq Q$. Recall that $\frac{1}{2}B(Q) \cap \frac{1}{2}B(P) = \emptyset$, by Remark 5.2, and thus for some constant C_6 , $P \subset B(x, C_6 r)$. Hence,

$$\nu(B(x, r)) \leq \sum_{P \in \mathcal{A}_0: P \subset B(x, C_6 r)} \nu(\tfrac{1}{4}S(P)) \lesssim \sum_{P \in \mathcal{A}_0: P \subset B(x, C_6 r)} \ell(P)^n.$$

Note now that by the definition of \mathcal{A}_0 , $\sigma(F \cap P) > 0$, which implies that $P \notin \text{LD}$ and that P is not contained in any cell from LD, and thus taking also into account that $P \in \mathcal{D}^{db}$,

$$(10.4) \quad \sigma(P) \gtrsim \sigma(3.5B_P) \gtrsim \theta_0 \ell(P)^n.$$

Together with the upper AD-regularity of σ this yields

$$\nu(B(x, r)) \lesssim \frac{1}{\theta_0} \sum_{P \in \mathcal{A}_0: P \subset B(x, C_6 r)} \sigma(P) \lesssim \frac{1}{\theta_0} \sigma(B(x, C_6 r)) \lesssim_{C_0, \theta_0} r^n,$$

which concludes the proof of (10.3).

It remains now to show the lower AD-regularity of ζ . First we will prove that

$$(10.5) \quad \zeta(2B_Q) \gtrsim_{\theta_0, \varepsilon_0, C_0} \ell(Q)^n \quad \text{if } Q \in \mathcal{D}_\sigma^{db} \text{ is not contained in any cell } Q_i, i \in J.$$

Indeed, note that by the definition of the cells $Q_i, i \in J$,

$$\sigma(Q \setminus F) \leq \left(1 - \frac{\varepsilon_0}{2}\right) \sigma(Q),$$

or equivalently,

$$\sigma(Q \cap F) \geq \frac{\varepsilon_0}{2} \sigma(Q).$$

Since Q is not contained in any cell from LD (by the definitions of F and \mathcal{A}_0) and is doubling,

$$(10.6) \quad \sigma(Q \cap F) \gtrsim_{\varepsilon_0} \sigma(3.5B_Q) \gtrsim_{\theta_0, \varepsilon_0} \ell(Q)^n.$$

On the other hand, by the construction of ζ ,

$$\sigma(Q \cap F) = \sigma(Q \cap \tilde{F}) + \sum_{P \in \mathcal{A}_0: P \subset Q} \sigma(P \cap F) \lesssim_{C_0} \zeta(Q \cap \tilde{F}) + \sum_{P \in \mathcal{A}_0: P \subset Q} \mathcal{H}^n(S(P)).$$

We may assume that all the cells $P \subset Q$ satisfy $S(P) \subset 2B_Q$, just by choosing the constant A_0 in the construction of the lattice \mathcal{D}_σ big enough. Then we get

$$\sigma(Q \cap F) \lesssim_{C_0} \zeta(Q \cap \tilde{F}) + \sum_{P \in \mathcal{A}_0: S(P) \subset 2B_Q} \zeta(S(P)) \lesssim_{C_0} \zeta(2B_Q).$$

Together with (10.6), this gives (10.5).

To prove the lower AD regularity of ζ , note that by Remark 10.1 there is some constant $C'(C_0, \theta_0)$ such that if $x \in S(Q)$, $Q \in \mathcal{A}_0$, and $C'(C_0, \theta_0) \ell(Q) < r \leq \text{diam}(Q_0)$, then there exists $P \in \mathcal{D}_\sigma^{db}$

not contained in any cell Q_i , $i \in J$, such that $2B_P \subset B(x, r)$, with $\ell(P) \approx_{\theta_0, C_0} r$. The same holds for $0 < r \leq \text{diam}(Q_0)$ if $x \in \tilde{F}$. From (10.5) we deduce that

$$\zeta(B(x, r)) \geq \zeta(2B_P) \gtrsim_{\theta_0, \varepsilon_0, C_0} \ell(P)^n \approx_{\theta_0, \varepsilon_0, C_0} r^n.$$

In the case that $r \leq C'(C_0, \theta_0) \ell(Q)$ for $x \in S(Q)$, $Q \in \mathcal{A}_0$, the lower AD-regularity of $\mathcal{H}^n|_{S(Q)}$ gives the required lower estimate for $\zeta(B(x, r))$. \square

Lemma 10.3. *The Riesz transform \mathcal{R}_ζ is bounded in $L^2(\zeta)$, with a bound on the norm depending on C_0 , C_1 , θ_0 , and ε_0 .*

To prove this result we will follow very closely the arguments in the last part of the proof of the Main Lemma 2.1 of [NToV2]. For completeness, we will show all the details.

For technical reasons, it will be convenient to work with an ε -regularized version $\widehat{\mathcal{R}}_{\nu, \varepsilon}$ of the Riesz transform \mathcal{R}_ν . For a measure ν with growth of order n , we set

$$\widehat{\mathcal{R}}_{\nu, \varepsilon} f(x) = \int \frac{x - y}{\max(|x - y|, \varepsilon)^{n+1}} f(y) d\nu(y).$$

It is easy to check that

$$|\widehat{\mathcal{R}}_{\nu, \varepsilon} f(x) - \mathcal{R}_{\nu, \varepsilon} f(x)| \leq c M_n f(x) \quad \text{for all } x \in \mathbb{R}^{n+1},$$

where c is independent of ε and M_n is the maximal operator defined by

$$M_n f(x) = \sup_{r>0} \frac{1}{r^n} \int_{B(x, r)} |f| d\nu.$$

Since M_n is bounded in $L^2(\nu)$ (because ν has growth of order n), it turns out that \mathcal{R}_ν is bounded in $L^2(\nu)$ if and only if the operators $\widehat{\mathcal{R}}_{\nu, \varepsilon}$ are bounded in $L^2(\nu)$ uniformly on $\varepsilon > 0$. The advantage of $\widehat{\mathcal{R}}_{\nu, \varepsilon}$ over $\mathcal{R}_{\nu, \varepsilon}$ is that the kernel

$$\widehat{K}_\varepsilon(x) = \frac{x}{\max(|x|, \varepsilon)^{n+1}}$$

is continuous and satisfies the smoothness condition

$$|\nabla \widehat{K}_\varepsilon(x)| \leq \frac{c}{|x|^{n+1}}, \quad |x| \neq \varepsilon$$

(with c independent of ε), which implies that $\widehat{K}_\varepsilon(x - y)$ is a standard Calderón-Zygmund kernel (with constants independent of ε), unlike the kernel of $\mathcal{R}_{\nu, \varepsilon}$.

Proof of Lemma 10.3. To shorten notation, in the arguments below we will allow all the implicit constants in the relations \lesssim and \approx to depend on $C_0, C_1, \theta_0, \varepsilon_0$.

Denote

$$\nu = \sum_{Q \in \mathcal{A}_0} \mathcal{H}^n|_{S(Q)},$$

so that $\zeta = \sigma|_{\tilde{F}} + \nu$. Since \mathcal{R}_σ is bounded in $L^2(\sigma)$, it is enough to show that \mathcal{R}_ν is bounded in $L^2(\nu)$. Indeed, the boundedness of both operators implies the boundedness of $\mathcal{R}_{\sigma+\nu}$ in $L^2(\sigma + \nu)$ (see Proposition 2.25 of [To2], for example).

As in (6.3), given $\kappa > 0$, for each $Q \in \mathcal{A}_0$, we consider the set

$$I_\kappa(Q) = \{x \in Q : \text{dist}(x, \text{supp } \sigma \setminus Q) \geq \kappa \ell(Q)\}.$$

By the small boundary condition of Q , the fact that Q is doubling, and that $\sigma(Q) \gtrsim \theta_0 \ell(Q)^n$ (as shown in (10.4)), we deduce there exists some $\kappa > 0$ small enough such that

$$(10.7) \quad \sigma(I_\kappa(Q)) \geq \frac{1}{2} \sigma(Q) \gtrsim \theta_0 \ell(Q)^n.$$

We consider the measure

$$\tilde{\sigma} = \sum_{Q \in \mathcal{A}_0} c_Q \sigma|_{I_\kappa(Q)},$$

with $c_Q = \mathcal{H}^n(S(Q))/\sigma(I_\kappa(Q))$. By (10.7) it follows that the constants c_Q , $Q \in \mathcal{A}_0$, have a uniform bound depending on θ_0 , and thus $\mathcal{R}_{\tilde{\sigma}}$ is bounded in $L^2(\tilde{\sigma})$ (with a norm possibly depending on θ_0). Further, $\nu(S(Q)) = \tilde{\sigma}(Q)$ for each $Q \in \mathcal{A}_0$.

It is clear that, in a sense, $\tilde{\sigma}$ can be considered as an approximation of ν (and conversely). To prove the boundedness of \mathcal{R}_ν in $L^2(\nu)$, we will prove that $\hat{\mathcal{R}}_{\nu,\varepsilon}$ is bounded in $L^2(\nu)$ uniformly on $\varepsilon > 0$ by comparing it to $\hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon}$. First we need to introduce some local and non local operators: given $z \in \bigcup_{Q \in \mathcal{A}_0} S(Q)$, we denote by $S(z)$ the sphere $S(Q)$, $Q \in \mathcal{A}_0$, that contains z . Then we write, for $z \in \bigcup_{Q \in \mathcal{A}_0} S(Q)$,

$$\mathcal{R}_{\nu,\varepsilon}^{loc} f(z) = \hat{\mathcal{R}}_{\nu,\varepsilon}(f \chi_{S(z)})(z), \quad \mathcal{R}_{\nu,\varepsilon}^{nl} f(z) = \hat{\mathcal{R}}_{\nu,\varepsilon}(f \chi_{\mathbb{R}^{n+1} \setminus S(z)})(z).$$

We define analogously $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc} f$ and $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl} f$: given $z \in \bigcup_{Q \in \mathcal{A}_0} Q$, we denote by $Q(z)$ the cell $Q \in \mathcal{A}_0$, that contains z . Then for $z \in \bigcup_{Q \in \mathcal{A}_0} Q$, we set

$$\mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc} f(z) = \hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon}(f \chi_{Q(z)})(z), \quad \mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl} f(z) = \hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon}(f \chi_{\mathbb{R}^{n+1} \setminus Q(z)})(z).$$

It is straightforward to check that $\mathcal{R}_{\nu,\varepsilon}^{loc}$ is bounded in $L^2(\nu)$, and that $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc}$ is bounded in $L^2(\tilde{\sigma})$, both uniformly on ε (in other words, \mathcal{R}_ν^{loc} is bounded in $L^2(\nu)$ and $\mathcal{R}_{\tilde{\sigma}}^{loc}$ is bounded in $L^2(\tilde{\sigma})$). Indeed,

$$\|\mathcal{R}_{\nu,\varepsilon}^{loc} f\|_{L^2(\nu)}^2 = \sum_{Q \in \mathcal{A}_0} \|\chi_{S(Q)} \hat{\mathcal{R}}_{\nu,\varepsilon}(f \chi_{S(Q)})\|_{L^2(\nu)}^2 \lesssim \sum_{Q \in \mathcal{A}_0} \|f \chi_{S(Q)}\|_{L^2(\nu)}^2 = \|f\|_{L^2(\nu)}^2,$$

by the boundedness of the Riesz transforms on $S(Q)$. Using the boundedness of \mathcal{R}_σ in $L^2(\sigma)$, it follows analogously that $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc}$ is bounded in $L^2(\tilde{\sigma})$.

Boundedness of \mathcal{R}_ν^{nl} in $L^2(\nu)$. We must show that \mathcal{R}_ν^{nl} is bounded in $L^2(\nu)$. To this end, we will compare \mathcal{R}_ν^{nl} to $\mathcal{R}_{\tilde{\sigma}}^{nl}$. Observe first that, since $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl} = \hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon} - \mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc}$, and both $\hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon}$ and $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{loc}$ are bounded in $L^2(\tilde{\sigma})$, it follows that $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl}$ is bounded in $L^2(\tilde{\sigma})$ (all uniformly on $\varepsilon > 0$).

Note also that for two different cells $P, Q \in \mathcal{A}_0$, we have

$$(10.8) \quad \text{dist}(S(P), S(Q)) \approx \text{dist}(I_\kappa(P), I_\kappa(Q)) \approx \text{dist}(S(P), I_\kappa(Q)) \approx D(P, Q),$$

where $D(P, Q) = \ell(P) + \ell(Q) + \text{dist}(P, Q)$ and the implicit constants may depend on κ . The arguments to prove this are exactly the same as the ones for (8.7), (8.8) and (8.9), and so we omit them. In particular, (10.8) implies that $(S(P) \cup I_\kappa(P)) \cap (S(Q) \cup I_\kappa(Q)) = \emptyset$, and thus for every $z \in \mathbb{R}^{n+1}$ there is at most one cell $Q \in \mathcal{A}_0$ such that $z \in S(Q) \cup I_\kappa(Q)$, which we denote by $Q(z)$. Hence we can extend $\mathcal{R}_{\nu,\varepsilon}^{nl}$ and $\mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl}$ to $L^2(\tilde{\sigma} + \nu)$ by setting

$$\mathcal{R}_{\nu,\varepsilon}^{nl} f(z) = \hat{\mathcal{R}}_{\nu,\varepsilon}(f \chi_{\mathbb{R}^{n+1} \setminus S(Q(z))})(z), \quad \mathcal{R}_{\tilde{\sigma},\varepsilon}^{nl} f(z) = \hat{\mathcal{R}}_{\tilde{\sigma},\varepsilon}(f \chi_{\mathbb{R}^{n+1} \setminus Q(z)})(z).$$

We will prove below that, for all $f \in L^2(\tilde{\sigma})$ and $g \in L^2(\nu)$ satisfying

$$(10.9) \quad \int_{I_\kappa(P)} f \, d\tilde{\sigma} = \int_{S(P)} g \, d\nu \quad \text{for all } P \in \mathcal{A}_0,$$

we have

$$(10.10) \quad I(f, g) := \int |\mathcal{R}_{\tilde{\sigma}, \varepsilon}^{nl} f - \mathcal{R}_{\nu, \varepsilon}^{nl} g|^2 \, d(\tilde{\sigma} + \nu) \lesssim \|f\|_{L^2(\tilde{\sigma})}^2 + \|g\|_{L^2(\nu)}^2,$$

uniformly on ε . Let us see how the boundedness of \mathcal{R}_ν^{nl} in $L^2(\nu)$ follows from this estimate. As a preliminary step, we show that $\mathcal{R}_\nu^{nl} : L^2(\nu) \rightarrow L^2(\tilde{\sigma})$ is bounded. To this end, given $g \in L^2(\nu)$, we consider a function $f \in L^2(\tilde{\sigma})$ satisfying (10.9) that is constant on each ball B_j . It is straightforward to check that

$$\|f\|_{L^2(\tilde{\sigma})} \leq \|g\|_{L^2(\nu)}.$$

Then from the $L^2(\tilde{\sigma})$ boundedness of $\mathcal{R}_{\tilde{\sigma}}^{nl}$ and (10.10), we obtain

$$\|\mathcal{R}_{\nu, \varepsilon}^{nl} g\|_{L^2(\tilde{\sigma})} \leq \|\mathcal{R}_{\tilde{\sigma}, \varepsilon}^{nl} f\|_{L^2(\tilde{\sigma})} + I(f, g)^{1/2} \lesssim \|f\|_{L^2(\tilde{\sigma})} + \|g\|_{L^2(\nu)} \lesssim \|g\|_{L^2(\nu)},$$

which proves that $\mathcal{R}_\nu^{nl} : L^2(\nu) \rightarrow L^2(\tilde{\sigma})$ is bounded.

It is straightforward to check that the adjoint of $(\mathcal{R}_{\nu, \varepsilon}^{nl})_j : L^2(\nu) \rightarrow L^2(\tilde{\sigma})$ (where $(\mathcal{R}_{\nu, \varepsilon}^{nl})_j$ stands for the j -th component of $(\mathcal{R}_{\nu, \varepsilon}^{nl})$) equals $-(\mathcal{R}_{\tilde{\sigma}, \varepsilon}^{nl})_j : L^2(\tilde{\sigma}) \rightarrow L^2(\nu)$. So by duality we deduce that $\mathcal{R}_{\tilde{\sigma}}^{nl} : L^2(\tilde{\sigma}) \rightarrow L^2(\nu)$ is also bounded.

To prove now the $L^2(\nu)$ boundedness of \mathcal{R}_ν^{nl} , we consider an arbitrary function $g \in L^2(\nu)$, and we construct $f \in L^2(\tilde{\sigma})$ satisfying (10.9) which is constant in each ball P . Again, we have $\|f\|_{L^2(\tilde{\sigma})} \leq \|g\|_{L^2(\nu)}$. Using the boundedness of $\mathcal{R}_{\tilde{\sigma}}^{nl} : L^2(\tilde{\sigma}) \rightarrow L^2(\nu)$ together with (10.10), we obtain

$$\|\mathcal{R}_{\nu, \varepsilon}^{nl} g\|_{L^2(\nu)} \leq \|\mathcal{R}_{\tilde{\sigma}, \varepsilon}^{nl} f\|_{L^2(\nu)} + I(f, g)^{1/2} \lesssim \|f\|_{L^2(\tilde{\sigma})} + \|g\|_{L^2(\nu)} \lesssim \|g\|_{L^2(\nu)},$$

as wished.

It remains to prove that (10.10) holds for $f \in L^2(\tilde{\sigma})$ and $g \in L^2(\nu)$ satisfying (10.9). For $z \in \bigcup_{P \in \mathcal{A}_0} P$, we have

$$|\mathcal{R}_{\tilde{\sigma}, \varepsilon}^{nl} f(z) - \mathcal{R}_{\nu, \varepsilon}^{nl} g(z)| \leq \sum_{P \in \mathcal{A}_0 : P \neq Q(z)} \left| \int \widehat{K}_\varepsilon(z - y) (f(y) \, d\tilde{\sigma}|_{I_\kappa(P)}(y) - g(y) \, d\nu|_{S(P)}(y)) \right|,$$

where $\widehat{K}_\varepsilon(z)$ is the kernel of the ε -regularized n -Riesz transform. By standard estimates, using (10.9) and (10.8), and the smoothness of \widehat{K}_ε , it follows that

$$\begin{aligned} & \left| \int \widehat{K}_\varepsilon(z - y) (f(y) \, d\tilde{\sigma}|_{I_\kappa(P)}(y) - g(y) \, d\nu|_{S(P)}(y)) \right| \\ &= \left| \int_P (\widehat{K}_\varepsilon(z - y) - \widehat{K}_\varepsilon(z - x)) (f(y) \, d\tilde{\sigma}|_{I_\kappa(P)}(y) - g(y) \, d\nu|_{S(P)}(y)) \right| \\ &\lesssim \int \frac{|x - y|}{|z - y|^{n+1}} (|f(y)| \, d\tilde{\sigma}|_{I_\kappa(P)}(y) + |g(y)| \, d\nu|_{S(P)}(y)) \\ &\approx \frac{\ell(P)}{D(Q(z), P)^{n+1}} \int (|f(y)| \, d\tilde{\sigma}|_{I_\kappa(P)}(y) + |g(y)| \, d\nu|_{S(P)}(y)). \end{aligned}$$

Recall that $Q(z)$ stands for the cell Q , $Q \in \mathcal{A}_0$, such that $z \in S(Q) \cup I_\kappa(Q)$.

We consider the operators

$$T_{\tilde{\sigma}}(f)(z) = \sum_{P \in \mathcal{A}_0: P \neq Q(z)} \frac{\ell(P)}{D(Q(z), P)^{n+1}} \int f d\tilde{\sigma}|_{I_\kappa(P)},$$

and T_ν , which is defined in the same way with $\tilde{\sigma}|_{I_\kappa(P)}$ replaced by $\nu|_{S(P)}$. Observe that

$$\begin{aligned} I(f, g) &\leq c \|T_{\tilde{\sigma}}(|f|) + T_\nu(|g|)\|_{L^2(\tilde{\sigma}+\nu)}^2 \\ &\leq 2c \|T_{\tilde{\sigma}}(|f|)\|_{L^2(\tilde{\sigma}+\nu)}^2 + 2c \|T_\nu(|g|)\|_{L^2(\tilde{\sigma}+\nu)}^2 \\ &= 4c \|T_{\tilde{\sigma}}(|f|)\|_{L^2(\tilde{\sigma})}^2 + 4c \|T_\nu(|g|)\|_{L^2(\nu)}^2, \end{aligned}$$

where, for the last equality, we took into account that both $T_{\tilde{\sigma}}(|f|)$ and $T_\nu(|g|)$ are constant on $I_\kappa(P) \cup S(P)$ and that $\tilde{\sigma}(I_\kappa(P)) = \nu(S(P))$ for all $P \in \mathcal{A}_0$.

To complete the proof of (10.10) it is enough to show that $T_{\tilde{\sigma}}$ is bounded in $L^2(\tilde{\sigma})$ and T_ν in $L^2(\nu)$. We only deal with $T_{\tilde{\sigma}}$, since the arguments for T_ν are analogous. We argue by duality again. So we consider non-negative functions $f, h \in L^2(\tilde{\sigma})$ and we write

$$\begin{aligned} \int T_{\tilde{\sigma}}(f) h d\tilde{\sigma} &= \int \left(\sum_{P \in \mathcal{A}_0: P \neq Q(z)} \frac{\ell(P)}{D(P, Q(z))^{n+1}} \int_P f d\tilde{\sigma} \right) h(z) d\tilde{\sigma}(z) \\ &\lesssim \sum_{P \in \mathcal{A}_0} \ell(P) \int_P f d\tilde{\sigma} \int_{\mathbb{R}^{n+1} \setminus P} \frac{1}{(\text{dist}(z, P) + \ell(P))^{n+1}} h(z) d\tilde{\sigma}(z). \end{aligned}$$

From the growth of order n of $\tilde{\sigma}$, it follows easily that

$$\int_{\mathbb{R}^{n+1} \setminus P} \frac{1}{(\text{dist}(z, P) + \ell(P))^{n+1}} h(z) d\tilde{\sigma}(z) \lesssim \frac{1}{\ell(P)} M_{\tilde{\sigma}} h(y) \quad \text{for all } y \in P,$$

where $M_{\tilde{\sigma}}$ stands for the (centered) maximal Hardy-Littlewood operator (with respect to $\tilde{\sigma}$). Then we deduce that

$$\int T_{\tilde{\sigma}}(f) h d\tilde{\sigma} \lesssim \sum_{P \in \mathcal{A}_0} \int_P f(y) M_{\tilde{\sigma}} h(y) d\tilde{\sigma}(y) \lesssim \|f\|_{L^2(\tilde{\sigma})} \|h\|_{L^2(\tilde{\sigma})},$$

by the $L^2(\tilde{\sigma})$ boundedness of $M_{\tilde{\sigma}}$. Thus $T_{\tilde{\sigma}}$ is bounded in $L^2(\tilde{\sigma})$. \square

Proof of the Main Lemma 3.1. By Lemmas 10.2, 10.3, and the Nazarov-Tolsa-Volberg theorem of [NTov1], ζ is a uniformly rectifiable measure. So it only remains to note that the set $\Gamma := \tilde{F}$ satisfies the required properties from the Main Lemma: it is contained in $\text{supp } \zeta$, which is uniformly rectifiable and, by (10.2), $\mu(\Gamma) = \sigma(\tilde{F}) \geq \frac{\varepsilon_0}{2} \mu(Q_0)$. \square

REFERENCES

- [AMT] J. Azzam, M. Mourgoglou and X. Tolsa, *Mutual absolute continuity of interior and exterior harmonic measure implies rectifiability*, arXiv:1602.01397 (2016).
- [BCGJ] C. J. Bishop, L. Carleson, J. B. Garnett, and P. W. Jones, *Harmonic measures supported on curves*, Pacific J. Math. **138** (1989), no. 2, 233–236.
- [Bi1] C. J. Bishop, *A characterization of Poissonian domains*, Ark. Mat. **29**(1) (2006), 1–24.

- [Bi2] C. J. Bishop, *Some questions concerning harmonic measure*. Partial differential equations with minimal smoothness and applications (Chicago, IL, 1990) IMA Vol. Math. Appl., vol. 42, Springer, New York, 1992, pp. 89–97.
- [Da] G. David, *Unrectifiable 1-sets have vanishing analytic capacity*. Revista Mat. Iberoamericana 14(2) (1998), 369–479.
- [DM] G. David and P. Mattila. *Removable sets for Lipschitz harmonic functions in the plane*. Rev. Mat. Iberoamericana 16(1) (2000), 137–215.
- [DS1] G. David and S. Semmes, *Singular integrals and rectifiable sets in R_n : Beyond Lipschitz graphs*, Astérisque No. 193 (1991).
- [DS2] G. David and S. Semmes, *Analysis of and on uniformly rectifiable sets*, Mathematical Surveys and Monographs, 38, American Mathematical Society, Providence, RI, 1993.
- [ENV] V. Eiderman, F. Nazarov and A. Volberg. *The s -Riesz transform of an s -dimensional measure in \mathbb{R}^2 is unbounded for $1 < s < 2$* , to appear in J. Anal. Math.
- [JK] D. S. Jerison and C.E. Kenig, *Boundary behavior of harmonic functions in nontangentially accessible domains*, Adv. in Math. 46 (1982), no. 1, 80–147.
- [KPT] C. E. Kenig, D. Preiss, and T. Toro, *Boundary structure and size in terms of interior and exterior harmonic measures in higher dimensions*, J. Amer. Math. Soc. **22** (2009), no. 3, 771–796.
- [Lé] J.C. Léger, *Menger curvature and rectifiability*, Ann. of Math. 149 (1999), 831–869.
- [Me] M.S. Melnikov. *Analytic capacity: discrete approach and curvature of a measure*. Sbornik: Mathematics 186(6) (1995), 827–846.
- [MV] M.S. Melnikov and J. Verdera. *A geometric proof of the L^2 boundedness of the Cauchy integral on Lipschitz graphs*. Internat. Math. Res. Notices (1995), 325–331.
- [NTov1] F. Nazarov, X. Tolsa and A. Volberg, *On the uniform rectifiability of AD-regular measures with bounded Riesz transform operator: the case of codimension 1*, Acta Math. **213** (2014), no. 2, 237–321.
- [NTov2] F. Nazarov, X. Tolsa and A. Volberg. *The Riesz transform Lipschitz harmonic functions*. Publ. Mat. 58 (2014), 517–532.
- [Pa] H. Pajot, *Théorème de recouvrement par des ensembles Ahlfors-réguliers et capacité analytique*, C. R. Acad. Sci. Paris Sér. I Math. 323(2) (1996), 133–135.
- [To1] X. Tolsa, *Uniform rectifiability, Calderón-Zygmund operators with odd kernel, and quasiorthogonality*. Proc. London Math. Soc. 98(2) (2009), 393–426.
- [To2] X. Tolsa. *Analytic capacity, the Cauchy transform, and non-homogeneous Calderón-Zygmund theory*, volume 307 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2014.
- [To3] X. Tolsa. *Rectifiable measures, square functions involving densities, and the Cauchy transform*. To appear in Mem. Amer. Math. Soc.
- [Vo] A. Volberg, *Calderón-Zygmund capacities and operators on nonhomogeneous spaces*. CBMS Regional Conf. Ser. in Math. 100, Amer. Math. Soc., Providence, 2003.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA (BARCELONA, CATALONIA)

E-mail address: ddgirela@gmail.com

ICREA, UNIVERSITAT AUTÒNOMA DE BARCELONA AND BGSMATH
DEPARTAMENT DE MATEMÀTIQUES, 08193 BELLATERRA (BARCELONA, CATALONIA)

E-mail address: xtolsa@mat.uab.cat