

# INTEGRAL HOMOLOGY 3-SPHERES AND THE JOHNSON FILTRATION

WOLFGANG PITTSCH

ABSTRACT. The mapping class group  $\mathcal{M}_{g,1}$  of an oriented surface  $\Sigma_{g,1}$  of genus  $g$  with one boundary component has a natural decreasing filtration  $\mathcal{M}_{g,1} \supset \mathcal{M}_{g,1}(1) \supset \mathcal{M}_{g,1}(2) \supset \mathcal{M}_{g,1}(3) \supset \dots$ , where  $\mathcal{M}_{g,1}(k)$  is the kernel of the action of  $\mathcal{M}_{g,1}$  on the  $k^{\text{th}}$  nilpotent quotient of  $\pi_1(\Sigma_{g,1})$ . Using a tree Lie algebra approximating the graded Lie algebra  $\bigoplus_k \mathcal{M}_{g,1}(k)/\mathcal{M}_{g,1}(k+1)$  we prove that any integral homology sphere of dimension 3 has for some  $g$  a Heegaard decomposition of the form  $M = \mathcal{H}_g \amalg_{\iota_g \phi} -\mathcal{H}_g$ , where  $\phi \in \mathcal{M}_{g,1}(3)$  and  $\iota_g$  is such that  $\mathcal{H}_g \amalg_{\iota_g} -\mathcal{H}_g = S^3$ . This proves a conjecture due to S. Morita [11] and shows that the "core" of the Casson invariant introduced in [11] is indeed the Casson invariant.

## 1. INTRODUCTION

In the early 80's, A. Casson succeeded to lift the classical Rohlin invariant of integral homology 3-spheres, which is  $\mathbf{Z}/2\mathbf{Z}$ -valued, to a  $\mathbf{Z}$ -valued invariant. Later S. Morita [11] used the theory of Heegaard splittings combined with techniques from group cohomology to give a new construction of this invariant related to algebraic properties of the mapping class group of an oriented surface.

More precisely, let  $\Sigma_g$  be a closed oriented surface of genus  $g$ . For technical reasons, fix a small disc  $D^2$  on  $\Sigma_g$  and consider the mapping class group  $\mathcal{M}_{g,1}$ , i.e. the group of orientation preserving diffeomorphisms of  $\Sigma_g$  which are the

---

2000 *Mathematics Subject Classification*. Primary 57M99; Secondary 20F38, 20F12.


The author is supported MEC grant MTM2004-06686 and by the program Ramón y Cajal, MEC, Spain.

identity on  $D^2$  modulo isotopies which are also the identity on  $D^2$ . Let  $h : \Sigma_g \rightarrow \mathbf{S}^3$  be a fixed Heegaard embedding of  $\mathbf{S}^3$ , and for  $\phi \in \mathcal{M}_{g,1}$  let  $\mathbf{S}_\phi^3$  denote the manifold obtained by cutting  $\mathbf{S}^3$  along  $h(\Sigma_g)$  and gluing back the two pieces by a map representing  $\phi$ . It is a classical fact ([7], chap. 8) that the diffeomorphism class of  $\mathbf{S}_\phi^3$  is well-defined, and that any diffeomorphism class of closed oriented 3-manifolds can be obtained in this way.

The mapping class group has a natural filtration by normal subgroups, called the Johnson filtration :  $\cdots \mathcal{M}_{g,1}(k+1) \subset \mathcal{M}_{g,1}(k) \subset \cdots \mathcal{M}_{g,1}$ , and it is a natural question to try to relate this filtration to the theory of Heegaard splittings. For instance, if we restrict the map  $\phi \mapsto \mathbf{S}_\phi^3$  to the group  $\mathcal{M}_{g,1}(1)$ , also called the Torelli group, what we get are exactly the integral homology 3-spheres  $\mathcal{S}(3)$ . In [11] S. Morita proved that the elements of  $\mathcal{M}_{g,1}(2)$  suffice to obtain the whole set  $\mathcal{S}(3)$ . Then he proved that the composite of the map  $\mathcal{M}_{g,1}(2) \rightarrow \mathcal{S}(3)$  with the Casson invariant  $\mathcal{S}(3) \rightarrow \mathbf{Z}$ , is a homomorphism of groups, and moreover that it is the sum of two morphisms  $d$  and  $q$ . The first morphism  $d$ , which is related to cohomological properties of the mapping class group [16], was called by S. Morita “the core of the Casson invariant”, the second term  $q$  appeared as a correcting term and was shown by S. Morita to vanish on the subgroup  $\mathcal{M}_{g,1}(3)$ . Therefore he asked if this latter group suffices to construct all integral homology spheres. The purpose of this paper is to give an affirmative answer to this question :

**Main Theorem.** *Any integral homology sphere is diffeomorphic to  $\mathbf{S}_\phi^3$  for some  $\phi \in \mathcal{M}_{g,1}(3)$  with  $g \geq 9$ .*

Our starting point is a classical theorem of Singer. Roughly speaking it asserts that the sentence “the manifolds  $\mathbf{S}_\phi^3$  and  $\mathbf{S}_\psi^3$  are diffeomorphic” can be algebraically translated as “the mapping classes  $\phi$  and  $\psi$  lie in the same double coset in  $\mathcal{A}_{g+h,1} \backslash \mathcal{M}_{g+h,1} / \mathcal{B}_{g+h,1}$  for all sufficiently large  $h$ ”, where the subgroups  $\mathcal{A}_{g,1}$  and  $\mathcal{B}_{g,1}$  are known. Unfortunately, this algebraic statement is difficult to handle in practice. Therefore, we first construct an injective



 FIGURE 1. The torus  $\mathcal{H}_{1,2}$ 

homomorphism  $\tilde{\tau}_2$  from  $\mathcal{M}_{g,1}(2)/\mathcal{M}_{g,1}(3)$  to a commutative group  $\mathcal{A}_2(N_1)$ . By commutativity, to check that an element in  $\mathcal{M}_{g,1}(2)$  is equivalent to some element in  $\mathcal{M}_{g,1}(3)$  it suffices then to prove that  $\tilde{\tau}_2(\phi)$  belongs to the subgroup  $\text{Can}(\mathcal{A}_2(N_1)) \subset \mathcal{A}_2(N_1)$  generated by the images of  $(\mathcal{M}_{g,1}(2) \cap \mathcal{A}_{g,1})/\mathcal{M}_{g,1}(3)$  and  $(\mathcal{M}_{g,1}(2) \cap \mathcal{B}_{g,1})/\mathcal{M}_{g,1}(3)$ .

This paper is organised as follows. In section 2 we have compiled some standard facts about mapping class groups, the Johnson filtration and Heegaard decompositions. In section 3 we introduce Lie algebra structures on the graded group  $\oplus_k \mathcal{M}_{g,1}(k)/\mathcal{M}_{g,1}(k+1)$  and on the natural target of the Johnson's homomorphisms  $\oplus_k D_k(N_1)$ . We also introduce a graded injective homomorphism  $\oplus_k \tilde{\tau}_k : \oplus_k D_k(N_1) \rightarrow \oplus_k \mathcal{A}_k(N_1)$  into a Lie algebra of labelled trees. In section 4 we give some properties of the group  $\text{Can}(\mathcal{A}_2(N_1))$ , in particular we study its invariance under some natural actions. The final section 5 is devoted to the proof of the main theorem.

## 2. PRELIMINARIES

**2.1. Johnson's homomorphisms.** Let  $\mathcal{H}_{1,2}$  be an oriented solid torus with two small discs  $D_1^2$  and  $D_2^2$  on its boundary intersecting in one point  $x_0$ , and fix two based loops on the boundary: a parallel  $\alpha$  and a meridian  $\beta$ , as in Figure 1.

FIGURE 2. Model for  $\Sigma_{g,1}$ 

Gluing  $g$  copies of  $\mathcal{H}_{1,2}$  by identifying  $(D_2^2, x_0)$  on copy  $i$  to  $(D_1^2, x_0)$  on copy  $i + 1$  we get an oriented genus  $g$  handlebody  $\mathcal{H}_g$  with a small disc  $D^2$  on its surface (the  $D_2^2$  of the  $g^{\text{th}}$  term of the sum). Let  $\Sigma_g$  denote the boundary of  $\mathcal{H}_g$ , and  $\Sigma_{g,1} = \Sigma_g \setminus \overset{\circ}{D}^2$ , these two oriented surfaces are naturally pointed by  $x_0$ . By construction the  $2g$  curves  $\alpha_i, \beta_i$  (see Figure 2), which are the images of the different curves  $\alpha$  and  $\beta$ , are free generators of the free group  $\pi_1(\Sigma_{g,1}, x_0)$ .

Consider the central series of the fundamental group  $\pi_1(\Sigma_{g,1}, x_0)$ , inductively defined by  $\Gamma_0 = \pi_1(\Sigma_{g,1}, x_0)$ , and  $\forall k \geq 1 \Gamma_{k+1} = [\Gamma_k, \Gamma_0]$ . Denote by  $N_k$  the nilpotent quotient  $\Gamma_0/\Gamma_k$ . For instance  $N_1$  is isomorphic to the abelianization of  $\pi_1(\Sigma_{g,1}, x_0)$ .

Let  $\mathcal{M}_{g,1}$  denote the group of isotopy classes of diffeomorphisms which fix  $D^2$  point-wise modulo isotopies which also fix  $D^2$  point-wise, also called mapping class group. Note that as they fix  $D^2$  these mapping classes preserve the orientation.

The natural action of  $\mathcal{M}_{g,1}$  on  $\pi_1(\Sigma_{g,1}, x_0)$  was described by Nielsen ([13], [14], [15]) :

**Theorem 2.1.** *The canonical morphism*

$$\mathcal{M}_{g,1} \longrightarrow \text{Aut}(\pi_1(\Sigma_{g,1}, x_0))$$

*is injective. Its image consists of those automorphisms which fix  $\prod_{i=1}^g [\alpha_i, \beta_i]$ .*

As the commutator subgroups  $\Gamma_k$  are preserved by any automorphism of  $\pi_1(\Sigma_{g,1}, x_0)$ , the above action induces a family of compatible morphisms :

$$\begin{array}{ccc} \mathcal{M}_{g,1} & & \\ \downarrow & \searrow & \\ \text{Aut}N_{k+1} & \longrightarrow & \text{Aut}N_k. \end{array}$$

There is a canonical surjection  $N_{k+1} \longrightarrow N_k$  whose kernel  $\mathcal{L}_{k+1}$  is known to be the centre of  $N_{k+1}$  ([10] p. 343). As the centre of a group is preserved by any automorphism one can prove (cf. [17]) that the central extension

$$0 \longrightarrow \mathcal{L}_{k+1} \longrightarrow N_{k+1} \longrightarrow N_k \longrightarrow 1,$$

induces an exact sequence

$$0 \longrightarrow \text{Hom}(N_1, \mathcal{L}_{k+1}) \longrightarrow \text{Aut}(N_{k+1}) \longrightarrow \text{Aut}(N_k) \longrightarrow 1.$$

Let  $\mathcal{M}_{g,1}(k)$  denote the kernel of the canonical homomorphism  $\mathcal{M}_{g,1} \longrightarrow \text{Aut}(N_k)$ . We have an induced morphism, called the Johnson homomorphism  $\tau_k : \mathcal{M}_{g,1}(k) \longrightarrow \text{Hom}(N_1, \mathcal{L}_{k+1})$ . Explicitly,  $\tau_k(f)$  is given as follows :

If  $c \in N_1$  lifts to  $\gamma \in \pi_1(\Sigma_{g,1}, x_0)$ , then

$$\tau_k(f)(c) = f(\gamma)\gamma^{-1} \text{ mod } \Gamma_{k+1}.$$

The mapping class group acts on the kernels  $\mathcal{M}_{g,1}(k)$  by conjugation and on  $\text{Hom}(N_1, \mathcal{L}_{k+1})$  through the conjugation action by  $\text{Aut}(N_{k+1})$ . These actions are obviously compatible, so that :

**Proposition 2.1.** *The collection of Johnson homomorphisms reassemble into a monomorphism of graded groups and  $\mathcal{M}_{g,1}$ -modules*

$$\bigoplus_{k \geq 1} \tau_k : \bigoplus_{k \geq 1} \mathcal{M}_{g,1}(k) / \mathcal{M}_{g,1}(k+1) \longrightarrow \bigoplus_{k \geq 1} \text{Hom}(N_1, \mathcal{L}_{k+1}).$$

Recall that the algebraic intersection of closed paths on  $\pi_1(\Sigma_{g,1}, x_0)$  induces a symplectic form  $\omega : \Lambda^2 N_1 \rightarrow \mathbf{Z}$ , where we identify  $N_1$  with  $H_1(\Sigma_{g,1})$  via the Hurewicz homomorphism. The image of the free basis  $\alpha_1, \beta_1 \dots \alpha_g, \beta_g$  in  $N_1$  is a symplectic basis  $a_1, b_1 \dots a_g, b_g$  for  $\omega$  and the image of the canonical

morphism  $\mathcal{M}_{g,1} \rightarrow \text{Aut}N_1$  is known to be the associated symplectic group  $\text{Sp}\omega$  (see [10] p. 178, the proof for the one boundary component case is the same). In the sequel we will denote the two Lagrangians generated by  $\{a_i \mid 1 \leq i \leq g\}$  and  $\{b_i \mid 1 \leq i \leq g\}$  respectively by  $L_A$  and  $L_B$ . By definition we have a decomposition by  $\omega$ -dual Lagrangians  $N_1 = L_A \oplus L_B$ .

**2.2. Heegaard splittings.** Let  $\mathcal{V}(3)$  denote the set of all diffeomorphism classes of closed oriented 3-manifolds and  $\mathcal{S}(3)$  the subset of all integral homology spheres. By virtue of the classical theorem of Heegaard splittings, one can describe these two sets as the direct limit of a system of double cosets of mapping class groups.

Our inductive construction  $\mathcal{H}_{g+1} = \mathcal{H}_g \sharp \mathcal{H}_{1,2}$  makes  $\Sigma_{g,1}$  a canonical sub-surface of  $\Sigma_{g+1,1}$ , thus extending an orientation preserving diffeomorphism of  $\Sigma_{g,1}$  by the identity over its complement we get a well defined morphism  $\mathcal{M}_{g,1} \rightarrow \mathcal{M}_{g+1,1}$  compatible with the action on the fundamental group. The Nielsen theorem implies that this morphism is injective.

Consider the following morphism :

$$\begin{aligned} \tilde{\iota}_g : \pi_1(\Sigma_{g,1}, x_0) &\longrightarrow \pi_1(\Sigma_{g,1}, x_0) \\ \alpha_i &\longmapsto \beta_i^{-1} \\ \beta_i &\longmapsto \beta_i \alpha_i \beta_i^{-1}. \end{aligned}$$

It fixes  $(\prod_{i=1}^g [\alpha_i, \beta_i])$  and so corresponds to an orientation preserving diffeomorphism  $\Sigma_{g,1} \rightarrow \Sigma_{g,1}$ . Let  $\iota_g : \Sigma_{g,1} \rightarrow -\Sigma_{g,1}$  be the composite  $-Id \circ \tilde{\iota}_g$  where  $-\Sigma_{g,1}$  denotes the surface  $\Sigma_{g,1}$  with opposite orientation.

**Lemma 2.1.** *The oriented manifold  $\mathcal{H}_g \amalg_{\iota_g} -\mathcal{H}_g$  obtained by identifying the boundaries of  $\mathcal{H}_g$  and  $-\mathcal{H}_g$  via  $\iota_g$  is diffeomorphic to  $\mathbf{S}^3$ . Here  $-\mathcal{H}_g$  denotes  $\mathcal{H}_g$  with opposite orientation.*

*Proof.* Almost by definition  $\tilde{\iota}_{g+1} = \tilde{\iota}_g \# \tilde{\iota}_1$ , so  $\mathcal{H}_g \amalg_{\iota_g} -\mathcal{H}_g$  is diffeomorphic to the connected sum of  $g$  copies of  $\mathcal{H}_1 \amalg_{\iota_1} -\mathcal{H}_1$ . The diffeomorphism  $\tilde{\iota}_1$  exchanges the meridian  $\alpha$  and the parallel  $\beta$  and it is well known that this construction yields  $\mathbf{S}^3$ . Q. E. D.

Therefore we have a map  $\mathcal{M}_{g,1} \rightarrow \mathcal{V}(3)$ , given by  $\phi \mapsto \mathbf{S}_\phi^3 = \mathcal{H}_g \amalg_{\iota_g \circ \phi} -\mathcal{H}_g$ , which is compatible with the monomorphisms  $\mathcal{M}_{g,1} \rightarrow \mathcal{M}_{g+1,1}$ , thus yielding a well defined and surjective map

$$\lim_{g \rightarrow +\infty} \mathcal{M}_{g,1} \longrightarrow \mathcal{V}(3).$$

Note that as we identify  $\mathbf{S}^3$  with  $\mathcal{H}_g \amalg_{\iota_g} -\mathcal{H}_g$  the surface  $\Sigma_g$  becomes the common boundary of the two solid handlebodies  $\mathcal{H}_g$ , and  $-\mathcal{H}_g$ . Denote the image of the restriction maps  $\text{Diff}(\mathcal{H}_g, \text{rel. } D^2) \rightarrow \text{Diff}(\Sigma_g, \text{rel. } D^2)$  and  $\text{Diff}(-\mathcal{H}_g, \text{rel. } D^2) \rightarrow \text{Diff}(\Sigma_g, \text{rel. } D^2)$  in  $\mathcal{M}_{g,1}$  respectively by  $\mathcal{B}_{g,1}$  and  $\mathcal{A}_{g,1}$ . It is easily seen that  $\mathcal{A}_{g,1} = \iota_g^{-1} \mathcal{B}_{g,1} \iota_g$ .

**Definition 1.** *Two mapping classes  $\phi, \psi \in \mathcal{M}_{g,1}$  will be called equivalent, denoted by  $\phi \approx \psi$ , if there exists  $f_a \in \mathcal{A}_{g,1}$  and  $f_b \in \mathcal{B}_{g,1}$  such that  $\phi = f_a \circ \psi \circ f_b$ . This is equivalent to say that  $\phi$  and  $\psi$  lie in the same double coset in  $\mathcal{A}_{g,1} \backslash \mathcal{M}_{g,1} / \mathcal{B}_{g,1}$ .*

The relation  $\approx$  is clearly an equivalence relation. It is a classical fact that if  $\phi \approx \psi$  then  $\mathbf{S}_\phi^3$  is diffeomorphic to  $\mathbf{S}_\psi^3$  (cf. [7] chap. 8). Moreover two equivalent maps in  $\mathcal{M}_{g,1}$  are also equivalent when considered as maps in  $\mathcal{M}_{g+1,1}$ . The key result in the theory of Heegaard splittings is :

**Theorem 2.2.** *The map*

$$\begin{aligned} \lim_{g \rightarrow +\infty} \mathcal{M}_{g,1} / \approx & \longrightarrow \mathcal{V}(3) \\ \phi & \longmapsto \mathbf{S}_\phi^3 \end{aligned}$$

*is bijective.*

For historical proofs of this theorem we refer the reader to Redemeister [18] or Singer [19]. For a shorter proof see for instance Craggs [2]. The historical proofs of this theorem by are somewhat obscure, for a modern proof

Using the Mayer-Vietoris sequence one can prove that the image of the restriction of this map to the equivalence classes of maps in  $\mathcal{M}_{g,1}(1)$  is precisely  $\mathcal{S}(3)$ . This result was refined by S. Morita who proved :

**Theorem 2.3.** [11] *The map*

$$\lim_{g \rightarrow +\infty} \mathcal{M}_{g,1}(2)/\approx \longrightarrow \mathcal{S}(3)$$

*is a bijection.*

**Remark 2.1.** *In the above theorem  $\mathcal{M}_{g,1}(2)/\approx$  stands for the equivalence classes under the equivalence relation  $\approx$  restricted to  $\mathcal{M}_{g,1}(2)$ . In particular if  $\phi \in \mathcal{M}_{g,1}(2)$  and  $\psi \in \mathcal{M}_{g,1}(2)$  are equivalent, that is  $\phi = f_a \circ \psi \circ f_b$  for some  $(f_a, f_b) \in \mathcal{A}_{g,1} \times \mathcal{B}_{g,1}$ , we do not require  $f_a$  nor  $f_b$  to belong to  $\mathcal{M}_{g,1}(2)$ .*

Our aim is to prove that the theorem still holds when we replace  $\mathcal{M}_{g,1}(2)$  by  $\mathcal{M}_{g,1}(3)$ , to do so we first need to enrich the structure of the Johnson homomorphisms.

### 3. A LIE ALGEBRA APPROXIMATING THE MAPPING CLASS GROUP

Both graded groups  $\oplus_{k \geq 1} \mathcal{M}_{g,1}(k)$  and  $\oplus_{k \geq 1} \text{Hom}(N_1, \mathcal{L}_{k+1})$  have natural structures of graded Lie algebras, that these structures are compatible with the Johnson homomorphism was first observed by S. Morita in [12]. Let us briefly recall the involved structures, for the details we refer the reader to [12].

The Lie algebra structure on  $\oplus_{k \geq 1} \mathcal{M}_{g,1}(k)$  is induced by the commutator bracket  $f, g \mapsto [f, g] = f \circ g \circ f^{-1} \circ g^{-1}$ , and it is clearly compatible with the action by  $\mathcal{M}_{g,1}$ .

The algebra  $\oplus_{k \geq 1} \mathcal{L}_{k+1}$  is the graded subalgebra of elements of degree  $k \geq 2$  in the free graded Lie algebra  $\mathcal{L}$  generated by  $N_1$ . Recall that a derivation of a Lie algebra  $L$  is a homomorphism of abelian groups  $d : L \rightarrow L$ , such that

for any  $u, v \in L$ ,  $d([u, v]) = [d(u), v] + [u, d(v)]$ . A derivation of a graded Lie algebra is graded of degree  $s$  if for all  $k \geq 0$   $d(L_k) \subset L_{k+s}$ . In particular, the abelian group of graded derivations of degree  $\geq 1$  of  $\mathcal{L}$  is isomorphic to  $\bigoplus_{k \geq 1} \text{Hom}(N_1, \mathcal{L}_{k+1})$  (note that by definition an element  $f \in \text{Hom}(N_1, \mathcal{L}_{k+1})$  is of degree  $k$ ). There is a natural graded product on the group of graded derivations :

**Definition 2.** *Let  $f$  and  $g$  be two derivations of degree  $k$  and  $l$ , then  $f\{g\}$  is the unique derivation of degree  $k+l$  given by  $f\{g\}(u) = f(g(u))$  for all  $u \in N_1$ .*

The anti-symmetrisation of this product  $[f, g] = f\{g\} - g\{f\}$  induces a Lie bracket and in our case it is compatible with the action of  $\mathcal{M}_{g,1}$ , as was proved by S. Morita.

Recall that we have a symplectic form  $\omega : \Lambda^2 N_1 \rightarrow \mathbf{Z}$ , as usual this form induces an isomorphism of  $\text{Sp}\omega$ -modules

$$\begin{aligned} N_1 &\rightarrow \text{Hom}(N_1, \mathbf{Z}) \\ a &\mapsto (b \rightsquigarrow \omega(a, b)). \end{aligned}$$

This in turn induces isomorphisms of  $\text{Sp}\omega$ -modules  $\text{Hom}(N_1, \mathcal{L}_{k+1}) \simeq N_1 \otimes \mathcal{L}_{k+1}$ . Let  $D_k(N_1)$  be the kernel of the Lie bracket  $N_1 \otimes \mathcal{L}_{k+1} \xrightarrow{[\cdot, \cdot]} \mathcal{L}_{k+2}$ . Denote by  $\theta = \sum_{i=1}^g a_i \wedge b_i \in \mathcal{L}_2$  the class of  $\prod_{i=1}^g [\alpha_i, \beta_i]$ . A typical derivation  $\phi \in \text{Hom}(N_1, \mathcal{L}_k)$ , such that  $\phi(a_i) = l_{a_i}$  and  $\phi(b_i) = l_{b_i}$  sends  $\theta$  to  $\phi(\theta) = \sum_{i=1}^g [l_{a_i}, b_i] + [a_i, l_{b_i}]$ . In particular  $D(N_1) = \bigoplus_{k=1}^{\infty} D_k(N_1)$  is the sub-Lie algebra of  $\bigoplus_{k \geq 1} \text{Hom}(N_1, \mathcal{L}_{k+1})$  of elements that send  $\theta$  to 0. As  $\mathcal{M}_{g,1}$  fixes  $\prod_{i=1}^g [\alpha_i, \beta_i]$ , the images of the Johnson homomorphisms lie in  $D(N_1)$ .

**Theorem 3.1** ([12]). *The Johnson homomorphisms reassemble into a graded monomorphism of graded Lie algebras and  $\mathcal{M}_{g,1}$ -modules :*

$$\bigoplus_{k \geq 1} \tau_k : \bigoplus_{k \geq 1} \mathcal{M}_{g,1}(k) / \mathcal{M}_{g,1}(k+1) \longrightarrow \bigoplus_{k \geq 1} D_k(N_1).$$

Moreover, the two structures are mutually compatible.

By definition of  $D(N_1)$  the action of  $\mathcal{M}_{g,1}$  factors through  $\text{Sp}\omega$  and therefore it is also the case for the action on  $\bigoplus_{k \geq 1} \mathcal{M}_{g,1}(k) / \mathcal{M}_{g,1}(k+1)$ .

**3.1. Interpretation by tree Lie algebras.** The Lie algebra  $D(N_1)$  is still difficult to handle for explicit computations. To overcome this problem we embed it into a Lie algebra generated by uni-trivalent trees, labelled by  $N_1$ . Most of the material in this paragraph is known, we refer the interested reader to Levine's paper [9] for a more detailed investigation of the relationship between  $D(N_1)$  and tree Lie algebras. Nevertheless we will give a somewhat extended discussion as our proof makes full use of the involved structures.

In the sequel a *tree* will always mean a finite, uni-trivalent tree equipped with a cyclic ordering of its trivalent vertices. Recall that a *uni-trivalent tree* is a tree whose vertices are all of index one or three. Unless otherwise specified, the trivalent vertices of the trees we draw are oriented according to the usual trigonometric orientation of the plane. A *rooted tree* is a tree with a distinguished univalent vertex called the root and represented by  $*$ . A *labelled tree* with labels in the set  $E$  is a tree  $T$  together with a function  $l : V_1(T) \rightarrow E$ , where  $V_1(T)$  stands for the set of univalent vertices excepting the root in case  $T$  is rooted. The label of the univalent vertex  $x$  will be denoted by  $l_x$ .

**Definition 3.** Let  $\mathcal{A}(N_1)$  (resp.  $\mathcal{A}^r(N_1)$ ) denote the free abelian group generated by the uni-trivalent, oriented (resp. rooted) trees with labels in  $N_1$ , modulo multi-linearity with respect to the labels, and the relations IHX, AS and " $T \wedge T = 0$ " of Figure 3.

In relation " $T \wedge T = 0$ ",  $S$  may be either a tree, a label or the root.

**Remark 3.1.** Usually (cf. [1] [5]), the third relation is omitted as it is a consequence of the previous one when multiplication by two is an isomorphism on the group of labels, e.g. when  $N_1$  is replaced by  $N_1 \otimes \mathbf{Q}$ .

Both groups  $\mathcal{A}^r(N_1)$  and  $\mathcal{A}(N_1)$  are naturally graded by the internal degree of trees, that is the number of trivalent vertices in a tree. In the case of  $\mathcal{A}(N_1)$ , we will only consider the subgroup of elements of degree at least 1, and still denote it by  $\mathcal{A}(N_1)$ . The following Proposition is well-known :




FIGURE 3. Relations among trees




FIGURE 4. Lie bracket in  $\mathcal{A}^r(N_1)$

**Proposition 3.1.** *The bracket in Figure 4 endows  $\mathcal{A}^r(N_1)$  with a graded Lie algebra structure, compatible with the action of  $\text{Sp}\omega$ . As  $\mathcal{A}_0^r(N_1) \simeq N_1$ , there is a canonical epimorphism of graded Lie algebras and  $\text{Sp}\omega$ -modules  $\mathcal{L} \rightarrow \mathcal{A}^r(N_1)$ , which is in fact an isomorphism.*

In the sequel we identify the graded Lie algebras  $\mathcal{L}$  and  $\mathcal{A}^r(N_1)$ . There is a fundamental operation on trees given by gluing two univalent vertices of two distinct trees. Consider two trees  $S$  and  $T$ , where  $S$  is not rooted but  $T$  may be, and choose two univalent vertices  $x \in V_1(S)$  and  $w \in V_1(T)$  (recall that by definition the root  $*$  does not belong to  $V_1(T)$ ). We forget the labelling and glue these two vertices to get a new labelled tree  $S - xw - T$ , naturally rooted if  $T$  was. This gluing operation satisfies :

(1) Symmetry :

$$S - xw - T = T - wx - S.$$

(2) Associativity : If at least two of the three trees  $S, T, R$  are non-rooted

$$S - xw - (T - vy - R) = (S - xw - T) - vy - R.$$

Extending by linearity, this allows to define :

(1) A bracket on  $\mathcal{A}(N_1)$ , which is given on trees by :

$$[S, T] = \sum_{x, y \in V_1(S) \times V_1(T)} \omega(l_x \wedge l_y) S - xy - T.$$

(2) A natural graded linear action of  $\mathcal{A}(N_1)$  on  $\mathcal{A}^r(N_1)$ , which is given on trees by :

$$S\{T\} = \sum_{x, y \in V_1(S) \times V_1(T)} \omega(l_x \wedge l_y) S - xy - T.$$

It is readily seen from the definition that this is an action by graded derivations of degree at least 1. We call the resulting linear map :

$$\mathcal{A}(N_1) \longrightarrow \bigoplus_{k \geq 1} \text{Hom}(N_1, \mathcal{L}_{k+1}) \simeq N_1 \otimes \bigoplus_{k \geq 1} \mathcal{L}_{k+1}$$

the expansion map. On a tree  $T \in \mathcal{A}_k(N_1)$  the expansion map is given by :

$$T \longmapsto \sum_{x \in v_1(T)} l_x \otimes T^x,$$

where  $T^x$  denotes the tree  $T$  rooted by its vertex  $x$ .

In fact more is true, namely the above bracket is a Lie bracket and turns the expansion map into a Lie algebra homomorphism. This was first observed by Garoufalidis and Levine [3] for trees labelled by  $N_1 \otimes \mathbf{Q}$ . The proof is straightforward, for the details we refer the interested reader to [6].

**Theorem 3.2.**

- *The bracket on  $\mathcal{A}(N_1)$  induces a graded Lie algebra structure compatible with the action of  $\text{Sp}\omega$  on the labels.*
- *The expansion map induces a graded homomorphism of Lie algebras and  $\text{Sp}\omega$ -modules. Its image is contained in  $D(N_1)$ .*

In degree 1 the expansion map is an isomorphism, the group  $D_1(N_1)$  is isomorphic to  $\Lambda^3 N_1$ , and the expansion map is given by :

$$\begin{array}{ccc} \mathcal{A}_1(N_1) & \longrightarrow & D_1(N_1) = \Lambda^3 N_1 \\ \begin{array}{c} u \\ \diagdown \quad \diagup \\ v \quad w \end{array} & \longmapsto & u \wedge v \wedge w. \end{array}$$

In higher degrees the expansion map is no more an isomorphism. Nevertheless there is still a linear homomorphism  $D(N_1) \rightarrow \mathcal{A}(N_1)$  which is an inverse after taking tensor product with the rationals. It is compatible with the  $\mathrm{Sp}\omega$ -action but not with the Lie bracket.

Recall that for each  $k \geq 1$   $\mathrm{Hom}(N_1, \mathcal{L}_{k+1}) \simeq N_1 \otimes \mathcal{L}_{k+1}$  as an  $\mathrm{Sp}\omega$ -module. Moreover we have identified  $\mathcal{L}_{k+1}$  with  $\mathcal{A}_{k+1}^r(N_1)$ . Therefore we view  $D_k(N_1) \subset \mathrm{Hom}(N_1, \mathcal{L}_{k+1})$  as an  $\mathrm{Sp}\omega$ -submodule of  $N_1 \otimes \mathcal{A}_{k+1}^r(N_1)$ . To each elementary tensor  $u \otimes T \in N_1 \otimes \mathcal{A}_{k+1}^r(N_1)$  we associate the tree  $T_u \in \mathcal{A}_k(N_1)$  which is obtained by labelling the root of  $T$  by  $u$ . Extending by linearity we get the “labelling map”,  $Lab : N_1 \otimes \mathcal{A}_{k+1}^r(N_1) \longrightarrow \mathcal{A}_k(N_1)$ .

**Lemma 3.1.** (1) *For any  $k \geq 1$ , the composite*

$$\mathcal{A}_k(N_1) \xrightarrow{\text{expan.}} D_k(N_1) \xrightarrow{Lab} \mathcal{A}_k(N_1)$$

*is multiplication by  $k + 2$ .*

(2) *The Labelling map  $Lab : D_k(N_1) \longrightarrow \mathcal{A}_k(N_1)$  is injective.*

*Proof.* (1) A direct computation shows that the image of a tree  $T \in \mathcal{A}_{k+1}(N_1)$  in  $N_1 \otimes \mathcal{L}_{k+1}$  is  $\sum_{x \in V_1(T)} l_x \otimes T^x$  so that the composite  $\mathcal{A}_k(N_1) \longrightarrow D_k(N_1) \longrightarrow \mathcal{A}_k(N_1)$  sends  $T$  to  $(k + 2)T$ .

(2) Multiplication by  $k + 2$  certainly induces an isomorphism after tensorisation by the rationals. In [5] it is proved that the expansion map is also an isomorphism after tensorisation by the rationals. It follows that the map  $Lab \otimes \mathbf{Q}$  is an isomorphism. Consider the following commutative diagram :

$$\begin{array}{ccc}
D_k(N_1) & \xrightarrow{Lab} & \mathcal{A}_k(N_1) \\
\downarrow & & \downarrow \\
D_k(N_1) \otimes \mathbf{Q} & \xrightarrow{Lab \otimes \mathbf{Q}} & \mathcal{A}_k(N_1) \otimes \mathbf{Q}.
\end{array}$$

The group  $D_k(N_1)$  is free abelian as it is a subgroup of the free abelian group  $\text{Hom}(N_1, \mathcal{L}_{k+1})$ . In particular the left vertical arrow is an injection. The bottom arrow is an isomorphism. In particular it is injective. By commutativity of the diagram the composite  $D_k(N_1) \xrightarrow{Lab} \mathcal{A}_k(N_1) \rightarrow \mathcal{A}_k(N_1) \otimes \mathbf{Q}$  is therefore injective, and we may conclude that the labelling map is injective. Q. E. D.

#### 4. THE GROUP $\text{Can}(\mathcal{A}_2(N_1))$

We focus on the low degree Johnson homomorphisms. In the preceding sections we have constructed homomorphisms that fit into a commutative diagram :

$$\begin{array}{ccccc}
\mathcal{M}_{g,1}(3) & \hookrightarrow & \mathcal{M}_{g,1}(2) & \twoheadrightarrow & \mathcal{M}_{g,1}(2)/\mathcal{M}_{g,1}(3) \\
& & \searrow^{\tau_2} & & \downarrow \\
& & & & D_2(N_1) \hookrightarrow \mathcal{A}_2(N_1).
\end{array}$$

Moreover, all arrows are  $\mathcal{M}_{g,1}$ -equivariant and the action of  $\mathcal{M}_{g,1}$  on the three rightmost groups factors through  $\text{Sp}\omega$ .

From now on we denote the composite  $Lab \circ \tau_2$  by  $\tilde{\tau}_2$ . We denote by  $\mathcal{A}_{g,1}(k)$  (respectively  $\mathcal{B}_{g,1}(k)$ ) the subgroup  $\mathcal{A}_{g,1} \cap \mathcal{M}_{g,1}(k)$  (respectively  $\mathcal{B}_{g,1} \cap \mathcal{M}_{g,1}(k)$ ). By definition  $\mathcal{A}_{g,1}(k)$  (resp.  $\mathcal{B}_{g,1}(k)$ ) is normal in  $\mathcal{A}_{g,1}$  (resp.  $\mathcal{B}_{g,1}$ ).

**Definition 4.** Let  $\text{Can}(\mathcal{A}_2(N_1))$  denote the subgroup of  $\mathcal{A}_2(N_1)$  generated by  $\tilde{\tau}_2(\mathcal{A}_{g,1}(2))$  and  $\tilde{\tau}_2(\mathcal{B}_{g,1}(2))$ . It is the subgroup of cancelable trees.

The study of this group is justified by the next proposition.

**Proposition 4.1.** If  $\tilde{\tau}_2(\mathcal{M}_{g,1}(2)) \subset \text{Can}(\mathcal{A}_2(N_1))$ , then for any mapping class  $\phi \in \mathcal{M}_{g,1}(2)$  there exists a mapping class  $\psi \in \mathcal{M}_{g,1}(3)$  such that  $\phi \approx \psi$  (cf Definition 1), that is :  $S_\phi^3$  is diffeomorphic to  $S_\psi^3$ .

*Proof.* By definition of  $Can(\mathcal{A}_2(N_1))$  there exist elements  $\phi_a \in \mathcal{A}_{g,1}(2)$  and  $\phi_b \in \mathcal{B}_{g,1}(2)$  such that  $\tilde{\tau}_2(\phi) = \tilde{\tau}_2(\phi_a) + \tilde{\tau}_2(\phi_b)$ . Setting  $\psi = \phi_a^{-1} \circ \phi \circ \phi_b^{-1}$ , we have that  $\psi \in \mathcal{M}_{g,1}(3)$  and  $\psi \approx \phi$ . Q. E. D.

To fully exploit this proposition, we need first to have a description of a large class of elements in  $Can(\mathcal{A}_2(N_1))$  and second to have a description of the image of  $\tilde{\tau}_2 : \mathcal{M}_{g,1}(2) \rightarrow \mathcal{A}_2(N_1)$ .

#### 4.1. Some families of elements in $Can(\mathcal{A}_2(N_1))$ .

4.1.1. *Elements arising from bounding simple closed curves.* A bounding simple closed curve  $\gamma$ , BSCC for short, is a simple closed curve such that  $\Sigma_{g,1} - \{\gamma\}$  has two connected components. One of the components of  $\Sigma_{g,1} - \{\gamma\}$ , denoted by  $\Sigma_{h,\gamma}$ , is a surface with only one boundary component, which we may identify with  $\gamma$ . The genus of  $\gamma$  or the genus of the Dehn twist  $T_\gamma$  is the genus  $h$  of  $\Sigma_{h,\gamma}$ . This terminology was introduced by Johnson, who proved :

**Proposition 4.2.** [8] *The group  $\mathcal{M}_{g,1}(2)$  is generated by the Dehn twists around BSCC.*

There is a very nice, algebraic characterisation of the subgroups  $\mathcal{A}_{g,1}$  and  $\mathcal{B}_{g,1}$  due to Griffiths [4]. Choose a map  $\phi \in \mathcal{B}_{g,1}$ . Then  $\phi$  is the restriction of some diffeomorphism  $\Phi : \mathcal{H}_g \rightarrow \mathcal{H}_g$ , and we can assume that  $\Phi(x_0) = x_0$ , where  $x_0$  is the common base point of  $\Sigma_{g,1}$  and  $\mathcal{H}_g$ . The inclusion  $\Sigma_{g,1} \subset \mathcal{H}_g$  induces a surjective homomorphism  $\pi_1(\Sigma_{g,1}, x_0) \rightarrow \pi_1(\mathcal{H}_g, x_0)$ , its kernel is the normal subgroup generated by the set of curves  $\{\beta_i\}_{1 \leq i \leq g}$ . As  $\phi$  is the restriction of  $\Phi$ ,  $\pi_1(\phi)$  induces a morphism on  $\pi_1(\mathcal{H}_g, x_0)$ , in fact it induces precisely  $\pi_1(\Phi)$ . Therefore,  $\pi_1(\phi)$  preserves the kernel of  $\pi_1(\Sigma_{g,1}, x_0) \rightarrow \pi_1(\mathcal{H}_g, x_0)$ . It turns out that this last property characterises the group  $\mathcal{B}_{g,1}$  :

**Proposition 4.3.** [4] *An element in  $\mathcal{M}_{g,1}$  is in  $\mathcal{B}_{g,1}$  (resp.  $\mathcal{A}_{g,1}$ ), if and only if it preserves the normal subgroup of  $\pi_1(\Sigma_{g,1}, x_0)$  generated by the set  $\{\beta_i\}_{1 \leq i \leq g}$  (resp.  $\{\alpha_i\}_{1 \leq i \leq g}$ ), which is the kernel of the canonical morphism  $\pi_1(\Sigma_{g,1}, x_0) \rightarrow \pi_1(\mathcal{H}_g, x_0)$  (resp.  $\pi_1(\Sigma_{g,1}, x_0) \rightarrow \pi_1(-\mathcal{H}_g, x_0)$ ).*

We are now ready to describe our first family :

**Proposition 4.4.** *Let  $\gamma$  be a simple closed curve on  $\Sigma_{g,1}$  which is contractible in  $\mathcal{H}_g$ , then  $T_\gamma$  belongs to  $\mathcal{B}_{g,1}$ . In particular if  $\gamma$  is a BSCC, then  $\tilde{\tau}_2(T_\gamma)$  belongs to  $\text{Can}(\mathcal{A}_2(N_1))$ . Similarly, if  $\gamma \in \Sigma_{g,1}$  is contractible in  $-\mathcal{H}_g$ , then  $T_\gamma \in \mathcal{A}_{g,1}$ , and if  $\gamma$  is moreover a BSCC, then  $\tilde{\tau}_2(T_\gamma) \in \text{Can}(\mathcal{A}_2(N_1))$ .*

*Proof.* By Griffiths' result, Proposition 4.3, it is enough to prove that  $\forall 1 \leq i \leq g$ ,  $T_\gamma(\beta_i) \in \ker(\pi_1(\Sigma_{g,1}, x_0) \rightarrow \pi_1(\mathcal{H}_g, x_0))$ . As  $\beta_i$  belongs to the kernel, it is enough to prove that  $T_\gamma(\beta_i)$  is homotopic to  $\beta_i$  in  $\mathcal{H}_g$ . Up to homotopy the curves  $\gamma$  and  $\beta_i$  meet transversally in a finite number of points  $x_1, \dots, x_s$ . For each  $1 \leq j \leq s$  consider  $\gamma$  as a based loop at  $x_j$ , and choose a based homotopy  $H_j : S^1 \times [0, 1] \rightarrow \mathcal{H}_g$  between  $\gamma$  and the constant loop at  $x_j$ . This is possible since  $\gamma$  is contractible in  $\mathcal{H}_g$  by assumption. These homotopies reassemble into a homotopy between  $T_\gamma(\beta_i)$  and  $\beta_i$ .

If  $\gamma$  is a BSCC, then according to Johnson ,  $T_\gamma$  belongs also to  $\mathcal{M}_{g,1}(2)$ .

Q. E. D.

**Remark 4.1.** *There is an easy way to produce a BSCC of genus 1 in  $\Sigma_{g,1}$ . Consider two simple closed curves  $\lambda$  and  $\mu$  intersecting exactly in one point on  $\Sigma_{g,1}$ . A tubular neighbourhood of the union  $\lambda \cup \mu$  is a subsurface of genus 1 in  $\Sigma_{g,1}$ , its boundary  $\delta$  is the desired BSCC of genus 1. If  $[\lambda], [\mu] \in N_1$  are the homology classes of the curves  $\lambda$  and  $\mu$ , then the image of  $T_\delta$  in  $\mathcal{A}_2(N_1)$  is the tree*

$$\begin{array}{cc} [\lambda] & [\mu] \\ 2 & \text{---} \\ [\mu] & [\lambda] \end{array} .$$


*The last assertion follows from a direct computation in the case  $\lambda = \alpha_1$  and  $\mu = \beta_1$  together with the fact that  $\mathcal{M}_{g,1}$  acts transitively on the pairs of simple closed curves intersecting exactly in one point.*

In the sequel, when producing a BSCC of genus 1, we will only draw the two curves  $\lambda$  and  $\mu$ , and label them by their homology class. The curves  $\{\lambda, \mu\}$

are usually called the spine of the sub-surface, as we focus on the boundary we call  $\{\lambda, \mu\}$  the spine of the boundary.

4.1.2. *Elements arising from commutators.* Our second family is more algebraic in nature, it comes from commutators of pairs of elements both sitting in  $\mathcal{A}_{g,1}(1)$  or in  $\mathcal{B}_{g,1}(1)$ .

Recall that we have an isomorphism  $\mathcal{A}_1(N_1) \simeq \Lambda^3 N_1$  induced by the expansion map. According to S. Morita, the images of both  $\tau_1(\mathcal{B}_{g,1}(1))$  and  $\tau_1(\mathcal{A}_{g,1}(1))$  have a very simple characterisation in  $\Lambda^3 N_1 \simeq \mathcal{A}_1(N_1)$ .

**Proposition 4.5.** [11] A tree  represents an element of  $\tau_1(\mathcal{A}_{g,1}(1))$  (resp.  $\mathcal{B}_{g,1}(1)$ ) if and only if at least one of its labels belongs to the Lagrangian  $L_A$  (resp.  $L_B$ ).

**Corollary 4.1.** The group  $\tilde{\tau}_2([\mathcal{A}_{g,1}(1), \mathcal{A}_{g,1}(1)])$  (resp.  $\tilde{\tau}_2([\mathcal{B}_{g,1}(1), \mathcal{B}_{g,1}(1)])$ ) is generated by the brackets

$$4 \left[ \begin{array}{c} c \\ \diagup \quad \diagdown \\ d \quad e \end{array}, \begin{array}{c} f \\ \diagup \quad \diagdown \\ g \quad h \end{array} \right],$$

where at least one of the labels  $\{c, d, e\}$  and one of the labels  $\{f, g, h\}$  are in the Lagrangian  $L_A$  (resp.  $L_B$ ).

*Proof.* As the Johnson homomorphism defines a Lie algebra homomorphism we have  $\tau_2([\mathcal{B}_{g,1}(1), \mathcal{B}_{g,1}(1)]) = [\tau_1(\mathcal{B}_{g,1}(1)), \tau_1(\mathcal{B}_{g,1}(1))]$ , whence the condition on the labels. This subgroup is obviously contained in  $\mathcal{A}_2(N_1) \subset D_2(N_1)$  and as the composite  $\mathcal{A}_2(N_1) \hookrightarrow D_2(N_1) \rightarrow A_2(N_1)$  is multiplication by 4 we get the factor 4 in front of the bracket. Q. E. D.

4.1.3. *Invariance properties of  $Can(\mathcal{A}_2(N_1))$ .* From the above two families of cancelable trees it is easy to construct more cancelable trees by making use of the elements in  $\mathcal{M}_{g,1}$  that preserve the group  $Can(\mathcal{A}_2(N_1))$ .

In Lemma 2.1 we introduced a mapping class  $\iota_g \in \mathcal{M}_{g,1}$  such that  $S^3 = \mathcal{H}_g \coprod_{\iota_g} -\mathcal{H}_g$  and we remarked that  $\mathcal{A}_{g,1} = \iota_g^{-1}\mathcal{B}_{g,1}\iota_g$ . By equivariance of the map  $\tilde{\tau}_2$ , the group  $Can(\mathcal{A}_2(N_1))$  is invariant under the action of  $\iota_g$ .

In the same spirit, if  $\phi \in \mathcal{A}_{g,1}(2)$  and  $\Phi \in \mathcal{A}_{g,1}$  then  $\Phi \cdot \tilde{\tau}_2(\phi) = \tilde{\tau}_2(\Phi\phi\Phi^{-1}) \in Can(\mathcal{A}_2(N_1))$ , because  $\mathcal{A}_{g,1}(2)$  is normal in  $\mathcal{A}_{g,1}$ . The same result holds for  $\mathcal{B}_{g,1}(2)$ .

If  $T \in Can(\mathcal{A}_2(N_1))$  and  $\Phi \in \mathcal{A}_{g,1} \cap \mathcal{B}_{g,1}$  then  $\Phi \cdot T \in Can(\mathcal{A}_2(N_1))$ . Indeed, by definition of  $Can(\mathcal{A}_2(N_1))$ , we can decompose  $T$  into a sum  $T = T_a + T_b$ , where  $T_a \in \tilde{\tau}_2(\mathcal{A}_{g,1}(2))$  and  $T_b \in \tilde{\tau}_2(\mathcal{B}_{g,1}(2))$  and then apply the preceding remarks to  $T_a$  and  $T_b$ .

Recall that the action of  $\mathcal{M}_{g,1}$  on  $\mathcal{A}_2(N_1)$  factors through  $Sp\omega$ , so to fully exploit the preceding theorem we need to know the images of  $\mathcal{A}_{g,1}$ ,  $\mathcal{B}_{g,1}$  and  $\iota_g$  in  $Sp\omega$ .

The action of  $\iota_g$  is easy to deduce from its action in homology which is given by :  $\iota(a_i) = -b_i$  and  $\iota(b_i) = a_i$ , for all  $1 \leq i \leq g$ .

In [20] Suzuki describes generators for the group  $\mathcal{B}_{g,1}$  and therefore for  $\mathcal{A}_{g,1} = \iota_g\mathcal{B}_{g,1}\iota_g^{-1}$ , their action on the basis  $\{a_i, b_i\}_{1 \leq i \leq g}$  of  $N_1$  is summarised in the following table (if an element of the basis is omitted, it is fixed under the action). The names of the generators are those given by Suzuki.

Common generators to $\mathcal{A}_{g,1}$ and $\mathcal{B}_{g,1}$	Action on the basis of $N_1$
Twist of the knob $i$	$a_i \rightarrow -a_i$ $b_i \rightarrow -b_i$
Exchange of knobs $i$ and $j$	$a_i \leftrightarrow a_j$ $b_i \leftrightarrow b_j$
Generators of $\mathcal{A}_{g,1}$ alone	Action on the basis of $N_1$
A-twist of the $i$ handle	$b_i \rightarrow b_i - a_i$
Sliding $\theta_{A,i,j}$	$a_i \rightarrow a_i + a_j$ $b_j \rightarrow b_j - b_i$
Sliding $\xi_{A,i,j}$	$b_i \rightarrow b_i - a_j$ $b_j \rightarrow b_j - a_i$
Generators of $\mathcal{B}_{g,1}$ alone	Action on the basis of $N_1$
B-twist of the $i$ handle	$a_i \rightarrow a_i - b_i$
Sliding $\theta_{B,i,j}$	$a_i \rightarrow a_i - a_j$ $b_j \rightarrow b_j + b_i$
Sliding $\xi_{B,i,j}$	$a_i \rightarrow a_i - b_j$ $a_j \rightarrow a_j - b_i$

It is a standard fact that the subgroup of the elements of  $\mathrm{Sp}\omega$  that respect the orthogonal decomposition  $N_1 = L_A \oplus L_B$  is naturally isomorphic to  $\mathrm{GL}_g(\mathbf{Z})$ . In particular, if  $P$  denotes the canonical projection  $P : \mathcal{M}_{g,1} \rightarrow \mathrm{Sp}\omega$ , we have  $P(\mathcal{A}_{g,1}) \cap P(\mathcal{B}_{g,1}) \subset \mathrm{GL}_g(\mathbf{Z})$ . The ‘‘sliding’’ and the ‘‘twists of the knobs’’ morphisms hit the standard generators of  $\mathrm{GL}_g(\mathbf{Z})$ , therefore :

**Lemma 4.1.** *In  $\mathrm{Sp}\omega$  we have  $\mathrm{GL}_g(\mathbf{Z}) = P(\mathcal{A}_{g,1}) \cap P(\mathcal{B}_{g,1})$ .*

Summarising, we have :

**Theorem 4.1.** *The group  $\mathrm{Can}(\mathcal{A}_2(N_1))$  is invariant :*

- (1) *Under the action of  $\iota_g$ .*
- (2) *Under the action of  $\mathrm{GL}_g(\mathbf{Z}) = P(\mathcal{A}_{g,1}) \cap P(\mathcal{B}_{g,1})$ .*
- (3) *If  $T \in \mathrm{Can}(\mathcal{A}_2(N_1))$  is contained in  $\tilde{\tau}_2(\mathcal{A}_{g,1}(2))$ , then for all  $\Phi \in \mathcal{A}_{g,1}$   $\Phi \cdot T \in \mathrm{Can}(\mathcal{A}_2(N_1))$ .*

- (4) If  $T \in \text{Can}(\mathcal{A}_2(N_1))$  is contained in  $\tilde{\tau}_2(\mathcal{B}_{g,1}(2))$ , then for all  $\Phi \in \mathcal{B}_{g,1}$   
 $\Phi \cdot T \in \text{Can}(\mathcal{A}_2(N_1))$ .

4.2. **The image of  $\tilde{\tau}_2 : \mathcal{M}_{g,1}(2) \rightarrow \mathcal{A}_2(N_1)$ .** Here is a stronger version of Proposition 4.2, also due to Johnson.

**Theorem 4.2.** [8] *For  $g \geq 3$ , the group  $\mathcal{M}_{g,1}(2)$  is generated by the Dehn twists around BSCC of genus 1 and 2.*

The mapping class group acts naturally on the set of simple closed curves, two BSCC are in the same orbit if and only if they have the same genus. It follows easily from this that two Dehn twists around BSCC are conjugated in  $\mathcal{M}_{g,1}$  if and only if they have the same genus. Therefore  $\mathcal{M}_{g,1}(2)$  is generated by the conjugates of any two Dehn twists  $T_{\gamma_1}$  and  $T_{\gamma_2}$  of genus 1 and 2 respectively and  $\tilde{\tau}_2(\mathcal{M}_{g,1}(2))$  is generated by the orbits of  $\tilde{\tau}_2(T_{\gamma_1})$  and  $\tilde{\tau}_2(T_{\gamma_2})$  under the action of  $\mathcal{M}_{g,1}$ . The algebraic version, in terms of trees, of these results is given by :

**Theorem 4.3.** *The image of  $\tilde{\tau}_2$  in  $\mathcal{A}_2(N_1)$  is the  $\mathcal{M}_{g,1}$ -module generated by the elements of the form*

$$2 \begin{array}{c} u \quad v \\ | \text{---} | \\ v \quad u \end{array} \text{ with } \omega(u \wedge v) = 1$$

$$4 \begin{array}{c} u_1 \quad v_2 \\ | \text{---} | \\ v_1 \quad u_2 \end{array} \text{ with } \omega(u_i \wedge v_j) = \delta_{ij} \text{ and } \omega(u_1 \wedge u_2) = 0 = \omega(v_1 \wedge v_2).$$

*Proof.* We choose as curves  $\gamma_1$  and  $\gamma_2$  the curves of genus 1 and 2 parallel to the boundary as in Figure 5.

A direct computation shows that  $T_{\gamma_1}$  acts on  $\alpha_1, \beta_1 \in \pi_1(\Sigma_{g,1})$  as the conjugation by  $[\alpha_1, \beta_1]$ , and leaves the other generators  $\alpha_i, \beta_i$   $2 \leq i \leq g$  unchanged.




FIGURE 5. Curves  $\gamma_1$  and  $\gamma_2$

By definition of our maps,

$$\tau_2(T_{\gamma_1}) = a_1 \otimes [b_1, [a_1, b_1]] + b_1 \otimes [a_1, [b_1, a_1]],$$

and

$$\tilde{\tau}_2(T_{\gamma_1}) = 2 \begin{array}{c} a_1 \quad b_1 \\ | \text{---} | \\ b_1 \quad a_1 \end{array} .$$

The orbit of this last element is clearly the set of trees of the form  $2 \begin{array}{c} u \quad v \\ | \text{---} | \\ v \quad u \end{array}$

such that  $\omega(u \wedge v) = 1$ .

Similarly, writing down the action of  $T_{\gamma_2}$  on curves  $\alpha_1, \beta_1, \alpha_2, \beta_2$  one gets :

$$\begin{aligned} \tau_2(T_{\gamma_2}) &= b_1 \otimes [a_1, [b_1, a_1]] + b_1 \otimes [a_1, [b_2, a_2]] + \\ & a_1 \otimes [b_1, [a_1, b_1]] + a_1 \otimes [b_1, [a_2, b_2]] + \\ & b_2 \otimes [a_2, [b_1, a_1]] + b_2 \otimes [a_2, [b_2, a_2]] + \\ & a_2 \otimes [b_2, [a_1, b_1]] + a_2 \otimes [b_2, [a_2, b_2]], \end{aligned}$$

and :

$$\tilde{\tau}_2(T_{\gamma_2}) = 2 \begin{array}{c} a_1 \quad b_1 \\ | \text{---} | \\ b_1 \quad a_1 \end{array} + 4 \begin{array}{c} a_1 \quad b_2 \\ | \text{---} | \\ b_1 \quad a_2 \end{array} + 2 \begin{array}{c} a_2 \quad b_2 \\ | \text{---} | \\ b_2 \quad a_2 \end{array} .$$

The orbit of this element gives, up to elements coming from the orbit of  $\tilde{\tau}_2(T_{\gamma_1})$ ,

all trees of the form  $4 \begin{array}{cc} u_1 & v_2 \\ | & | \\ \hline & \\ | & | \\ v_1 & u_2 \end{array}$  where  $\omega(u_1 \wedge v_1) = 1 = \omega(u_2 \wedge v_2)$  and

$$\omega(u_1 \wedge u_2) = \omega(u_1 \wedge v_2) = 0 = \omega(u_2 \wedge v_1). \quad \text{Q. E. D.}$$

## 5. PROOF OF THE MAIN THEOREM

**Main Theorem.** *Any integral homology sphere is diffeomorphic to  $S_\phi^3$  for some  $\phi \in \mathcal{M}_{g,1}(3)$  with  $g \geq 9$ .*

By Morita's result, we have a bijection (cf. Theorem 2.3) :

$$\lim_{g \rightarrow +\infty} \mathcal{M}_{g,1}(2)/\approx \longrightarrow \mathcal{S}(3).$$

Thus it suffices to show that the inclusions  $\mathcal{M}_{g,1}(3) \hookrightarrow \mathcal{M}_{g,1}(2)$  induce a bijection :

$$\lim_{g \rightarrow +\infty} \mathcal{M}_{g,1}(3)/\approx \longrightarrow \lim_{g \rightarrow +\infty} \mathcal{M}_{g,1}(2)/\approx .$$

This is the case if for large enough values of  $g$  any element in  $\mathcal{M}_{g,1}(2)$  is equivalent to some element in  $\mathcal{M}_{g,1}(3)$ . By Proposition 4.1 this will follow from :

**Theorem 5.1.** *For  $g \geq 9$ , the image of  $\tilde{\tau}_2 : \mathcal{M}_{g,1}(2) \rightarrow \mathcal{A}_2(N_1)$  belongs to the subgroup  $\text{Can}(\mathcal{A}_2(N_1))$ .*

**5.1. Three technical lemmas.** In the sequel we will work “modulo the subgroup  $\text{Can}(\mathcal{A}_2(N_1))$ ”. Two elements, say  $S, T$  in  $\mathcal{A}_2(N_1)$ , which differ by an element of  $\text{Can}(\mathcal{A}_2(N_1))$  will be called *equivalent*, denoted by  $S \approx T$ . An element equivalent to 0 will be called *cancelable*. It is obvious that being equivalent is invariant under the action of  $\text{GL}_g(\mathbf{Z})$ . We also assume that  $g \geq 9$ .

**Lemma 5.1.** *For all  $i \neq j$  the following trees are in  $\tilde{\tau}_2(\mathcal{A}_{g,1}(2)) \cap \tilde{\tau}_2(\mathcal{B}_{g,1}(2)) \subset \text{Can}(\mathcal{A}_2(N_1))$ .* :

$$2 \begin{array}{cc} a_i & b_i \\ | & | \\ \hline & \\ | & | \\ b_i & a_i \end{array} , \quad 4 \begin{array}{cc} a_i & b_j \\ | & | \\ \hline & \\ | & | \\ b_i & a_j \end{array} .$$

*Proof.* The trees  $2 \begin{array}{c} a_i \quad b_i \\ | \text{---} | \\ b_i \quad a_i \end{array}$  are all in the  $\mathrm{GL}_g(\mathbf{Z})$ -orbit of  $2 \begin{array}{c} a_1 \quad b_1 \\ | \text{---} | \\ b_1 \quad a_1 \end{array}$ ,

is cancelable since the curve  $\gamma_1$  in Figure 5 is contractible in both  $\mathcal{H}_g$  and  $-\mathcal{H}_g$ . By Proposition 4.4 they are all cancelable. The same argument holds for

$4 \begin{array}{c} a_i \quad a_i \\ | \text{---} | \\ b_i \quad a_j \end{array}$  since, up to trees of the preceding form, it belongs to the  $\mathrm{GL}_g(\mathbf{Z})$ -orbit of the curve  $\gamma_2$ , which is contractible too.

Q. E. D.

**Column Intersection Lemma 5.2.** *Suppose that the tree  $4 \begin{array}{c} c \quad f \\ | \text{---} | \\ d \quad e \end{array}$  is labelled by elements of the symplectic basis  $\{a_i, b_i\}_{1 \leq i \leq g}$ . Assume furthermore*

*that  $\omega(c \wedge e) = \omega(c \wedge f) = 0 = \omega(d \wedge e) = \omega(d \wedge f)$ . Then  $4 \begin{array}{c} c \quad f \\ | \text{---} | \\ d \quad e \end{array}$  is*

*cancelable.*

*Proof.* As  $g \geq 9$ , there is some index  $j$  such that the sets  $\{a_j, b_j\}$  and  $\{c, d, e, f\}$  are disjoint. Assume  $c \in \{b_i\}_{1 \leq i \leq g}$ . According to Proposition 4.5 both elements

$\begin{array}{c} a_j \\ / \quad \backslash \\ c \quad d \end{array}$  and  $\begin{array}{c} b_j \\ / \quad \backslash \\ e \quad f \end{array}$  represent elements of  $\tau_1(\mathcal{B}_{g,1}(1))$ . By definition of the bracket :

$$4 \left[ \begin{array}{c} a_j \\ / \quad \backslash \\ c \quad d \end{array}, \begin{array}{c} b_j \\ / \quad \backslash \\ e \quad f \end{array} \right] = 4 \begin{array}{c} c \quad f \\ | \text{---} | \\ d \quad e \end{array},$$

which is cancelable by Corollary 4.1.

If  $c \in \{a_i\}_{1 \leq i \leq g}$ , just replace the pair  $(a_j, b_j)$  in the last equality by  $(-b_j, a_j)$ .

Q. E. D.

**Chain Equivalences Lemma 5.3.** *For all pairwise distinct indices  $i, j, k$ , we have three chain of equivalences :*

$$1) \ 2 \begin{array}{c} a_i \quad b_j \\ | \text{---} | \\ b_j \quad a_i \end{array} \approx 4 \begin{array}{c} a_i \quad b_j \\ | \text{---} | \\ b_j \quad a_j \end{array} \approx 4 \begin{array}{c} a_k \quad b_k \\ | \text{---} | \\ a_i \quad b_j \end{array} \approx 4 \begin{array}{c} a_k \quad b_k \\ | \text{---} | \\ b_j \quad a_i \end{array} ;$$

$$\begin{aligned}
 2) \quad & 2 \begin{array}{c} a_i \quad a_j \\ \text{---} \\ a_j \quad a_i \end{array} \approx 4 \begin{array}{c} b_j \quad a_j \\ \text{---} \\ a_j \quad a_i \end{array} \approx 4 \begin{array}{c} a_k \quad b_k \\ \text{---} \\ a_i \quad a_j \end{array} \approx 4 \begin{array}{c} a_k \quad b_k \\ \text{---} \\ a_j \quad a_i \end{array} ; \\
 3) \quad & 2 \begin{array}{c} b_i \quad b_j \\ \text{---} \\ b_j \quad b_i \end{array} \approx 4 \begin{array}{c} a_j \quad b_j \\ \text{---} \\ b_j \quad b_i \end{array} \approx 4 \begin{array}{c} b_k \quad a_k \\ \text{---} \\ b_i \quad b_j \end{array} \approx 4 \begin{array}{c} b_k \quad a_k \\ \text{---} \\ b_j \quad b_i \end{array} .
 \end{aligned}$$

Moreover, any of this elements is equivalent to its opposite, so that 2 times any of this elements is cancelable.

*Proof.* Using suitable exchange of knobs we may suppose that  $i = 1, j = 2$  and  $k = 3$ .

**First chain of equivalences**

Observe that the curve  $\delta$ , which spine is drawn in Figure 6, is contractible in  $\mathcal{H}_g$ , in particular  $\tilde{\tau}_2(T_\delta)$  is cancelable.




FIGURE 6. Spine of the curve  $\delta$

Now,

$$\tilde{\tau}_2(T_\delta) = 2 \begin{array}{c} a_1 + a_2 \quad b_2 \\ \text{---} \\ b_2 \quad a_1 + a_2 \end{array} = 2 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_1 \end{array} + 2 \begin{array}{c} a_2 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} + 4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} .$$

As  $2 \begin{array}{c} a_2 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array}$  is cancelable, we get  $2 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_1 \end{array} \approx -4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array}$ .

Applying the “twist of the knob 2” to both sides removes the minus sign.

The equality

$$4 \left[ \begin{array}{c} a_3 \\ \diagup \quad \diagdown \\ a_1 \quad b_2 \end{array}, \begin{array}{c} b_3 \\ \diagup \quad \diagdown \\ a_2 \quad b_2 \end{array} \right] = 4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} - 4 \begin{array}{c} a_3 \quad b_3 \\ \text{---} \\ a_1 \quad b_2 \end{array},$$

shows that  $4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} \approx 4 \begin{array}{c} a_3 \quad b_3 \\ \text{---} \\ a_1 \quad b_2 \end{array}$ . Exchanging the roles of  $a_3$  and  $b_3$  in

the previous bracket we get  $4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} \approx -4 \begin{array}{c} b_3 \quad a_3 \\ \text{---} \\ a_1 \quad b_2 \end{array}$ .

Notice that we proved that  $4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} \approx 2 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_1 \end{array} \approx -4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array}$ . In

particular  $4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_1 \end{array} \approx 4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} - 4 \begin{array}{c} a_1 \quad b_2 \\ \text{---} \\ b_2 \quad a_2 \end{array} = 0$ . This proves that any

tree in this chain is equivalent to its opposite, and therefore that two times any tree involved in this Chain of Equivalences is cancelable. A similar argument works for the other two Chains of Equivalences

**Second and third chains of equivalences**

The second chain of equivalences is proved in the same way as the first one. One has just to replace the curve  $\delta$  by the curve  $\delta'$  which spine is drawn in Figure 7 and which is contractible in  $-\mathcal{H}_g$ . The third chain of equivalences is




FIGURE 7. Spine of the curve  $\delta'$

obtained from the second one by applying  $\iota_g$  termwise.

Q. E. D.

**5.2. Proof of Theorem 5.1.** We are now in position to prove that the image of  $\tilde{\tau}_2$  belongs to  $Can(\mathcal{A}_2(N_1))$ . It is obviously enough to prove that the image of the generators of  $\mathcal{M}_{g,1}(2)$ , that is Dehn twists of genus 1 and 2 belong to  $Can(\mathcal{A}_2(N_1))$ .

The proof of Theorem 5.1 is quite straightforward once we have in hand the three above technical Lemmas 5.1 5.2 and 5.3 but nevertheless a bit long and proceeds by "case by case" computations. Therefore we will only write the main lines of the proof leaving the details of the computations to the interested reader.

**5.2.1. Images of genus 1 twists are in  $Can(\mathcal{A}_2(N_1))$ .** By the description of the images of genus 1 BSCC (see Theorem 4.3) it is enough to prove

**Theorem 5.2.** *Assume  $g \geq 9$ . For any pair of elements  $u, w \in N_1$  such that  $\omega(u \wedge w) = 1$ ,*

$$\begin{array}{ccc} u & & w \\ 2 & \text{---} & \\ w & & u \end{array}$$

*is cancelable.*

If  $x, y \in N_1$  we denote by  $\langle x, y \rangle \subset N_1$  the submodule they generate.

*Proof. First step :* Without loss of generality we may suppose that  $u, w \in \langle a_1, a_2, b_1, b_2 \rangle$ .

Recall that  $N_1 = L_A \oplus L_B$  and decompose  $u$  and  $w$  as  $u = a_u + b_u$  and  $w = a_w + b_w$ . The elements  $a_u$  and  $a_w$  are contained in a direct factor of  $A$  of rank at most 2, therefore there exists  $\phi \in GL_g(\mathbf{Z})$  such that  $\phi(a_u)$  and  $\phi(a_w)$  belong to  $\langle a_1, a_2 \rangle$ . Similarly  $\phi(b_u)$  and  $\phi(b_w)$  are the sum of elements contained in  $\langle b_1, b_2 \rangle$  and of two elements  $\hat{b}_u$  and  $\hat{b}_w$  contained in  $\langle b_3, \dots, b_g \rangle$ . As  $\hat{b}_u$  and  $\hat{b}_w$  are contained in a direct factor of  $\langle b_3, \dots, b_g \rangle$  of rank at most two, there exists  $\psi \in GL_g(\mathbf{Z})$  such that  $\psi|_{\langle b_1, b_2 \rangle} = Id = {}^t \psi^{-1}|_{\langle a_1, a_2 \rangle}$  and  $\psi(\langle \hat{b}_u, \hat{b}_w \rangle) \subset \langle b_3, b_4 \rangle$ .

By construction the labels of the trees

$${}^t\psi\phi \cdot \left( 2 \begin{array}{c} u \quad w \\ | \text{---} | \\ w \quad u \end{array} \right)$$

are contained in the desired submodule. As a consequence of Theorem 4.1,

$$2 \begin{array}{c} u \quad w \\ | \text{---} | \\ w \quad u \end{array} \text{ is cancelable if and only if } {}^t\psi\phi(2 \begin{array}{c} u \quad w \\ | \text{---} | \\ w \quad u \end{array}) \text{ is.}$$

As an intermediary step, we may thus assume that  $u, w \in \langle a_1, a_2, b_1, b_2, b_3, b_4 \rangle$ .

By symmetry, to reduce the module to  $\langle a_1, a_2, b_1, b_2 \rangle$  it is enough to prove that if  $w = w' + \hat{b}_w$  with  $w' \in \langle a_1, a_2, b_1, b_2 \rangle$  and  $\hat{b}_w \in \langle b_3, b_4 \rangle$  then

$$2 \begin{array}{c} u \quad w \\ | \text{---} | \\ w \quad u \end{array} \approx 2 \begin{array}{c} u \quad w' \\ | \text{---} | \\ w' \quad u \end{array} .$$

By multilinearity of the labels :

$$2 \begin{array}{c} u \quad w \\ | \text{---} | \\ w \quad u \end{array} - 2 \begin{array}{c} u \quad w' \\ | \text{---} | \\ w' \quad u \end{array} = 2 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad \hat{b}_w \end{array} + 4 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad w' \end{array} .$$

Computing the cancellable bracket

$$4 \left[ \begin{array}{c} a_5 \quad b_5 \\ \diagdown \quad \diagup \\ \hat{b}_w \quad u \end{array} , \begin{array}{c} b_5 \\ \diagdown \quad \diagup \\ w' \quad u \end{array} \right] = 4 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad w' \end{array} + 4 \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad u \end{array}$$

we see that

$$2 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad \hat{b}_w \end{array} + 4 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad w' \end{array} \approx 2 \begin{array}{c} \hat{b}_w \quad u \\ | \text{---} | \\ u \quad \hat{b}_w \end{array} - 4 \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad u \end{array} .$$

We have to prove that the righthand term is cancelable.

Write  $u = n_1 a_1 + n_2 a_2 + \sum_{i=1}^4 m_i b_i$  and expand  $2 \begin{array}{c} \hat{b}_w \quad u \quad a_5 \quad b_5 \\ | \text{---} | \\ u \quad \hat{b}_w \quad \hat{b}_w \quad u \end{array} - 4 \begin{array}{c} | \text{---} | \\ | \text{---} | \end{array}$

by multilinearity. It is equal to

$$\begin{aligned} & \sum_{i=1}^2 (2n_i^2 \begin{array}{c} \hat{b}_w \quad a_i \\ | \text{---} | \\ a_i \quad \hat{b}_w \end{array} - 4n_i \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad a_i \end{array}) + \sum_{i=1}^4 (2m_i^2 \begin{array}{c} \hat{b}_w \quad b_i \\ | \text{---} | \\ b_i \quad \hat{b}_w \end{array} - 4m_i \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad b_i \end{array}) \\ & + \sum_{1 \leq i \neq j \leq 4} 4n_i m_j \begin{array}{c} \hat{b}_w \quad b_j \\ | \text{---} | \\ a_i \quad \hat{b}_w \end{array}. \end{aligned}$$

By the Column Intersection Lemma 5.2, each term in the third sum is cancelable. The terms in two remaining sums are handled in the same way. We only write the details for terms in the first sum.

**Lemma 5.4.** *For  $1 \leq i \leq 2$ , the sum,*

$$2n_i^2 \begin{array}{c} \hat{b}_w \quad a_i \\ | \text{---} | \\ a_i \quad \hat{b}_w \end{array} - 4n_i \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad a_i \end{array}.$$

*is cancelable.*

*Proof.* Write  $\hat{b}_w = s b_3 + t b_4$ . Expanding by multilinearity one finds that

$$2n_i^2 \begin{array}{c} \hat{b}_w \quad a_i \\ | \text{---} | \\ a_i \quad \hat{b}_w \end{array} - 4n_i \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ \hat{b}_w \quad a_i \end{array} \text{ is equivalent to :}$$

$$4((n_i s)^2 - n_i s) \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ b_3 \quad a_i \end{array} + 4((n_i t)^2 - n_i t) \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ b_4 \quad a_i \end{array}.$$

We have proved in the Chain Equivalences Lemma 5.3 that the 2 times any

term that appears in a ‘‘Chain Equivalence’’ is cancelable. As  $4 \begin{array}{c} a_5 \quad b_5 \\ | \text{---} | \\ b_3 \quad a_i \end{array}$

$a_5$   $b_5$   
 and  $4 \begin{array}{c} | \\ \hline | \end{array}$  both appear in the first chain and both  $((n_i s)^2 - n_i s)$  and  $b_4$   $a_i$   
 $((n_i t)^2 - n_i t)$  are even, the sum is cancelable. Q. E. D.

**Second step** Assume that  $u, w \in \langle a_1, a_2, b_1, b_2 \rangle$ . Set  $u = u_1^a a_1 + u_2^a a_2 + u_1^b b_1 + u_2^b b_2$  and  $w = w_1^a a_1 + w_2^a a_2 + w_1^b b_1 + w_2^b b_2$ .

$u$   $w$   
 We expand the tree  $2 \begin{array}{c} | \\ \hline | \end{array}$  by multilinearity and group the trees by  $w$   $u$   
 type, according to the number of  $a_i$ 's and  $b_i$ 's that appear as labels. The types are thus (4a), (3a, 1b) (2a, 2b), (1a, 3b) and (4b). We will inspect these trees by type and show that after grouping equivalent trees, they are all cancelable.

$x_i$   $t_i$   
 The coefficient of the tree  $\begin{array}{c} | \\ \hline | \end{array}$  is  $(u_i^x w_j^y - u_j^y w_i^x)(u_k^z w_l^t - u_l^t w_k^z)$  up to  $y_j$   $z_k$   
 a coefficient 2 or 4 depending on the symmetry of the tree.

#### Types (4a) and (3a,1b)

Tree	$2 \begin{array}{c} a_1 \quad a_2 \\   \\ \hline   \\ a_2 \quad a_1 \end{array}$	$4 \begin{array}{c} a_1 \quad b_1 \\   \\ \hline   \\ a_2 \quad a_1 \end{array}$	$4 \begin{array}{c} a_1 \quad b_2 \\   \\ \hline   \\ a_2 \quad a_2 \end{array}$
Coefficient	$(u_1^a w_2^a - u_2^a w_1^a) \times (u_1^a w_2^a - u_2^a w_1^a)$	$(u_1^a w_2^a - u_2^a w_1^a) \times (u_1^a w_1^b - u_1^b w_1^a)$	$(u_1^a w_2^a - u_2^a w_1^a) \times (u_2^a w_2^b - u_2^b w_2^a)$

Tree	$4 \begin{array}{c} a_1 \quad b_1 \\   \\ \hline   \\ a_2 \quad a_2 \end{array}$	$4 \begin{array}{c} a_2 \quad b_2 \\   \\ \hline   \\ a_1 \quad a_1 \end{array}$
Coefficient	Not needed	not needed

By the second chain of equivalences the first 3 trees are mutually equivalent. As in Lemma 5.4, to prove that their sum is cancelable it is enough to prove that the sum of their coefficients is even. A direct computation using the fact that  $\omega(u \wedge v) = 1$  yields the result.

The last two trees are in the same  $Gl_g(\mathbf{Z})$ -orbit, therefore one is cancelable

if and only if the other is. The proof of the cancellation of the tree  $4 \begin{array}{c} a_1 \quad b_1 \\ | \text{---} | \\ a_2 \quad a_2 \end{array}$

is obtained by applying the Chain Equivalences Lemma 5.3 to the trees that appear from the spine of the curve  $\delta''$  on Figure 8.




FIGURE 8. Spine of  $\delta''$

**Type (4b) and (3a,1b)**

These trees are obtained from the preceding list by interchanging the roles of  $a$  and  $b$ . The same argument as above shows that their sum is cancelable.

**Type (2a,2b)**

There are 11 trees of this type to consider. Among them, 3 are cancelable by Lemma 5.1,

$$2 \begin{array}{c} a_1 \quad b_1 \\ | \text{---} | \\ b_1 \quad a_1 \end{array} , 2 \begin{array}{c} a_2 \quad b_2 \\ | \text{---} | \\ b_2 \quad a_2 \end{array} \text{ and } 4 \begin{array}{c} a_1 \quad b_2 \\ | \text{---} | \\ b_1 \quad a_2 \end{array} .$$

For the remaining trees, as above, we have two groups of 3 trees that are handled by the Chain Equivalences Lemma, and two that need a particular argument to be cancelled. The two groups of 3 trees are :

Tree	$\begin{array}{c} a_2 \quad b_1 \\ 4 \text{ ---} \\ b_1 \quad a_1 \end{array}$	$\begin{array}{c} a_2 \quad b_1 \\ 2 \text{ ---} \\ b_1 \quad a_2 \end{array}$	$\begin{array}{c} a_2 \quad b_2 \\ 4 \text{ ---} \\ b_1 \quad a_2 \end{array}$
Coefficient	$(u_2^a w_1^b - u_1^b w_2^a) \times (u_1^a w_1^b - u_1^b w_1^a)$	$(u_2^a w_1^b - u_1^b w_2^a) \times (u_2^a w_1^b - u_1^b w_2^a)$	$(u_2^a w_1^b - u_1^b w_2^a) \times (u_2^a w_2^b - u_2^b w_2^a)$

Tree	$\begin{array}{c} a_1 \quad b_2 \\ 4 \text{ ---} \\ b_2 \quad a_2 \end{array}$	$\begin{array}{c} a_1 \quad b_2 \\ 2 \text{ ---} \\ b_2 \quad a_1 \end{array}$	$\begin{array}{c} a_1 \quad b_2 \\ 4 \text{ ---} \\ b_1 \quad a_1 \end{array}$
Coefficient	$(u_1^a w_2^b - u_2^b w_1^a) \times (u_2^a w_2^b - u_2^b w_2^a)$	$(u_1^a w_2^b - u_2^b w_1^a) \times (u_1^a w_2^b - u_2^b w_1^a)$	$(u_1^a w_1^b - u_1^b w_1^a) \times (u_1^a w_2^b - u_2^b w_1^a)$

In each of these tables the 3 trees appear in a Chain of Equivalences as in the Chain Equivalences Lemma 5.3l. Therefore they are mutually equivalent. As before to prove that their sum is cancellable it is enough to prove that the sums of the coefficients in each table is even. Again a direct computation using the fact that  $\omega(u \wedge w) = 1$  yields the result.

The remaining trees are :

Tree	$\begin{array}{c} a_1 \quad b_1 \\ 4 \text{ ---} \\ b_2 \quad a_2 \end{array}$	$\begin{array}{c} a_1 \quad b_2 \\ 4 \text{ ---} \\ a_2 \quad b_1 \end{array}$
Coefficient	$(u_1^a w_2^b - u_2^b w_1^a) \times (u_2^a w_1^b - u_1^b w_2^a)$	$(u_1^a w_2^a - u_2^a w_1^a) \times (u_2^b w_1^b - u_1^b w_2^b)$

The IHX relation applied for instance to the first tree shows that the difference of the above two trees is cancellable. Therefore the sum of the two trees is equivalent to say the first tree with coefficient the difference of the coefficients.

Summing the two cancellable brackets

$$4 \left[ \begin{array}{c} a_3 \quad b_3 \\ \diagdown \quad \diagup \\ a_1 \quad b_2 \end{array}, \begin{array}{c} b_3 \quad a_2 \\ \diagdown \quad \diagup \\ b_1 \quad a_2 \end{array} \right], 4 \left[ \begin{array}{c} -b_3 \quad a_3 \\ \diagdown \quad \diagup \\ a_1 \quad b_2 \end{array}, \begin{array}{c} a_3 \quad a_2 \\ \diagdown \quad \diagup \\ b_1 \quad a_2 \end{array} \right]$$

one finds that

$$8 \begin{array}{c} a_1 \quad b_1 \\ | \quad | \\ \hline | \quad | \\ b_2 \quad a_2 \end{array} \approx 4 \begin{array}{c} a_3 \quad a_2 \\ | \quad | \\ \hline | \quad | \\ b_3 \quad b_2 \end{array} - 4 \begin{array}{c} a_3 \quad a_1 \\ | \quad | \\ \hline | \quad | \\ b_3 \quad b_1 \end{array}$$

is cancellable by the Column Intersection Lemma 5.2.

Therefore it is enough to show that the aforementioned difference of coefficients is even, again this is proved by direct computation.

Q. E. D.

**5.3. Images of genus 2 twists are in  $Can(\mathcal{A}_2(N_1))$ .** As one can see from Theorem 4.3 it remains to prove :

**Theorem 5.3.** *Assume  $g \geq 9$ . For any elements  $u_1, w_1, u_2, w_2 \in N_1$  such that  $\omega(u_1 \wedge w_1) = 1 = \omega(u_2 \wedge w_2)$ , and  $\omega(u_1 \wedge w_2)\omega(u_1 \wedge u_2) = 0 = \omega(w_1 \wedge u_2) = \omega(w_1 \wedge w_2)$ , the tree*

$$4 \begin{array}{c} u_1 \quad w_2 \\ | \quad | \\ \hline | \quad | \\ w_1 \quad u_2 \end{array}$$

is cancellable.

*Proof.* As in the genus 1 case, Using a suitable element in  $GL_g(\mathbf{Z})$  we may suppose that  $u_1, u_2, w_1, w_2 \in \langle a_i, b_i | 1 \leq i \leq 8 \rangle$ .

Write  $u_1$  as  $\tilde{a} + \tilde{b}$  with  $\tilde{a} \in L_A$  and  $\tilde{b} \in L_B$ . Then

$$4 \begin{array}{c} u_1 \quad w_2 \\ | \quad | \\ \hline | \quad | \\ w_1 \quad u_2 \end{array} = 4 \begin{array}{c} \tilde{a} \quad w_2 \\ | \quad | \\ \hline | \quad | \\ w_1 \quad u_2 \end{array} + 4 \begin{array}{c} \tilde{b} \quad w_2 \\ | \quad | \\ \hline | \quad | \\ w_1 \quad u_2 \end{array}$$

Computing the sum of the two cancellable brackets

$$4 \left[ \begin{array}{c} b_9 \quad a_9 \\ \diagdown \quad \diagup \\ \tilde{a} \quad w_1 \quad u_2 \quad w_2 \end{array} \right], 4 \left[ \begin{array}{c} -a_9 \quad b_9 \\ \diagdown \quad \diagup \\ \tilde{b} \quad w_1 \quad u_2 \quad w_2 \end{array} \right]$$

we get that

$$\begin{aligned}
 & 4 \begin{array}{c} u_1 \quad w_2 \\ | \text{---} | \\ w_1 \quad u_2 \end{array} \approx -4\omega(\tilde{a} \wedge u_2) \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array} + 4\omega(\tilde{a} \wedge w_2) \begin{array}{c} w_1 \quad u_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array} + \\
 & 4\omega(\tilde{b} \wedge u_2) \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} - 4\omega(\tilde{b} \wedge w_2) \begin{array}{c} w_1 \quad u_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} .
 \end{aligned}$$

The equations  $\omega(\tilde{a} \wedge u_2) + \omega(\tilde{b} \wedge u_2) = 0 = \omega(\tilde{a} \wedge w_2) + \omega(\tilde{b} \wedge w_2)$  imply that

$$\begin{aligned}
 & 4 \begin{array}{c} u_1 \quad w_2 \\ | \text{---} | \\ w_1 \quad u_2 \end{array} \approx \omega(\tilde{b} \wedge u_2) \left( 4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} + 4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array} \right) + \\
 & \omega(\tilde{a} \wedge w_2) \left( 4 \begin{array}{c} w_1 \quad u_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array} + 4 \begin{array}{c} w_1 \quad u_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} \right) .
 \end{aligned}$$

We only prove that  $4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} + 4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array}$  is cancelable, the proof for the remaining sum is the same.

Applying the IHX relation to this sum we get that

$$4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} + 4 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ b_9 \quad a_9 \end{array} \approx 8 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array} .$$

Expanding by multilinearity  $8 \begin{array}{c} w_1 \quad w_2 \\ | \text{---} | \\ a_9 \quad b_9 \end{array}$ , we find that it is the sum of the following trees, where  $1 \leq i \neq j \leq 8$ , up to a multiplicative coefficient which is irrelevant to the proof :

$$8 \begin{array}{c} a_i \quad b_i \\ | \text{---} | \\ a_9 \quad b_9 \end{array} , 8 \begin{array}{c} b_i \quad a_i \\ | \text{---} | \\ a_9 \quad b_9 \end{array} , 8 \begin{array}{c} a_i \quad b_j \\ | \text{---} | \\ a_9 \quad b_9 \end{array} , 8 \begin{array}{c} b_j \quad a_i \\ | \text{---} | \\ a_9 \quad b_9 \end{array} , 8 \begin{array}{c} a_i \quad a_j \\ | \text{---} | \\ a_9 \quad b_9 \end{array} , 8 \begin{array}{c} b_i \quad b_j \\ | \text{---} | \\ a_9 \quad b_9 \end{array} .$$

Applying the IHX relation to the two first trees we find that they are both the sum of cancelable trees. The last 4 trees are each equal to 2 times a tree

that is involved in a Chain of Equivalences (cf Lemma 5.3), hence they are all cancelable. Q. E. D.

#### ACKNOWLEDGEMENTS

The author would like to thank Professor Jean Barge for his support during the different stages of this work. This paper also benefited from various mathematical discussions on its subject and many others with Professors Jacques Thévenaz from the University of Lausanne, Pierre de la Harpe from the University of Geneva and Jérôme Scherer from the Universitat Autònoma de Barcelona.

#### REFERENCES

1. D. Bar-Natan, *On the Vassiliev knot invariants*, *Topology* **34** (1995), no. 2, 423–472.
2. R. Craggs, *A new proof of the Reidemeister-Singer theorem on stable equivalence of Heegaard splittings*, *Proc. Amer. Math. Soc.* **57** (1976), no. 1, 143–147.
3. S. Garoufalidis and J. Levine, *Tree-level invariants of three-manifolds, Massey products and the Johnson homomorphism*, *Graphs and patterns in mathematics and theoretical physics*, *Proc. Sympos. Pure Math.*, vol. 73, Amer. Math. Soc., Providence, RI, 2005, pp. 173–203.
4. H. B. Griffiths, *Automorphisms of a 3-dimensional handlebody*, *Abh. Math. Sem. Univ. Hamburg* **26** (1963/1964), 191–210.
5. N. Habegger and G. Masbaum, *The Kontsevich integral and Milnor’s invariants*, *Topology* **39** (2000), no. 6, 1253–1289.
6. N. Habegger and W. Pitsch, *Tree level Lie algebra structures of perturbative invariants*, *J. Knot Theory Ramifications* **12** (2003), no. 3, 333–345.
7. Morris W. Hirsch, *Differential topology*, *Graduate Texts in Mathematics*, vol. 33, Springer-Verlag, New York, 1994, Corrected reprint of the 1976 original.
8. D. Johnson, *The structure of the Torelli group. II. A characterization of the group generated by twists on bounding curves*, *Topology* **24** (1985), no. 2, 113–126.
9. J. Levine, *Addendum and correction to: “Homology cylinders: an enlargement of the mapping class group”* [*Algebr. Geom. Topol.* **1** (2001), 243–270; *MR1823501* (2002m:57020)], *Algebr. Geom. Topol.* **2** (2002), 1197–1204 (electronic).

10. W. Magnus, A. Karrass, and D. Solitar, *Combinatorial group theory*, second ed., Dover Publications Inc., Mineola, NY, 2004, Presentations of groups in terms of generators and relations.
11. S. Morita, *Casson's invariant for homology 3-spheres and characteristic classes of surface bundles. I*, *Topology* **28** (1989), no. 3, 305–323.
12. ———, *Abelian quotients of subgroups of the mapping class group of surfaces*, *Duke Math. J.* **70** (1993), no. 3, 699–726.
13. J. Nielsen, *Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen I*, *Acta Math* **50** (1927), 189–358.
14. ———, *Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen II*, *Acta Math* **53** (1929), 1–76.
15. ———, *Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen III*, *Acta Math* **58** (1931), 87–167.
16. W. Pitsch, *Une construction intrinsèque du cœur de l'invariant de Casson*, *Ann. Inst. Fourier (Grenoble)* **51** (2001), no. 6, 1741–1761.
17. ———, *Extensions verselles et automorphismes des groupes nilpotents libres*, *J. Algebra* **249** (2002), no. 2.
18. K. Reidemeister, *Zur dreidimensionalen Topologie*, *Abh. Math. Sem. Univ. Hamburg* (1933), 189–194.
19. J. Singer, *Three-dimensional manifolds and their Heegaard diagrams*, *Trans. Amer. Math. Soc.* **35** (1933), no. 1, 88–111.
20. S. Suzuki, *On homeomorphisms of a 3-dimensional handlebody*, *Canad. J. Math.* **29** (1977), no. 1, 111–124.

WOLFGANG PITSCHE, UNIVERSITAT AUTÒNOMA DE BARCELONA, DEPARTAMENT DE MATEMÀTIQUES E-08193 BELLATERRA, SPAIN

*E-mail address:* pitsch@mat.uab.es