

NOWHERE SCATTERED MULTIPLIER ALGEBRAS

EDUARD VILALTA

ABSTRACT. We study sufficient conditions under which a nowhere scattered C^* -algebra A has a nowhere scattered multiplier algebra $\mathcal{M}(A)$, that is, we study when $\mathcal{M}(A)$ has no nonzero, elementary ideal-quotients. In particular, we prove that a σ -unital C^* -algebra A of

- (i) finite nuclear dimension, or
 - (ii) real rank zero, or
 - (iii) stable rank one with k -comparison,
- is nowhere scattered if and only if $\mathcal{M}(A)$ is.

1. INTRODUCTION

The study of regularity properties of multiplier algebras appears throughout the literature; see, for example, [Ell74], [Zha90], [Rør91], and [KNZ17]. One notable instance of this is the study of pure infiniteness and, more concretely, of when a C^* -algebra has a purely infinite multiplier algebra; see [Lin04], [Lin91], [KNP10], and [KNZ19]. When this condition is relaxed to weak pure infiniteness (in the sense of [KR02]) it was shown in [KR02, Proposition 4.11] that a C^* -algebra is weakly purely infinite if and only if its multiplier algebra is. In general, given a certain property \mathcal{P} , one can ask: If a C^* -algebra A satisfies \mathcal{P} , when does $\mathcal{M}(A)$ satisfy \mathcal{P} ?

In this paper we study the question above for the property of being *nowhere scattered*. This notion ensures sufficient noncommutativity of the algebra, and can be characterized in a number of ways. As shown in [TV21b, Theorem 3.1], a C^* -algebra A is nowhere scattered if and only if it has no nonzero elementary ideal-quotients. This is in turn equivalent to no hereditary sub- C^* -algebra of A admitting a one-dimensional representation. Every weakly purely infinite C^* -algebra is nowhere scattered ([TV21b, Example 3.3]) but, in contrast to the weakly purely infinite case, the multiplier algebra of a nowhere scattered C^* -algebra need not be nowhere scattered; see Examples 5.2 and 5.3.

Question 1.1. Let A be nowhere scattered. When is $\mathcal{M}(A)$ nowhere scattered?

One of the motivations behind Question 1.1 is the study of when a multiplier algebra has no characters, since such a property leads to important structure results. For example, it follows from [Rob16, Theorem 3.2] that every element in a nowhere scattered multiplier algebra can be written as the finite sum of commutators and products of two commutators. Moreover, knowing that $\mathcal{M}(A)$ is nowhere scattered would have implications on its unitary group; see [CR21].

Nowhere scatteredness of a C^* -algebra A can also be characterized in terms of its Cuntz semigroup $\text{Cu}(A)$, a powerful invariant for C^* -algebras introduced in [Cun78] and further developed in [CEI08]; see also [APT18], [APRT22], [TV22a] and [Vil22]. Explicitly, it was shown in [TV21b, Theorem 8.9] that a C^* -algebra is nowhere scattered if and only if its Cuntz semigroup is *weakly* $(2, \omega)$ -divisible, a

Date: 27 April 2023.

The author was partially supported by MINECO (grant No. PID2020-113047GB-I00 and No. PRE2018-083419), and by the Departament de Recerca i Universitats de la Generalitat de Catalunya (grant No. 2021-SGR-01015).

notion defined by Robert and Rørdam in [RR13] to study when certain C^* -algebras have characters; see Paragraph 2.3. Consequently, the study of Question 1.1 leads naturally to the study of divisibility properties of $\text{Cu}(\mathcal{M}(A))$. The main divisibility properties at play in this case are weak $(2, \omega)$ -divisibility and its bounded counterpart, known as *weak (m, n) -divisibility*; see Section 3.

Although the C^* -algebras from Examples 5.2 and 5.3 are nowhere scattered (and thus have a $(2, \omega)$ -divisible Cuntz semigroup), they both have *unbounded divisibility*, that is, for every pair $m, n \in \mathbb{N}$ there exists a Cuntz class that is not weakly (m, n') -divisible for any $n' \leq n$. Thus, we ask:

Question 1.2. Let A be a nowhere scattered C^* -algebra. When does there exist $n \in \mathbb{N}$ such that $[a]$ is weakly $(2, n)$ -divisible for every $a \in A_+$?

More generally, when can one find $m, n \in \mathbb{N}$ such that $[a \otimes 1_m]$ is weakly $(2m, n)$ -divisible for every $a \in A_+$?

This question is studied in detail in Section 4, where we provide a number of examples where one can answer Question 1.2 affirmatively. Further, we also show that having bounded divisibility is characterized by the soft part of the monoid; see Theorem 4.10.

We prove in Section 5 that Question 1.1 has a positive answer whenever Question 1.2 does.

Theorem 1.3 (5.11). *Let A be a σ -unital C^* -algebra. Assume that for every orthogonal sequence $(a_i)_i$ of positive elements in A there exist $m, n \in \mathbb{N}$ such that $[a_i \otimes 1_m]$ is weakly $(2m, n)$ -divisible for every i . Then $\mathcal{M}(A)$ is nowhere scattered.*

Using the study of bounded divisibility from Section 4, and that nowhere scatteredness passes to ideals ([TV21b, Proposition 4.2]), one obtains:

Theorem 1.4 (5.12). *Let A be a σ -unital C^* -algebra. Assume that A is of*

- (i) *real rank zero, or*
- (ii) *finite nuclear dimension, or*
- (iii) *stable rank one with k -comparison.*

Then A is nowhere scattered if and only if $\mathcal{M}(A)$ is nowhere scattered.

Further, one can generalize (iii) above by changing stable rank one for the condition of having a *surjective rank map*; see Proposition 4.3 and Remark 4.4.

We finish the section by proving a weak converse of Theorem 1.3 for stable C^* -algebras:

Theorem 1.5 (5.18). *Let A be a σ -unital, stable C^* -algebra. Assume that $\mathcal{M}(A)$ is nowhere scattered. Then, for every $a \in A_+$ and $m \in \mathbb{N}$ there exists n such that $[a]$ is weakly (m, n) -divisible.*

Most of the results in Sections 3 and 4 can be translated directly to the more general setting of abstract Cuntz semigroups, or Cu -semigroups for short; see for example [APT18] and [GP22]. However, since in this paper we focus on multiplier algebras (which have no known Cu -counterpart), we state all the results in the language of C^* -algebras to ease the notation.

Acknowledgements. The author thanks Hannes Thiel for his comments on a first draft of this paper.

2. PRELIMINARIES

2.1 (Nowhere scatteredness and the Global Glimm Property). As defined in [TV21b, Definition A], a C^* -algebra is *nowhere scattered* if none of its quotients contains a

minimal open projection. By [TV21b, Theorem 3.1], a C^* -algebra A is nowhere scattered if and only if no nonzero ideal-quotient of A is elementary.

One also says that a C^* -algebra A has the *Global Glimm Property* ([KR02, Definition 4.12]) if for every $a \in A_+$ and $\varepsilon > 0$ there exists a $*$ -homomorphism $\varphi: M_2(C_0(0, 1]) \rightarrow \overline{aAa}$ such that the image of φ contains $(a - \varepsilon)_+$.

A C^* -algebra is nowhere scattered whenever it has the Global Glimm Property. The converse remains open, and is known as the *Global Glimm Problem*; see [APRT22], [ER06], and [TV22b].

Examples of C^* -algebras with the Global Glimm Property include simple, non-elementary C^* -algebras, \mathcal{Z} -stable C^* -algebras, and purely infinite C^* -algebras. Traceless C^* -algebras are all nowhere scattered.

2.2 (The Cuntz semigroup). For any pair of positive elements a, b in a C^* -algebra A one writes $a \precsim b$ if $a = \lim_n r_n b r_n^*$ for some sequence $(r_n)_n$ in A . One also says that a is *Cuntz equivalent* to b , in symbols $a \sim b$, whenever $a \precsim b$ and $b \precsim a$.

The *Cuntz semigroup* $\text{Cu}(A)$ is defined as the quotient $(A \otimes \mathcal{K})_+ / \sim$ equipped with the order induced by \precsim and the addition induced by diagonal addition; see [Cun78] and [CEI08] for details.

Given elements $x, y \in \text{Cu}(A)$, we write $x \ll y$ whenever there exists $a \in (A \otimes \mathcal{K})_+$ and $\varepsilon > 0$ such that $x \leq [(a - \varepsilon)_+]$ and $y = [a]$. As shown in [CEI08], every increasing sequence in $\text{Cu}(A)$ has a supremum, and every element can be written as the supremum of a \ll -increasing sequence.

In recent years, the Cuntz semigroup has benefitted from the study of abstract Cuntz semigroups, or *Cu-semigroups* for short. Good references for this include [APT11], [APT18], and [GP22]. In this paper, we will not use the language of Cu-semigroups (except in Corollary 4.12), but we will still make use of some of the abstract properties that the Cuntz semigroup of a C^* -algebra always satisfies. These are:

- (O5) Given $x', x, y', y, z \in \text{Cu}(A)$ such that $x + y \leq z$ with $x' \ll x$ and $y' \ll y$, there exists $c \in \text{Cu}(A)$ such that $y' \ll c$ and $x' + c \leq y \leq x + c$.
- (O6) Given $x' \ll x \ll y + z$, there exist $v \leq x, y$ and $w \leq x, z$ such that $x' \leq v + w$.

The reader is referred to [APT18, Proposition 4.6] (also [RW10]) and [Rob13] for the respective proofs.

2.3 (Divisibility in the Cuntz semigroup). Let A be a C^* -algebra, let $x \in \text{Cu}(A)$ and take $n \geq 1$ and $m \geq 2$. Following [RR13], we will say that x is *weakly (m, n) -divisible* if, whenever $x' \ll x$, there exist elements y_1, \dots, y_n such that $x' \leq y_1 + \dots + y_n$ and $my_j \leq x$ for each $j \leq n$. Similarly, x is *weakly (m, ω) -divisible* if the previous condition holds without any bound on n , that is, allowing n to depend on x' .

The element x is said to be *(m, n) -divisible* (resp. *(m, ω) -divisible*) if one can always set $y_1 = \dots = y_n$.

A C^* -algebra is nowhere scattered if and only if every element in its Cuntz semigroup is weakly (m, ω) -divisible for each m , whilst a C^* -algebra has the Global Glimm Property if and only if every element is (m, ω) -divisible for each m ; see [TV21b, Theorem 8.9] and [TV22b, Theorem 3.6] respectively.

3. FINITE DIVISIBILITY

In this section we recall the notions of finite divisibility introduced in [RR13]; see Definition 3.1. We prove their main properties (Lemma 3.3), and study some situations where $\text{Cu}(A)$ contains a sup-dense subset of elements with finite divisibility. We also find sufficient conditions for finite weak divisibility to imply finite divisibility; see Theorem 3.14.

Definition 3.1 ([RR13]). Let a be a positive element of a C^* -algebra A , and let $m \geq 2$. We let $\text{div}_m([a])$ and $\text{Div}_m([a])$ be the least positive integers n, n' such that $[a] \in \text{Cu}(A)$ is weakly (m, n) -divisible and (m, n') -divisible respectively.

If no such n or n' exist, we set $\text{div}_m([a]) = \infty$ or $\text{Div}_m([a]) = \infty$.

Remark 3.2. The class of a positive element $a \in A_+$ in $\text{Cu}(A)$ is said to be *compact* if $[a] \ll [a]$. Every projection gives rise to a compact Cuntz class and, in some cases, these are the only compact classes; see [BC09].

Given a compact element $[a] \in \text{Cu}(A)$, then $[a]$ is weakly (m, ω) -divisible if and only if $\text{div}_m([a]) < \infty$.

However, for a non-compact element $[a]$, $\text{div}_m([a]) < \infty$ is not equivalent to $[a]$ being weakly (m, ω) -divisible; see Example 3.6 and Remark 5.10.

Let us first summarize the main properties of $\text{div}_m()$. Part (iii) of the following lemma is in analogy to [KR02, Lemma 4.9], while (iv) is a Cu-analogue of [RR13, Proposition 3.6].

Recall that, given $x, y \in \text{Cu}(A)$, the infimum $x \wedge \infty y$ always exists; see [APRT21, Remark 2.6]. Here, ∞y denotes the element $\sup_n ny$.

Lemma 3.3. *Let A be a C^* -algebra, and let $x = [a] \in \text{Cu}(A)$ and $m \in \mathbb{N}$. Then,*

- (i) $\text{div}_m(x) \leq \text{div}_{m'}(x)$ whenever $m \leq m'$.
- (ii) $\text{div}_m([a]) \leq N(\text{div}_m([b_1]) + \dots + \text{div}_m([b_r]))$ whenever
$$[b_1], \dots, [b_r] \leq [a] \leq N([b_1] + \dots + [b_r]).$$
- (iii) $\text{div}_m([a + b]) \leq \text{div}_m([a]) + \text{div}_m([b])$.
- (iv) Given $x, y \in \text{Cu}(A)$ such that $y \leq x$ and $x \wedge \infty y = y$, then $\text{div}_m(y) \leq \text{div}_m(x)$.
- (v) $\text{div}_m(x) \leq \sup_k \text{div}_m(x_k)$ for every increasing sequence $(x_k)_k$ in $\text{Cu}(A)$ with supremum x .

Proof. (i) follows directly from the definition of $\text{div}_m([a])$.

To see (ii), let $n_j = \text{div}_m([b_j])$ for $j = 1, \dots, r$. We may assume that all quantities are finite, since otherwise we are done.

Take $x \ll [a]$. Since $[a] \leq N[b_1] + \dots + N[b_r]$, we can find elements x_j such that $x_j \ll [b_j]$ and $x \leq Nx_1 + \dots + Nx_r$. The element $[b_j]$ is weakly (m, n_j) -divisible, so we can find elements $y_{1,j}, \dots, y_{n_j,j}$ such that

$$my_{i,j} \leq [b_j], \quad \text{and} \quad x_j \leq y_{1,j} + \dots + y_{n_j,j}$$

for each i and j .

Using that $[b_j] \leq [a]$, one has $my_{i,j} \leq [a]$. Further, since we also have $x \leq N \sum_j x_j$, we deduce that $x \leq N \sum_{i,j} y_{i,j}$. It follows that $\text{Div}_m([a]) \leq N(n_1 + \dots + n_r)$, as desired.

Note that, given $a, b \in (A \otimes \mathcal{K})_+$, we have $[a + b] \leq [a] + [b]$ and $[a], [b] \leq [a + b]$. Thus, (iii) follows directly from (ii).

For (iv), let $x, y \in \text{Cu}(A)$ be as stated. As before, we may assume that $\text{div}_m(x) = n$ for some $n \in \mathbb{N}$. Take $y' \ll y$, and apply the weak (m, n) -divisibility of x to obtain elements y_1, \dots, y_n satisfying the properties from Paragraph 2.3 for y' and x .

Since $my_j \leq x$ for every j , and $x \wedge \infty y = y$, one obtains $m(y_j \wedge \infty y) \leq x \wedge \infty y = y$. It is now easy to check that the elements $y_j \wedge \infty y$ satisfy the conditions from Paragraph 2.3 for y' and y , as required.

To prove (v), assume that the supremum $\sup_k \text{div}_m(x_k)$ is finite, since otherwise there is nothing to prove. Let $n \in \mathbb{N}$ be such that $\sup_k \text{div}_m(x_k) \leq n$, and take $x' \in \text{Cu}(A)$ such that $x' \ll x$. Since $x = \sup_k x_k$, it follows that $x' \ll x_k$ for some $k \in \mathbb{N}$. Thus, using that $\text{div}_m(x_k) \leq n$, we obtain elements y_1, \dots, y_n with

$x' \leq y_1 + \dots + y_n$ and $my_j \leq x_k \leq x$ for each j . We get that $\text{div}_m(x) \leq n$, as desired. \square

Recall that a C^* -algebra A is said to have *strict comparison* if $[a] \leq [b]$ in $\text{Cu}(A)$ whenever there exists $\delta > 0$ such that $\lim_n \tau(a^{1/n}) \leq (1 - \delta) \lim_n \tau(b^{1/n})$ for every 2-quasitrace τ ; see [ERS11] for more details.

Lemma 3.4. *Let A be a C^* -algebra, let $[a] \in \text{Cu}(A)$ and $m \in \mathbb{N}$. Then,*

$$\text{div}_{km}(k[a]) \leq \text{div}_m([a])$$

for every $k \in \mathbb{N}$.

If A has strict comparison, one has $\text{div}_m([a]) \leq \text{div}_{k(m+1)}(k[a])$.

Proof. Let $n = \text{div}_m([a])$, which we may assume to be finite, and take $x \ll k[a]$. Then, there exists $\varepsilon > 0$ such that $x \ll k[(a - \varepsilon)_+]$. Since $[(a - \varepsilon)_+] \ll [a]$, there exist $y_1, \dots, y_n \in \text{Cu}(A)$ such that

$$[(a - \varepsilon)_+] \leq y_1 + \dots + y_n, \quad \text{and} \quad my_j \leq [a]$$

for each j .

In particular, one gets

$$x \ll k[(a - \varepsilon)_+] \leq ky_1 + \dots + ky_n, \quad \text{and} \quad m(ky_j) \leq k[a],$$

which implies $\text{div}_{km}(k[a]) \leq n$, as desired.

Now assume that A has strict comparison of positive elements, and let $n = \text{div}_{k(m+1)}(k[a])$. As before, we may assume n to be finite. Let $x \ll [a]$, which implies $kx \ll k[a]$. Then, there exist z_1, \dots, z_n such that $x \leq kx \leq z_1 + \dots + z_n$ and $k(m+1)z_j \leq k[a]$. Thus, one has $(km+1)mz_j \leq km(m+1)z_j \leq (km)[a]$.

Using [ERS11, Proposition 6.2], we get $mz_j \leq [a]$. This implies $\text{div}_m([a]) \leq \text{div}_{k(m+1)}(k[a])$. \square

Remark 3.5. Let A be a C^* -algebra and let $m \in \mathbb{N}$. Assume that $\text{div}_m([a]) < \infty$ for every $a \in A_+$. Lemma 3.3 (ii) implies, in particular, that $\text{div}_m([b]) < \infty$ for every $b \in M_n(A)_+$ and $n \in \mathbb{N}$.

Indeed, given $b \in M_n(A)_+$ there exist elements $b_1, \dots, b_n \in A_+$ and $N \in \mathbb{N}$ such that $[b_i] \leq [b] \leq N([b_1] + \dots + [b_n])$ for each $i \leq n$; see [APT20, 4.2] and [TV22b, Lemma 3.3].

By Lemma 3.3 (ii), we have that $\text{div}_m(b) < \infty$, as desired.

Example 3.6. Let $(A_k)_k$ be a family of unital C^* -algebras such that

$$k \leq \text{div}_2([1_{A_k}]) \leq 7k$$

for each k .

By [RR13, Theorem 7.9], such C^* -algebras can be taken to be simple, unital, infinite dimensional AH-algebras. Using [TV21b, Proposition 4.13], one sees that $A := \oplus_k A_k$ is nowhere scattered. Thus, every element in $\text{Cu}(A)$ is weakly $(2, \omega)$ -divisible.

Consider the element $x = \sup_n \sum_{k=1}^n [1_{A_k}] \in \text{Cu}(\oplus_k A_k)$, and denote by ι_k the induced inclusion $\text{Cu}(A_k) \rightarrow \text{Cu}(\oplus_k A_k)$. Then,

$$\iota_k([1_{A_k}]) \leq x, \quad \text{and} \quad x \wedge \iota_k([1_{A_k}]) = \iota_k([1_{A_k}]).$$

Thus, Lemma 3.3 (iv) implies that

$$k \leq \text{div}_2(\iota_k([1_{A_k}])) \leq \text{div}_2(x).$$

This shows that x is weakly $(2, \omega)$ -divisible, but $\text{div}_2(x) = \infty$; see also Corollary 5.16.

We now move our attention to C^* -algebras whose Cuntz semigroup $\text{Cu}(A)$ satisfies that every element (or, at least, a finite multiple of it) has finite weak divisibility.

To study such C^* -algebras, we will need the following definitions:

Definition 3.7. Let A be a C^* -algebra, and let $a \in A_+$. We set

$$\begin{aligned} \text{mDiv}([a]) &= \inf\{\text{Div}_{2m}(m[a]) \mid m \in \mathbb{N}\} \\ \text{mdiv}([a]) &= \inf\{\text{div}_{2m}(m[a]) \mid m \in \mathbb{N}\} \end{aligned}$$

We will denote by $\text{Cu}(A)_{\text{div}}$ (respectively $\text{Cu}(A)_{\text{mdiv}}$ and $\text{Cu}(A)_{\text{mDiv}}$) the subset of $\text{Cu}(A)$ consisting of the elements $[a] \in \text{Cu}(A)$ such that $\text{div}_{2m}([a]) < \infty$ for some $m \geq 1$ (respectively $\text{mdiv}([a]) < \infty$ and $\text{mDiv}([a]) < \infty$). By Lemma 3.3, the subsets $\text{Cu}(A)_{\text{div}}$ and $\text{Cu}(A)_{\text{mdiv}}$ are always subsemigroups of $\text{Cu}(A)$.

Note that one has $\text{Cu}(A)_{\text{div}} \subseteq \text{Cu}(A)_{\text{mdiv}}$, and it is easy to see that $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)_{\text{mdiv}}$ whenever $\text{Cu}(A)$ is unperforated.

Example 3.8. If $A = \mathbb{C}$, its Cuntz semigroup is isomorphic to $\overline{\mathbb{N}} := \mathbb{N} \cup \{\infty\}$. It is readily checked that $\text{div}_m(x) < \infty$ if and only if $x \geq m$. Thus, one has $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)_{\text{mdiv}} = \{0, 2, 3, \dots, \infty\}$.

Example 3.9. Every tracially \mathcal{Z} -stable C^* -algebra A satisfies $\text{Cu}(A) = \text{Cu}(A)_{\text{div}}$. More generally, recall that a C^* -algebra is said to be *N -almost divisible* (see [Win12]) if for each pair $x', x \in \text{Cu}(A)$ satisfying $x' \ll x$, and $k \in \mathbb{N}$, there exists $y \in \text{Cu}(A)$ such that

$$ky \leq x, \quad \text{and} \quad x' \leq (k+1)(N+1)y.$$

Thus, if A is N -almost divisible for some $N \in \mathbb{N}$, one also has $\text{Cu}(A) = \text{Cu}(A)_{\text{div}}$.

By [RT17, Theorem 3.1], every nowhere scattered C^* -algebra of nuclear dimension N that has no nonzero, simple, purely infinite ideal-quotients is N -almost divisible. This includes residually stably finite, nowhere scattered C^* -algebras of finite nuclear dimension, and nowhere scattered C^* -algebras of finite decomposition rank. Consequently, one has $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$.

As we will see in Proposition 4.1, $\text{Cu}(A)_{\text{mdiv}} = \text{Cu}(A)$ for any nowhere scattered C^* -algebra of finite nuclear dimension. In fact, one has more: $\text{Cu}(\oplus_{i=1}^{\infty} A)_{\text{mDiv}} = \text{Cu}(\oplus_{i=1}^{\infty} A)$.

Example 3.10. As shown in [TV21b, Theorem 9.1], every element x in the Cuntz semigroup of a nowhere scattered, real rank zero C^* -algebra is *weakly divisible*, that is, there exist y, z such that $x = 2y + 3z$. In particular, one gets

$$2(y+z) \leq x \leq 3(y+z).$$

Thus, it follows that $\text{div}_2(x) \leq 3$ for every x and, consequently, that $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$.

As noted in Example 3.10 above, every element in the Cuntz semigroup of a nowhere scattered, real rank zero C^* -algebra has finite divisibility. Lemmas 3.11 and 3.13 below show that the same is true for a sup-dense subset of (non-elementary) simple C^* -algebras and nowhere scattered C^* -algebras of topological dimension zero. However, this may not imply that every element in such a C^* -algebra has finite divisibility.

Recall that an element $x \in \text{Cu}(A)$ is *idempotent* if $x = 2x$. In the Cuntz semigroup of a simple C^* -algebra there is only one nonzero, idempotent element, which we denote by ∞ .

Lemma 3.11. *Let A be a simple, non-elementary C^* -algebra. Then, $\text{Cu}(A)_{\text{div}}$ is sup-dense in $\text{Cu}(A)$.*

If A is not stably finite, then $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$.

Proof. Let $m \in \mathbb{N}$. If A is simple and non-elementary, it follows from [Rob13, Proposition 5.2.1] that $\text{Cu}(A)$ is (m, ω) -divisible.

Let $x \in \text{Cu}(A)$ be such that $x \ll \infty$. Then, using that x is (m, ω) -divisible, there exists a nonzero element $y \in \text{Cu}(A)$ such that $my \leq x$. Using that $x \ll \infty = \infty y$, we find $n \in \mathbb{N}$ such that $x \leq ny$. This implies that $\text{div}_m(x) \leq n$. Since the subset of elements compactly contained in ∞ is sup-dense in $\text{Cu}(A)$, the result follows.

Now assume that A is not stably finite. Then, the element ∞ in $\text{Cu}(A)$ is compact; see, for example, [Thi17, Lemma 6.21] or the proof of [Bos22, Theorem 2.6].

It follows that every element in $\text{Cu}(A)$ is compactly contained in ∞ . Using the argument above, we see that $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$. \square

Remark 3.12. An inspection of the proof shows that Lemma 3.11 is slightly more general: For a C^* -algebra A , one has $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$ whenever every idempotent element in $\text{Cu}(A)$ is compact.

As defined in [BP09, Remark 2.5 (vi)], a C^* -algebra has *topological dimension zero* whenever its primitive ideal space has a basis of compact-open subsets. Examples include all C^* -algebras with the ideal property.

Lemma 3.13. *Let A be a separable nowhere scattered C^* -algebra of topological dimension zero. Then $\text{Cu}(A)_{\text{div}}$ is dense in $\text{Cu}(A)$.*

Proof. Take $x \in \text{Cu}(A)$, and assume that there exists $x' \in \text{Cu}(A)$ such that $x' \ll x \ll \infty x'$.

Let $M \in \mathbb{N}$ be such that $x \leq Mx'$. Using that x is weakly (m, ω) -divisible, one finds y_1, \dots, y_n such that $my_i \leq x$ and $x' \leq y_1 + \dots + y_n$. Thus, one has $x \leq My_1 + \dots + My_n$, and we get $\text{div}_m(x) \leq Mn$.

It follows from [TV22b, Proposition 4.18] that such elements x are sup-dense in $\text{Cu}(A)$, as required. \square

3.1. The Global Glimm Problem. We finish this section by studying when $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$ implies $\text{Cu}(A)_{\text{Div}} = \text{Cu}(A)$. More concretely, we study when an element of finite weak divisibility has finite divisibility. Given its similarities with the Global Glimm Problem (see Paragraph 2.1), one could call this the *discrete Global Glimm Problem*.

Following the ideas from [TV22b, Section 6], we obtain:

Theorem 3.14. *Let A be a C^* -algebra satisfying $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$. Assume that there exist $k \in \mathbb{N}$ and maps $N, M: \mathbb{N} \rightarrow \mathbb{N}$ such that*

- (i) *whenever $x' \ll x \leq ny, nz$, there exists $t \in \text{Cu}(A)$ such that $x' \ll N(n)t$ and $t \ll y, z$;*
- (ii) *whenever $x' \ll x$ and $2x \ll y + 2nz$, there exists $g \in \text{Cu}(A)$ such that $2g \ll y$ and $x' \ll g + M(n)z$;*
- (iii) *whenever $x_1 + f, x_2 + f \ll y$ and $x'_i \ll x_i \ll f$ for $i = 1, 2$, one can find $z_1, z_2 \in \text{Cu}(A)$ such that $z_1 + z_2 \ll y$ and $x'_1 + x'_2 \ll kz_1, kz_2$.*

Then, $\text{Cu}(A)_{\text{Div}} = \text{Cu}(A)$.

Proof. Assume that (i)-(iii) are satisfied, and let $x \in \text{Cu}(A)$. By assumption, we have $\text{div}_m(x) < \infty$, which implies $\text{div}_2(x) < \infty$. Set $n := \text{div}_2(x)$. We will show that $\text{Div}_2(x) < \infty$ by proving that

$$\text{Div}_2(x) \leq \left(N(\max\{N(2N_{2,n}), M(n)\}) + N(N(2N_{2,n})) \right) N(k),$$

where $N_{2,n} := (N \circ \cdot n \circ N)(2)$.

Let $x' \in \text{Cu}(A)$ be such that $x' \ll x$, and take y', y such that $x' \ll y' \ll y \ll x$. Then, since x is $(2, n)$ -divisible, we obtain y_1, \dots, y_n satisfying $y \ll \sum y_j$ and

$2y_j \leq x$ for each j . Take $y'_j \ll y_j$ such that $y \ll \sum y'_j$. By [TV22b, Lemma 2.2], for each j we obtain an element r_j such that $y_j + r_j \leq x \leq 2r_j$ and $y'_j \leq r_j$.

Using (i), we find $r \in \text{Cu}(A)$ satisfying $y \ll N_{2,n}r$ and $r \leq r_j$ for every j . Take $r' \ll r$ such that $y \ll N_{2,n}r'$. Using (O5) at $r' \ll r \leq x$, we obtain $c \in \text{Cu}(A)$ such that $r' + c \leq x \leq r + c$.

In particular, since one has $y_j + r \leq x \leq r + c$, we obtain

$$\begin{aligned} 2y &\leq 2y_1 + \dots + 2y_n \leq (2y_1 + \dots + 2y_n) + r \\ &\leq (y_1 + 2y_2 + \dots + 2y_n) + r + c \leq \dots \leq r + (2n)c. \end{aligned}$$

Using (ii), we find elements $g', g \in \text{Cu}(A)$ such that

$$2g \ll r, \quad y' \ll g' + M(n)c, \quad \text{and} \quad g' \ll g.$$

Applying [TV22b, Lemma 2.2] at $2g \ll r$, we find $d \in \text{Cu}(A)$ such that $g' + d \leq r \leq 2d$. Thus, we have $y' \ll y \leq N_{2,n}r' \leq 2N_{2,n}d$. By (i), we find $f \in \text{Cu}(A)$ such that

$$y' \leq N(2N_{2,n})f, \quad \text{and} \quad f \ll r', d.$$

By (O6) applied at $x' \ll y' \ll g' + M(n)c$, one finds elements $s', s, t', t \in \text{Cu}(A)$ such that

$$x' \ll s' + t', \quad s' \ll s \leq y', g', \quad \text{and} \quad t' \ll t \leq y', M(n)c.$$

Set $N_1 := N(\max\{N(2N_{2,n}), M(n)\})$ and $N_2 := N(N(2N_{2,n}))$. Applying (i) at $t' \ll t \leq M(n)c$ and $t' \ll t \leq y' \leq N(2N_{2,n})f$, we obtain $x_1 \in \text{Cu}(A)$ such that $x_1 \ll c, f$ and $t' \ll N_1 x_1$. Take $x'_1, x''_1 \in \text{Cu}(A)$ such that $x'_1 \ll x''_1 \ll x_1$ and $t' \leq N_1 x'_1$.

Using (i) again, but this time at $s' \ll s \leq y' \leq N(2N_{2,n})f$, we obtain $x_2 \in \text{Cu}(A)$ such that $x_2 \ll f, s$ and $s' \ll N_2 x_2$. As before, take $x'_2, x''_2 \in \text{Cu}(A)$ such that $x'_2 \ll x''_2 \ll x_2$ and $s' \leq N_2 x'_2$.

One has

$$x_1 + f \ll c + r' \leq x, \quad x_2 + f \ll s + d \leq g' + d \leq r \leq x, \quad \text{and} \quad x_1, x_2 \ll f.$$

By (iii), we find $z_1, z_2 \in \text{Cu}(A)$ such that

$$z_1 + z_2 \leq x, \quad \text{and} \quad x'_1 + x'_2 \ll kz_1, kz_2.$$

Applying (i) one last time, we find $z \in \text{Cu}(A)$ such that

$$z \leq z_1, z_2, \quad \text{and} \quad x'_1 + x'_2 \leq N(k)z.$$

The element z satisfies $2z \leq z_1 + z_2 \leq x$, and

$$x' \leq t' + s' \leq (N_1 + N_2)(x'_1 + x'_2) \leq (N_1 + N_2)N(k)z,$$

as desired. \square

One can check that real rank zero, or stable rank one C^* -algebras satisfy (i)-(iii) in Theorem 3.14 above. In particular, this recovers the discrete part of [APRT22, Theorem 5.5].

Corollary 3.15. *Let A be a C^* -algebra. Assume that A is either of stable rank one, or real rank zero. Then, $\text{Cu}(A)_{\text{div}} = \text{Cu}(A)$ if and only if $\text{Cu}(A)_{\text{Div}} = \text{Cu}(A)$.*

Proof. If A has real rank zero, use Example 3.10. For stable rank one C^* -algebras, this just amounts to an inspection of a number of proofs:

[APRT22, Lemma 5.3] implies that (i) is satisfied with $N = 2n - 1$.

The proof of [TV22b, Lemma 5.5], but using weak cancellation instead of residually stably finiteness, gives (ii) in Theorem 3.14 with $k = 2$.

Finally, (iii) follows from an inspection of [TV21b, 7.6-7.8] using that, when A has stable rank one, (O8) can be used without the assumption of an element being idempotent; see Definition 7.2 and Proposition 7.5 in [TV21b] for more details. \square

Remark 3.16. One can actually show that $\text{Cu}(A)_{\text{mdiv}} = \text{Cu}(A)$ if and only if $\text{Cu}(A)_{\text{mDiv}} = \text{Cu}(A)$ whenever A is of stable rank one. Indeed, given $x \in \text{Cu}(A)$ such that $\text{div}_{2m}(mx) < \infty$, it follows directly from [APRT22, Theorem 5.5] that $\text{Div}_{2m}(mx) < \infty$.

The same result holds for real rank zero C^* -algebras by Example 3.10.

Question 3.17. Let A be a C^* -algebra such that $\text{Cu}(A) = \text{Cu}_{\text{mdiv}}(A)$. When does A satisfy $\text{Cu}(A) = \text{Cu}_{\text{mDiv}}(A)$?

4. BOUNDED DIVISIBILITY

We focus in this section on C^* -algebras where there is a global bound for the divisibility of the elements. That is, we look at those C^* -algebras A such that there exist $n, m \in \mathbb{N}$ with $\text{div}_m([a]) \leq n$ for every $a \in A_+$. More generally, we focus on those that satisfy $\sup_{a \in A_+} \text{div}_{2m}(m[a]) < \infty$ for some $m \in \mathbb{N}$. As we will see in Theorem 5.11, these algebras will have a nowhere scattered multiplier algebra.

We begin the section by providing sufficient conditions for this bounded divisibility to occur (Propositions 4.1 and 4.3), and by showing that C^* -algebras with this property are closed under extensions (Proposition 4.5). Note that we have already seen some examples, such as tracially \mathcal{Z} -stable C^* -algebras, or C^* -algebras of real rank zero; see Examples 3.9 and 3.10.

In Theorem 4.10, we prove that the set of soft elements characterizes when a C^* -algebra has multiples of bounded divisibility, and we use this to deduce the property for simple, weakly cancellative C^* -algebras of Cuntz covering dimension zero; see Corollary 4.12.

As mentioned in [RT17], it is unclear if $\dim_{\text{nuc}}(A) < \infty$ (with A nowhere scattered) implies $\text{div}(a) < \infty$ for every $a \in A_+$. However, one stills gets the following result, which is a direct consequence of [RT17, Proposition 3.2].

Proposition 4.1. *Let A be a nowhere scattered C^* -algebra of finite nuclear dimension at most k . Then, for every $[a] \in \text{Cu}(A)$, one has*

$$\text{div}_{4(k+1)}(2(k+1)[a]) \leq 8(k+1)^2.$$

In particular, in a nowhere scattered C^ -algebra of finite nuclear dimension, one always gets $\text{mdiv}([a]) < \infty$.*

Proof. If A is nowhere scattered and of nuclear dimension at most k , so is $A \otimes \mathcal{K}$.

Let $[a] \in \text{Cu}(A)$. Upon passing to the hereditary C^* -algebra $a(A \otimes \mathcal{K})a$, we may assume that a is full in $A \otimes \mathcal{K}$; note that $\overline{a(A \otimes \mathcal{K})a}$ is still nowhere scattered by [TV21b, Proposition 4.1], and has nuclear dimension at most k by [WZ10, Proposition 2.5].

Recall from [TV21b, Theorem 3.1] that $A \otimes \mathcal{K}$ is nowhere scattered if and only if it has no finite-dimensional irreducible representations. By [RT17, Proposition 3.2], there exists $[b] \in \text{Cu}(A)$ such that

$$[(a - \varepsilon)_+] \leq 4(k+1)[b], \quad \text{and} \quad 4(k+1)[b] \leq 2(k+1)[a].$$

Thus, we obtain that $2(k+1)[(a - \varepsilon)_+] \leq 8(k+1)^2[b]$. Since $2(k+1)[a]$ can be written as the supremum of the elements $2(k+1)[(a - \varepsilon)_+]$ with $\varepsilon \rightarrow 0$, it follows that $\text{div}_{4(k+1)}(2(k+1)[a]) < 8(k+1)^2$, as desired. \square

Let us denote by $F(A)$ the central sequence algebra, as defined in [Kir06, Definition 1.1]. Given a separable, unital C^* -algebra A , it is not known if A being \mathcal{Z} -stable is equivalent to $F(A)$ admitting no characters; see [KR15]. Although this question remains open, one does have the following result from [Chr17, Part II, Article B, Proposition 2.8].

Proposition 4.2 ([Chr17]). *Let A be a separable C^* -algebra. Assume that $F(A)$ admits no characters. Then, for every $m \in \mathbb{N}$ there exists n such that $\text{div}_m([a]) \leq n$ for all $a \in A_+$.*

Proof. It follows from [RR13, Corollary 5.6] that $F(A)$ has no characters if and only if $\text{div}_2([1]) < \infty$ in $\text{Cu}(F(A))$. Let $n_0 = \text{div}_2([1])$. [Chr17, Part II, B, Proposition 2.8] implies that $\text{div}_2([a]) \leq n_0$ for every $a \in A_+$.

An inductive argument now shows that $\sup_{a \in A_+} \text{div}_m([a]) < \infty$ for every $m \in \mathbb{N}$; see, for example, [TV22b, Lemma 3.4]. \square

Given $[a], [b] \in \text{Cu}(A)$, recall that we write $[a] <_s [b]$ whenever there exists $\gamma < 1$ such that $\lambda([a]) \leq \gamma \lambda([b])$ for every $\lambda \in F(\text{Cu}(A)) \cong \text{QT}(A)$; see [RT17, 2.1].

Let $k \in \mathbb{N}$. As defined in [Win12, Definition 2.1], a C^* -algebra A is said to have *k-comparison* if for every $[a], [b_0], \dots, [b_k] \in \text{Cu}(A)$ such that $[a] <_s [b_i]$ for all i , we have $[a] \leq \sum_i [b_i]$.

Also recall that A has a *surjective rank map* if every element in $L(F(\text{Cu}(A)))$ can be realized as a rank function; see [APRT22, Section 7] for details.

Proposition 4.3. *Let A be a nowhere scattered C^* -algebra with k -comparison. Assume that the family of separable sub- C^* -algebras of A with a surjective rank map is σ -complete and cofinal. Then, $\text{Div}_{2(k+1)}((k+1)[a]) \leq 4(k+1)^3$ for every $[a] \in \text{Cu}(A)$.*

Proof. Let $x := [a]$. By [TV21a, Proposition 6.1] and [FHL⁺21, Proposition 3.8.1], there exists a separable sub- C^* -algebra $B \subseteq A$ that has a surjective rank map, is nowhere scattered, contains x , and is such that the induced inclusion map $\text{Cu}(B) \rightarrow \text{Cu}(A)$ is an order-embedding.

By definition, every element in $L(F(\text{Cu}(B)))$ can be realized as a rank function. Thus, there exists $y \in \text{Cu}(B)$ such that $\hat{y} = \frac{1}{3(k+1)}\hat{x}$. This implies, in particular, that

$$2(k+1)\hat{y} \leq \frac{2}{3}\hat{x}, \quad \text{and} \quad \hat{x} \leq 3(k+1)\hat{y} = \frac{3}{4}(4(k+1))\hat{y}.$$

Applying k -comparison, one obtains $2(k+1)y \leq (k+1)x$ and $x \leq 4(k+1)^2y$ in $\text{Cu}(B)$. Using that the map $\text{Cu}(B) \rightarrow \text{Cu}(A)$ is an order-embedding, the same is true in $\text{Cu}(A)$. Thus, we have

$$2(k+1)y \leq (k+1)x \leq 4(k+1)^3y,$$

as desired. \square

Remark 4.4. As shown in [APRT22, Theorem 7.14], a separable, nowhere scattered C^* -algebra of stable rank one always has a surjective rank map. Thus, Proposition 4.3 above applies to all stable rank one C^* -algebras with k -comparison.

In fact, it was shown in [APRT22, Theorem 8.12] that a separable, nowhere scattered C^* -algebra of stable rank one has k -comparison if and only if it has strict comparison. Consequently, we get that $\text{div}_2(x) \leq 4$ for every $x \in \text{Cu}(A)$. Further, a small modification of the proof shows that $\text{Cu}(A)$ is almost divisible (ie. 0-almost divisible).

The following is a generalization of [TV21b, Proposition 4.2].

Proposition 4.5. *Let I be an ideal of a C^* -algebra A . Assume that there exist $m_0, m_1, n_1, n_2 \in \mathbb{N}$ such that*

$$\text{div}_{2m_0}(m_0[a]) \leq n_0, \quad \text{and} \quad \text{div}_{2m_1}(m_1[b]) \leq n_1$$

for every $a \in (A/I)_+$ and every $b \in I_+$. Then,

$$\text{div}_{2m_0m_1}((m_0m_1)[c]) \leq m_0(m_1n_0 + n_1)$$

for every $c \in A_+$.

Conversely, $\text{div}_{2m}(m[a]), \text{div}_{2m}(m[b]) \leq n$ for every $a \in (A/I)_+$ and $b \in I_+$ whenever $\sup_{c \in A_+} \text{div}_{2m}(m[c]) \leq n$ for some $m, n \in \mathbb{N}$.

Proof. It follows from [CRS10] that $\text{Cu}(I)$ can be identified with an ideal of $\text{Cu}(A)$, and that $\text{Cu}(A/I)$ is naturally isomorphic to the quotient $\text{Cu}(A)/\text{Cu}(I)$. We will make these identifications without explicitly writing the isomorphisms.

Let $c \in A_+$, and set $x := [c]$. Denote by $\text{Cu}(\pi): \text{Cu}(A) \rightarrow \text{Cu}(A/I)$ the induced quotient map. Given x', x'' in $\text{Cu}(A)$ such that $x' \ll x'' \ll x$, we can apply the bounded divisibility of $\text{Cu}(A/I)$ to obtain elements $y_j'', y_j', y_j \in \text{Cu}(A)$ for $j = 1, \dots, n_0$ such that

$$(2m_0) \text{Cu}(\pi)(y_j) \leq \text{Cu}(\pi)(m_0 x), \quad \text{Cu}(\pi)(m_0 x'') \leq \text{Cu}(\pi)(y_1'') + \dots + \text{Cu}(\pi)(y_{n_0}''),$$

and $y_j'' \ll y_j' \ll y_j$ for each j . Note that this is possible because $\text{Cu}(\pi)(x) = [\pi(c)]$ and $\text{Cu}(\pi)(x)$ has a representative in $(A/I)_+$.

Take $w \in \text{Cu}(I)$ such that $w = 2w$, and

$$(2m_0)y_j' \ll (2m_0)y_j \leq m_0 x + w, \quad \text{and} \quad m_0 x'' \leq y_1'' + \dots + y_{n_0}'' + w.$$

Using [TV21b, Proposition 7.8], we find elements $z_j', z_j \in \text{Cu}(A)$ satisfying

$$(2m_0)z_j \ll m_0 x, \quad y_j'' \ll z_j' + w, \quad \text{and} \quad z_j' \ll z_j$$

for every $j = 1, \dots, n_0$.

In particular, we have

$$x' \ll x'' \leq m_0 x'' \leq z_1' + \dots + z_{n_0}' + w.$$

Applying (O6), we obtain an element $r \in \text{Cu}(A)$ such that

$$x' \ll z_1' + \dots + z_{n_0}' + r, \quad \text{and} \quad r \leq x'', w.$$

Note that, since $r \leq x''$, it follows that $r = [b]$ for some $b \in A_+$. Further, since we also have $[b] \leq w$, one gets $b \in I_+$. Take $r' \ll r$ such that $x' \leq z_1' + \dots + z_{n_0}' + r'$.

Now, using that $\text{div}_{2m_1}(m_1[b]) \leq n_1$, one gets elements r_1, \dots, r_{n_1} such that

$$(2m_1)r_i \ll m_1 r, \quad \text{and} \quad m_1 r' \ll r_1 + \dots + r_{n_1}$$

for each $i \leq n_1$.

Thus, it follows that

$$(m_0 m_1)x' \leq (m_0 m_1)z_1' + \dots + (m_0 m_1)z_{n_0}' + m_0 r_1 + \dots + m_0 r_{n_1}$$

and

$$(2m_0 m_1)z_j' \leq (m_0 m_1)x, \quad (2m_0 m_1)r_i \leq (m_0 m_1)r \leq (m_0 m_1)x$$

for each i and j .

This proves that

$$\text{div}_{2m_0 m_1}((m_0 m_1)x) \leq m_0(m_1 n_0 + n_1),$$

as desired.

The converse follows from a standard argument; see [TV22b, Proposition 3.9]. \square

The previous result also works, mutatis-mutandi, with $\text{Div}()$ instead of $\text{div}()$. In this case, an inspection of [TV22b, Theorem 3.10] gives the following:

Theorem 4.6. *Let I be an ideal of a C^* -algebra A . Assume that there exist $m_0, m_1, n_0, n_1 \in \mathbb{N}$ such that*

$$\text{Div}_{2m_0}(m_0[a]) \leq n_0, \quad \text{and} \quad \text{Div}_{2m_1}(m_1[b]) \leq n_1$$

for every $a \in I_+$ and every $b \in (A/I)_+$. Then, there exist $M, N \in \mathbb{N}$ depending only on m_0, m_1, n_0, n_1 such that

$$\text{Div}_{2M}(M[c]) \leq N$$

for every $c \in A_+$.

Conversely, $\text{Div}_{2m}(m[a]), \text{Div}_{2m}(m[b]) \leq n$ for every $a \in (A/I)_+$ and $b \in I_+$ whenever $\sup_{c \in A_+} \text{Div}_{2m}(m[c]) \leq n$ for some $m, n \in \mathbb{N}$.

As noted in Lemma 3.11, having a sup-dense subset of elements with finite divisibility does not imply that every element has such property. However, Proposition 4.8 below shows that it is enough to check finite divisibility for strongly soft elements (which, in general, are not sup-dense).

4.7 (Strongly soft elements and retracts). Let A be a C^* -algebra. As defined in [TV23], we denote by $\text{Cu}(A)_{\text{soft}}$ the set of *strongly soft elements*, that is, those elements $x \in \text{Cu}(A)$ such that for every $x' \ll x$ there exists $t \in \text{Cu}(A)$ satisfying $x' + t \leq x \leq \infty t$. When A is residually stably finite, $[a] \in \text{Cu}(A)$ is strongly soft if and only if \overline{aAa} has no nonzero, unital quotients; see [TV23, Proposition 4.16].

Under certain assumptions, given any $x \in \text{Cu}(A)$ one can find the largest strongly soft element below x . When this is the case, we denote by $\sigma: \text{Cu}(A) \rightarrow \text{Cu}(A)_{\text{soft}}$ the map that sends an element x to the largest strongly soft element dominated by x .

As shown in [Thi20, Proposition 2.9], this map can be defined whenever A is separable, simple, and of stable rank one. In fact, one can show that σ is an order- and suprema-preserving monoid morphism.

In [AVTV23, Theorem 5.6], it is shown that σ can always be defined whenever A is separable, has the Global Glimm Property, and its Cuntz semigroup is *left-soft separative*, that is, if for any triplet of elements $y, t \in S$ and $x \in S_{\text{soft}}$ satisfying

$$x + t \ll y + t, \quad t \ll \infty y, \quad \text{and} \quad t \ll \infty x,$$

we have $x \ll y$; see [AVTV23, Definition 3.2]. As explained in [AVTV23, Section 3], C^* -algebras with strict comparison or stable rank one have a left-soft separative Cuntz semigroup.

In this case, σ is an order- and suprema-preserving superadditive map that satisfies $x \leq \sigma(x) + t$ whenever $x \leq \infty t$.

Proposition 4.8. *Let A be a separable C^* -algebra with the Global Glimm Property such that $\text{Cu}(A)$ is left-soft separative, and let $x \in \text{Cu}(A)$ and $m \in \mathbb{N}$. Then*

$$\text{div}_m(\sigma(x)) - 1 \leq \text{div}_m(x) \leq \text{div}_m(\sigma(x)) + 1.$$

Proof. First, to see that $\text{div}_m(\sigma(x)) \leq \text{div}_m(x) + 1$, assume that $\text{div}_m(x) = n$ for some $n \in \mathbb{N}$, since otherwise there is nothing to prove.

Let $s \ll \sigma(x)$. Using that σ preserves suprema of increasing sequences, we can find $x' \ll x$ such that $s \ll \sigma(x')$.

By [TV23, Proposition 7.7], there exists $t \in \text{Cu}(A)_{\text{soft}}$ such that $(nm)t \leq x \leq \infty t$. Note that $(nm)t \leq \sigma(x)$ because t is strongly soft. Using that x is weakly (m, n) -divisible, one finds $z_1, \dots, z_n \in \text{Cu}(A)$ such that

$$\sigma(x') \leq z_1 + \dots + z_n, \quad \text{and} \quad mz_j \leq x$$

for every j .

Now, using that $z_i \leq \infty t$ for every i , we obtain

$$\sigma(x') \leq z_1 + \dots + z_n \leq (\sigma(z_1) + t) + z_2 + \dots + z_n \leq \sigma(z_1) + \dots + \sigma(z_n) + nt.$$

Further, using that σ is superadditive, we also get $m\sigma(z_j) \leq \sigma(mz_j) \leq \sigma(x)$. Since $n(mt) \leq \sigma(x)$, we deduce that $\text{div}_n(\sigma(x)) \leq n + 1$, as desired.

To prove that $\text{div}_m(x) \leq \text{div}_m(\sigma(x)) + 1$, take $t \in \text{Cu}(A)$ such that $mt \leq x \leq \infty t$. Note that this can be done because A has the Global Glimm Property and, consequently, $\text{Cu}(A)$ is (m, ω) -divisible; see Paragraph 2.3. Thus, we have $x \leq \sigma(x) + t$. Let $x' \ll x$. Since $x' \ll \sigma(x) + t$, we can use (O6) to obtain an element y such that $x' \ll y + t$ with $y \ll \sigma(x)$.

Let n be such that $\sigma(x)$ is weakly (m, n) -divisible. Then, there exist z_1, \dots, z_n satisfying

$$y \leq z_1 + \dots + z_n, \quad \text{and} \quad mz_j \leq \sigma(x)$$

for each j .

We obtain

$$x' \leq z_1 + \dots + z_n + t, \quad mt \leq x, \quad \text{and} \quad mz_j \leq \sigma(x) \leq x$$

for every j , which implies that $\text{div}_m(x) \leq n + 1$. \square

Lemma 4.9. *Let A be a C^* -algebra, and let $s, t \in \text{Cu}(A)$. Assume that s is strongly soft. Then, the element $s \wedge \infty t$ is strongly soft in $\text{Cu}(A)$.*

Proof. Let $x', x \in \text{Cu}(A)$ be such that $x' \ll x \ll s \wedge \infty t$. By definition, this implies that $x' \ll s$ and $x' \ll \infty t$.

Since s is strongly soft, we find $y \in \text{Cu}(A)$ such that $x' + y \leq s \leq \infty y$. Thus, using that taking infima with an idempotent is a monoid morphism ([APRT21, Section 2]), we have

$$x' + y \wedge \infty t = (x' + y) \wedge \infty t \leq s \wedge \infty t \leq \infty y \wedge \infty t = \infty(y \wedge \infty t).$$

This shows that $s \wedge \infty t$ is strongly soft. \square

When $\text{Cu}(A)$ is not left-soft separative, we still have the following:

Theorem 4.10. *Let A be a C^* -algebra with the Global Glimm Property. Then, the following are equivalent:*

- (i) *there exist $n, m \in \mathbb{N}$ such that $\text{div}_{2m}(m[a]) \leq n$ for every $[a] \in \text{Cu}(A)$;*
- (ii) *there exist $n, m \in \mathbb{N}$ such that $\text{div}_{2m}(m[a]) \leq n$ for every $[a] \in \text{Cu}(A)_{\text{soft}}$.*

Proof. That (i) implies (ii) is trivial.

To prove the converse, let $x', x \in \text{Cu}(A)$ be such that $x' \ll x$. It follows from [TV23, Proposition 7.7] that there exists $y \in \text{Cu}(A)_{\text{soft}}$ satisfying $y \leq x \leq \infty y$. In particular, one has $x' \ll \infty y$.

By [TV23, Proposition 5.6], we can take $y', y'' \in \text{Cu}(A)_{\text{soft}}$ such that $y' \ll y'' \ll y$ and $x' \ll \infty y'$. Since y'' is soft, we find $t \in \text{Cu}(A)_{\text{soft}}$ such that $y' + t \leq y'' \leq \infty t$; see [TV23, Proposition 4.13]. Now, by (O5), there exists $c \in \text{Cu}(A)$ satisfying

$$y'' + c \leq x \leq y + c.$$

Thus, we have

$$y' + (c + t) \leq x \leq y + c \leq y + (c + t).$$

Using that $x' \leq \infty y'$, we obtain

$$x' \leq x \wedge \infty y'' \leq (y + (c + t)) \wedge \infty y'' = y \wedge \infty y'' + (c + t) \wedge \infty y''$$

and

$$y \wedge \infty y'', (c + t) \wedge \infty y'' \leq x.$$

By Lemma 4.9, the elements $y \wedge \infty y''$ and $t \wedge \infty y''$ are strongly soft. Further, note that one gets

$$(c + t) \wedge \infty y'' = c \wedge \infty y'' + t \wedge \infty y''$$

with $c \wedge \infty y'' \leq \infty y'' \leq \infty(t \wedge \infty y'')$.

Using [TV23, Theorem 4.16], we deduce that $(c + t) \wedge \infty y''$ is also strongly soft. In other words, for every pair of elements $x', x \in \text{Cu}(A)$ such that $x' \ll x$, we find $s_1, s_2 \in \text{Cu}(A)_{\text{soft}}$ such that

$$s_1, s_2 \leq x, \quad \text{and} \quad x' \leq s_1 + s_2.$$

In particular, $ms_1, ms_2 \leq mx$ and $mx' \leq ms_1 + ms_2$.

By Lemma 3.3 (ii), we have

$$\text{div}_{2m}(mx) \leq \text{div}_{2m}(ms_1) + \text{div}_{2m}(ms_2) \leq 2n,$$

as desired. \square

Corollary 4.11. *Let A be a C^* -algebra with the Global Glimm Property. Assume that $\text{Cu}(A)_{\text{soft}}$ is a retract of a Cu -semigroup S satisfying $\sup_{s \in S} \text{div}_{2k}(ks) < \infty$ for some $k \in \mathbb{N}$. Then, there exist $n, m \in \mathbb{N}$ such that $\text{div}_{2m}(m[a]) \leq n$ for every $[a] \in \text{Cu}(A)$.*

Proof. Using that $\text{Cu}(A)_{\text{soft}}$ is a retract of S , it follows that

$$\text{div}_{2k}(ky) \leq \sup_{s \in S} \text{div}_{2k}(ks) < \infty$$

for every $y \in \text{Cu}(A)_{\text{soft}}$.

Applying Theorem 4.10, there exist $n, m \in \mathbb{N}$ such that $\text{div}_{2m}(mx) \leq n$ for every $x \in \text{Cu}(A)$. \square

Recall from [TV22a, Definition 3.1] that one says that $\text{Cu}(A)$ has *covering dimension zero* if, whenever $x' \ll x \ll y_1 + y_2$, there exist $z_1, z_2 \in \text{Cu}(A)$ such that $x' \leq z_1 + z_2 \leq x$. Examples of dimension zero include the Cuntz semigroup of any real rank zero C^* -algebra, as well as other semigroups, such as the Cuntz semigroup of the Jacelon-Razak algebra; see [TV22a, Section 5].

We say that a Cuntz semigroup is *weakly cancellative* if $x \ll y$ whenever $x + z \ll y + z$ for some element z . Stable rank one C^* -algebras have a weakly cancellative Cuntz semigroup by [RW10, Theorem 4.3].

Corollary 4.12. *Let A be a simple, non-elementary C^* -algebra of Cuntz covering dimension zero. Assume that $\text{Cu}(A)$ is weakly cancellative. Then there exists $n \in \mathbb{N}$ such that $\text{div}_2([a]) \leq n$ for every $[a] \in \text{Cu}(A)$.*

Proof. Assume first that A is separable. [TV22a, Theorem 7.10] shows that $\text{Cu}(A)_{\text{soft}}$ is a retract of an almost divisible Cu -semigroup S ; see also the proof of [TV22a, Proposition 7.13].

As explained in Example 3.10, almost divisibility implies $\text{div}_2(s) \leq 3$ for every element $s \in S$. Thus, the result in this case follows from Corollary 4.11 above.

If A is not separable, one can use the same techniques as in the proof of Proposition 4.3 to get the desired result. \square

5. NOWHERE SCATTERED CORONA AND MULTIPLIER ALGEBRAS

In this section we study when a multiplier algebra of a σ -unital, nowhere scattered C^* -algebra A is nowhere scattered. Our main result is that this happens whenever $\sup_{a \in A_+} \text{div}_{2m}(m[a]) < \infty$ for some $m \in \mathbb{N}$; see Theorem 5.11. Note that the supremum is not taken over all the elements of the Cuntz semigroup (ie. over $A \otimes K$), but only over those with a representative in A_+ . This set is called the *scale* of the Cuntz semigroup in [APT20, 4.2].

As an application, we prove in Theorem 5.12 that nowhere scattered C^* -algebras of finite nuclear dimension, or real rank zero, or stable rank one and k -comparison all have a nowhere scattered multiplier algebra. We also show in Theorem 5.18

that, for stable C^* -algebras, the multiplier algebra being nowhere scattered implies that every element in the algebra has finite divisibility.

Let us begin the section with some examples:

Example 5.1. Let A be a weakly purely infinite C^* -algebra. It was shown in [KR02, Proposition 4.11] that $\mathcal{M}(A)$ is weakly purely infinite, and thus nowhere scattered by [TV21b, Example 3.3].

As a similar example, it follows from [Lin04] that a simple, σ -unital, non-elementary (ie. nowhere scattered) C^* -algebra with continuous scale has a purely infinite corona algebra. This implies that $\mathcal{M}(A)$ is nowhere scattered by [TV21b, Proposition 4.2]; see also [KNZ19] and [Lin91].

The examples below appeared in [TV21b, Examples 4.14, 4.15]. We recall them here for the convenience of the reader.

Example 5.2. Let A_k be the family of separable, simple, AH-algebras from [RR13] (see Example 3.6). As shown in [RR13, Corollary 8.6], the product $\prod_k A_k$ has a one-dimensional, irreducible representation. By [TV21b, Theorem 3.1], this implies that $\prod_k A_k$ is not nowhere scattered. Consequently, $A := \oplus_k A_k$ is a nowhere scattered C^* -algebra with a multiplier algebra $\mathcal{M}(A) \cong \prod_k A_k$ that is not nowhere scattered.

Example 5.3. In [Sak71, Theorem 1], Sakai constructs a simple C^* -algebra A such that its derived algebra $D(A)$ satisfies $D(A)/A \cong \mathbb{C}$. Using that $D(A) \cong \mathcal{M}(A)$ for simple C^* -algebras (see the remarks after [Ped72, Proposition 2.6]), one sees that $\mathcal{M}(A)$ cannot be nowhere scattered.

It will follow from Theorem 5.11 and Proposition 4.1 that A is of infinite nuclear dimension. [Sak71, Corollary 1] already showed that A is not \mathcal{Z} -stable.

The following lemma will play an important role in the proof of Proposition 5.9.

Lemma 5.4. *Let A be a C^* -algebra. The following are equivalent:*

- (i) A is nowhere scattered.
- (ii) For every $a \in A_+$ and $n \in \mathbb{N}$, the class $[a \otimes 1_n] \in \text{Cu}(A)$ is weakly $(2n, \omega)$ -divisible.
- (iii) For every $a \in A_+$, there exists $n = n(a) \in \mathbb{N}$ such that the class $[a \otimes 1_n] \in \text{Cu}(A)$ is weakly $(n+1, \omega)$ -divisible.

Proof. We know from [TV21b, Theorem 8.9] that A is nowhere scattered if and only if every element in $\text{Cu}(A)$ is weakly $(2, \omega)$ -divisible. Using [TV21b, Theorem 8.9], it follows that every element is $(2n, \omega)$ -divisible for every $n \in \mathbb{N}$. This proves that (i) implies (ii), and it is trivial that (ii) implies (iii).

To see that (iii) implies (i), assume for the sake of contradiction that A is not nowhere scattered. Then, we know from [TV21b, Theorem 3.1] that there exist ideals $I \subseteq J \subseteq A$ such that J/I is elementary, that is, $\text{Cu}(I/J) \cong \text{Cu}(\mathbb{C}) \cong \overline{\mathbb{N}}$; see [Eng14, Theorem 4.4.4].

Note that, since we are assuming that A satisfies (iii), both I and I/J satisfy (iii) as well. Let ϕ be an isomorphism from $\text{Cu}(I/J)$ to $\overline{\mathbb{N}}$. Now, it follows from [TV22b, Lemma 3.3] that for every positive element $b \in (I/J) \otimes \mathcal{K}$ there exists $a \in (I/J)_+$ such that $[b] \leq \infty[a]$ in $\text{Cu}(I/J)$. Thus, let $a \in (I/J)_+$ be such that $\phi([a]) \neq 0$. We get $1 \ll 1 \leq \phi([a])$ and, using Rørdam's lemma (see, for example, [Thi17, Theorem 2.30]), there exists $a' \in (I/J)_+$ such that $1 = \phi([a'])$.

Take $n \in \mathbb{N}$ such that $[a' \otimes 1_n]$ is weakly $(n+1, \omega)$ -divisible. Since ϕ is an isomorphism, we must have that $\phi([a' \otimes 1_n]) = n$ is also weakly $(n+1, \omega)$ -divisible. However, the only element $x \in \overline{\mathbb{N}}$ such that $(n+1)x \leq n$ is zero. This contradicts the weak $(n+1, \omega)$ -divisibility of $[a' \otimes 1_n]$.

Thus, A has no nonzero elementary ideal-quotients, as desired. \square

Remark 5.5. In the proof of ‘(iii) \implies (i)’ in Lemma 5.4 above, we have used that a C^* -algebra is nowhere scattered if and only if it has no elementary ideal-quotients. As shown in [TV21b, Proposition 8.8], a Cu-semigroup S satisfying (O5)-(O8) has no elementary ideal-quotients if and only if every element in S is weakly $(2, \omega)$ -divisible.

In light of this, one might expect ‘(iii) \implies (i)’ to hold for every Cu-semigroup S satisfying (O5)-(O8), that is, that every element in S is weakly $(2, \omega)$ -divisible whenever for every element $x \in S$ there exists n_x such that $n_x x$ is weakly $(n_x + 1, \omega)$ -divisible.

However, this is not true: For example, $S = \{0, 1, \infty\}$ is a Cu-semigroup satisfying (O5)-(O8). S is not weakly $(2, \omega)$ -divisible, but every element in S has a properly infinite multiple. Thus, it satisfies (iii), but not (i).

The reason for this disparity is due to the fact that, in the context of abstract Cuntz semigroups, $\bar{\mathbb{N}}$ is not the only elementary Cu-semigroup. In fact, $\{0, 1, \infty\}$ is elementary; see [TV21b, Section 8].

Lemma 5.6 below is [KR02, Lemma 4.10], which uses some of the ideas from [Ell74]. It was stated with the assumption of weak pure infiniteness, but an inspection of their proof shows that this is not actually needed. This was also stated, in a different way and with a different proof, in [KNZ17, Theorem 4.2].

Lemma 5.6. *Let A be a σ -unital C^* -algebra, let $T \in \mathcal{M}(A)_+$, and let $\varepsilon > 0$. Then A has an increasing, countable, approximate unit $(e_n)_n$ of positive contractions with $e_{n+1}e_n = e_n$ such that*

$$T = a + \sum_{n=1}^{\infty} f_{2n-1}^{1/2} T f_{2n-1}^{1/2} + \sum_{n=1}^{\infty} f_{2n}^{1/2} T f_{2n}^{1/2}, \quad f_n := e_n - e_{n-1}$$

with $e_0 = 0$ and $a \in A$ satisfying $\|a\| \leq \varepsilon$.

In the proof of Proposition 5.9 we will also need the following:

Lemma 5.7. *Let A be a C^* -algebra, and let $a, b \in A_+$ be elements such that $\text{mdiv}([a]), \text{mdiv}([b]) < \infty$. Then $\text{mdiv}([a + b]) < \infty$.*

In particular, there exists $n \in \mathbb{N}$ such that $(a + b) \otimes 1_n$ has a $(n + 1, \omega)$ -divisible class.

Proof. Let $n_a, n_b \in \mathbb{N}$ be such that $\text{div}_{2n_a}(n_a[a]), \text{div}_{2n_b}(n_b[b]) < \infty$.

Using that $n_a n_b [a + b] = [a \otimes 1_{n_a n_b} + b \otimes 1_{n_a n_b}]$, we can apply Lemma 3.3 (iii) and get

$$\text{div}_{2n_a n_b}(n_a n_b [a + b]) \leq \text{div}_{2n_a n_b}(n_a n_b [a]) + \text{div}_{2n_a n_b}(n_a n_b [b]).$$

Thus, applying Lemma 3.4 to both summands, we obtain

$$\text{div}_{2n_a n_b}(n_a n_b [a + b]) \leq \text{div}_{2n_a}(n_a [a]) + \text{div}_{2n_b}(n_b [b]) < \infty,$$

as desired. \square

Lemma 5.8. *Let A be a σ -unital C^* -algebra, and let $(a_i)_{i=1}^{\infty}$ be a bounded sequence in A of pairwise orthogonal elements such that $R = \sum_i a_i$ is strictly convergent and $a_i \perp e_{i-1}$ for every $i \geq 1$, where $(e_i)_i$ is an approximate unit as in Lemma 5.6.*

Let $m \in \mathbb{N}$. Then for every $\varepsilon_0 > 0$, one has

$$\text{div}_{2m}(m[R]) \leq \sup_i \sup_{0 \leq \varepsilon \leq \varepsilon_0} \text{div}_{2m}(m[(a_i - \varepsilon)_+])$$

in $\text{Cu}(\mathcal{M}(A))$.

Proof. Let $\varepsilon_0 > 0$. Assume that $n := \sup_i \sup_{\varepsilon \leq \varepsilon_0} \text{div}_{2m}(m[(a_i - \varepsilon)_+])$ is finite, since otherwise there is nothing to prove. Fix $\varepsilon \leq \varepsilon_0$ positive. By assumption, $(a_i - \frac{\varepsilon}{4})_+ \otimes 1_m$ is weakly $(2m, n)$ -divisible for each i . Thus, we can find elements $b_{i,j} \in A \otimes \mathcal{K}$ for $j = 1, \dots, n$ such that

$$m[(a_i - \varepsilon/3)_+] \leq [b_{i,1}] + \dots + [b_{i,n}], \quad \text{and} \quad 2m[b_{i,j}] \leq m[(a_i - \varepsilon/4)_+]$$

for each i and j .

It follows from [RR13, Lemma 2.3(i)] that there exist elements $c_{i,j} \in M_m(A)_+$ with $c_{i,j} \preceq b_{i,j}$ such that

$$(a_i - 2\varepsilon/3)_+ \otimes 1_m = \sum_{j=1}^n c_{i,j}.$$

Set $d_{i,j} := ((a_i - 2\varepsilon/3)_+ \otimes 1_m)c_{i,j}((a_i - 2\varepsilon/3)_+ \otimes 1_m) \in M_m(A)$. Note that every entry in $d_{i,j}$ is in $\overline{a_i A a_i}$, and is thus orthogonal with e_{i-1} . Further, since the sequence $(d_{i,j})_i$ is bounded for every fixed j , so is each of the induced entry-wise sequences. This shows that the (strictly convergent) sum $R_j := \sum_i d_{i,j}$ is in $M_m(\mathcal{M}(A))$.

Thus, we have

$$(R - 2\varepsilon/3)_+^3 \otimes 1_m = \sum_{j=1}^n R_j$$

and, therefore, $m[(R - \varepsilon)_+] \ll \sum_{j=1}^n [R_j]$.

Let $\delta > 0$ be such that $m[(R - \varepsilon)_+] \leq \sum_{j=1}^n [(R_j - \delta)_+]$. Applying [KNZ17, Lemma 2.2] at

$$d_{i,j} \otimes 1_{2m} \preceq c_{i,j} \otimes 1_{2m} \preceq b_{i,j} \otimes 1_{2m} \preceq (a_i - \varepsilon/4)_+ \otimes 1_m$$

we find elements $r_{i,j}$ such that

$$(d_{i,j} - \delta)_+ \otimes 1_{2m} = r_{i,j}(a_i \otimes 1_m)r_{i,j}^*, \quad \text{and} \quad \sup_i \|r_{i,j}\|^2 \leq \frac{4}{\varepsilon} \sup_i \|d_{i,j} \otimes 1_{2m}\|.$$

Set $x_{i,j} := (a_i \otimes 1_m)^{1/2} r_{i,j}^* (d_{i,j} \otimes 1_{2m} - \delta)_+$. We get

$$x_{i,j}^* x_{i,j} = (d_{i,j} - \delta)_+^3 \otimes 1_{2m}, \quad \text{and} \quad x_{i,j} x_{i,j}^* \in \overline{(a_i \otimes 1_m) M_m(A) (a_i \otimes 1_m)},$$

with both $((a_i \otimes 1_m)^{1/2} r_{i,j}^*)_i$ and $(r_{i,j}^* (d_{i,j} \otimes 1_{2m} - \delta)_+)_i$ being bounded sequences for each fixed j .

Let $x_{i,j}(s, r)$ denote the (s, r) -th entry of $x_{i,j}$. Working as in [KNZ17, Proposition 4.4], one can now show that the following sums are all strictly convergent

$$x_j(s, r) := \sum_i x_{i,j}(s, r).$$

Set $x_j := (x_j(s, r))_{s,r} \in M_{m, 2m^2}(\mathcal{M}(A))$. One gets

$$x_j^* x_j = (R_j - \delta)_+^3 \otimes 1_{2m} \sim (R_j - \delta)_+ \otimes 1_{2m},$$

and

$$x_j x_j^* \in \overline{(R \otimes 1_m) M_m(\mathcal{M}(A)) (R \otimes 1_m)}.$$

It follows that, in $\text{Cu}(\mathcal{M}(A))$, we have $2m[(R_j - \delta)_+] \leq m[R]$ for each $j \in \mathbb{N}$. Since $[R] = \sup_{0 \leq \varepsilon \leq \varepsilon_0} [(R - \varepsilon)_+]$, we have that $m[R]$ is weakly $(2m, n)$ -divisible, as desired. \square

Let us say that a sequence $(x_i)_i$ in a C^* -algebra A is *orthogonal* if the elements in the sequence are pairwise orthogonal, that is, if $x_i x_j = 0$ whenever $i \leq j$.

Proposition 5.9. *Let A be a σ -unital C^* -algebra. Assume that for every orthogonal sequence $(a_i)_i$ of positive elements in A there exist $m, n \in \mathbb{N}$ such that $\text{div}_{2m}(m[a_i]) \leq n$ for each i . Then $\text{div}_{2m}(m[T]) < \infty$ for every $T \in \mathcal{M}(A)/A$.*

In particular, $\mathcal{M}(A)/A$ is nowhere scattered.

Proof. Let $\pi: \mathcal{M}(A) \rightarrow \mathcal{M}(A)/A$ denote the quotient map. Using Lemmas 5.6 and 5.8, we deduce that each positive element in the corona algebra can be written as $\pi(R_1) + \pi(R_2)$ with $\text{mdiv}(R_1), \text{mdiv}(R_2) < \infty$.

Using Lemma 5.7, one has $\text{mdiv}(\pi(T)) < \infty$ in $\mathcal{M}(A)/A$ for every element $T \in \mathcal{M}(A)$. Lemma 5.4 shows that the corona algebra is nowhere scattered, as desired. \square

Remark 5.10. Let A be the simple, nowhere scattered C^* -algebra from Example 5.3. It follows from Proposition 5.9 above that there must exist an element $a \in (\oplus_{i=1}^{\infty} A)_+$ such that $\text{mdiv}([a]) = \infty$.

Using Proposition 5.9, we can now prove the main result of the paper.

Theorem 5.11. *Let A be a σ -unital C^* -algebra. Assume that for every orthogonal sequence $(a_i)_i$ of positive elements in A there exist $m, n \in \mathbb{N}$ such that $\text{div}_{2m}(m[a_i]) \leq n$ for each i . Then $\mathcal{M}(A)$ is nowhere scattered.*

Proof. Using that nowhere scatteredness works well with extensions ([TV21b, Proposition 4.2]), the result is a direct consequence of Proposition 5.9. \square

It follows from the results in [PR04] that the multiplier algebra of a simple, separable C^* -algebra of real rank zero does not admit a character. Theorem 5.12 (i) below generalizes this result.

Theorem 5.12. *Let A be a σ -unital C^* -algebra. Assume that A has*

- (i) *real rank zero, or*
- (ii) *finite nuclear dimension, or*
- (iii) *k -comparison and a surjective rank map.*

Then A is nowhere scattered if and only if $\mathcal{M}(A)$ is nowhere scattered.

Proof. Assume that $\mathcal{M}(A)$ is nowhere scattered. [TV21b, Proposition 4.2] implies that A is nowhere scattered as well.

Conversely, assume that A is nowhere scattered. It follows from Example 3.10, Proposition 4.1 and Proposition 4.3 respectively that (i), (ii) and (iii) imply that every sequence of positive elements (orthogonal or not) in A_+ has uniformly bounded multiple divisibility. Thus, we can use Theorem 5.11 to get the desired result. \square

Question 5.13. Let A be a nowhere scattered C^* -algebra such that $\text{mdiv}([a]) < \infty$ for every $a \in A_+$. Does it follow that $\mathcal{M}(A)$ is nowhere scattered?

Note that this would imply that the simple C^* -algebra from Example 5.3 contains a $(2, \omega)$ -divisible element of infinite divisibility.

Remark 5.14. The assumption from Theorem 5.11 above is not equivalent to nowhere scatteredness, not even in the simple or σ -unital case:

- (1) The example from Example 5.2 is σ -unital. Thus, there exist σ -unital, nowhere scattered C^* -algebras that do not satisfy the condition from Theorem 5.11.
- (2) By Lemma 3.11, the set of classes with finite divisibility is sup-dense in the Cuntz semigroup of a simple, non-elementary C^* -algebra A . However, and as made explicit in Example 5.3, this is not enough for the multiplier algebra $\mathcal{M}(A)$ to be nowhere scattered.

Further, note that if the converse of Theorem 5.11 holds for some family of C^* -algebras, one must have that any unital, nowhere scattered C^* -algebra A in the family must satisfy $\sup_{a \in A_+} \text{div}_{2m}(m[a_i]) < \infty$ for some m .

As another application of Lemma 5.8, we can study when multiplier algebras have a character:

Proposition 5.15. *Let A be a σ -unital C^* -algebra, and let $(e_i)_i$ be an approximate unit as in Lemma 5.6. Assume that*

$$\sup_i \sup_{0 \leq \varepsilon \leq \varepsilon_0} \text{div}_{2m}(m[(e_i - e_{i-1} - \varepsilon)_+]) < \infty$$

for some $\varepsilon_0 > 0$.

Then, $\mathcal{M}(A)$ does not admit a character.

Proof. Using Lemma 5.8, we see that the unit $1 = \sum_i (e_i - e_{i-1})$ in $\mathcal{M}(A)$ satisfies $\text{div}_{2m}(m[1]) < \infty$.

The result now follows from [RR13, Corollary 5.6]. \square

In particular, Proposition 5.15 recovers [RR13, Corollary 8.5]:

Corollary 5.16. *The product of unital C^* -algebras A_i does not admit a character whenever the supremum of $\text{div}_2([1_i])$ is finite.*

5.1. Stable C^* -algebras. The multiplier algebra of a σ -unital, stable C^* -algebra A has been studied extensively. For example, [Rør91] studies when the corona algebra of A is simple, while in [NW10] it is shown, using its multiplier algebra, that a nowhere scattered C^* -algebra of finite nuclear dimension has the corona factorization property.

In what follows, we show that $\text{mdiv}([a])$ must be finite for every $a \in A_+$ if $\mathcal{M}(A)$ is to be nowhere scattered; see Theorem 5.18. Thus, if Question 5.13 has an affirmative answer, this would imply that the multiplier algebra of a σ -unital, stable C^* -algebra is nowhere scattered if and only if every element in the algebra has a multiple of finite divisibility.

Lemma 5.17 below was stated for (m, ω) -divisibility in [TV22b]. We now provide the analog statement for weak (m, ω) -divisibility, which only needs to assume the element (and not the whole Cuntz semigroup) to be divisible.

Lemma 5.17. *Let A be a C^* -algebra, and let $a \in (A \otimes \mathcal{K})_+$. Then, if $[a]$ is weakly (m, ω) -divisible, there exists a sequence $(y_n)_n \subseteq \text{Cu}(A)$ such that $my_j \leq [a] \leq \sum_{j=1}^{\infty} y_j$ for every $j \in \mathbb{N}$.*

Proof. Let ε_n be a strictly decreasing sequence converging to 0, and let $x_n := [(a - \varepsilon_n)_+]$. We will inductively find elements $y_{k,j}, c_n$ such that

$$x_n \leq \sum_{k=1}^n \sum_j y_{k,j}, \quad \text{and} \quad my_{k,j} \leq [a].$$

First, use weak (m, ω) -divisibility to obtain $y_{1,j}$ such that $x_1 \leq \sum_j y_{1,j}$ and $my_{1,j} \leq [a]$.

Assume now that we have proven the result for every $k \leq n$. Using (O5) at $x_{n-1} \ll x_n \leq [a]$, we obtain $c \in \text{Cu}(A)$ such that $x_{n-1} + c \leq [a] \leq x_n + c$. In particular, one has $x_{n+1} \ll [a] \leq x_n + c$. Take $c' \ll c$ such that $x_{n+1} \leq x_n + c'$.

Using (m, ω) -divisibility at $c' \ll c \leq x$, we obtain elements $y_{n+1,j}$ such that $c' \leq \sum_j y_{n+1,j}$ and $my_{n+1,j} \leq x$ for each j . We have

$$x_{n+1} \leq x_n + c' \leq \sum_{k=1}^n \sum_j y_{k,j} + \sum_j y_{n+1,j} = \sum_{k=1}^{n+1} \sum_j y_{k,j},$$

as desired.

By taking the supremum on n , one sees that the sequence $(y_{n,j})_{n,j}$ satisfies the desired conditions. \square

Note that, in general, one cannot adapt the previous proof to show that there exist y_1, \dots, y_m with $2y_j \leq [a] \leq y_1 + \dots + y_m$ whenever $[a]$ is $(2, m)$ -divisible.

However, this is the case for σ -unital C^* -algebras whose multiplier algebra is nowhere scattered:

Theorem 5.18. *Let A be a σ -unital, stable C^* -algebra. Assume that $\mathcal{M}(A)$ is nowhere scattered. Then, for every $a \in A_+$ and $m \in \mathbb{N}$ there exist finitely many elements $y_1, \dots, y_n \in \text{Cu}(A)$ such that*

$$my_j \leq [a] \leq y_1 + \dots + y_n$$

for every $j \leq n$.

In particular, $\text{mdiv}(a) < \infty$ for every $a \in A_+$.

Proof. Let $a \in A_+$ and $m \in \mathbb{N}$. By [APRT22, Proposition 2.8] there exists a projection $p_a \in \mathcal{M}(A)$ such that, for every $x \in \text{Cu}(A)$, one has $x \leq [a]$ if and only if $x \leq [p_a]$ in $\text{Cu}(\mathcal{M}(A))$.

Assume that $\mathcal{M}(A)$ is nowhere scattered. Thus, $\text{Cu}(\mathcal{M}(A))$ is weakly (m, ω) -divisible by [TV21b, Theorem 8.9]. It follows from Remark 3.2 that $\text{div}_m([p]) < \infty$ for every projection $p \in \mathcal{M}(A)$.

Then, since we have $[a] \leq [p_a] \ll [p_a]$, there exist elements $z_1, \dots, z_n \in \text{Cu}(\mathcal{M}(A))$ such that

$$[a] \leq [p_a] \leq z_1 + \dots + z_n, \quad \text{and} \quad mz_j \leq [p_a]$$

for each $j \leq n$.

Since $\mathcal{M}(A)$ is a C^* -algebra, we know from [APRT21, Remark 2.6] that the infimum $y_j := z_j \wedge \infty[a]$ exists for each j . Any representative of this element is contained in the ideal generated by a and, therefore, must be a positive element in A . Thus, we have $[a] \leq y_1 + \dots + y_n$ in $\text{Cu}(\mathcal{M}(A))$ with $y_j \in \text{Cu}(A)$ for each j .

Further, we also get $my_j = m(z_j \wedge \infty[a]) \leq [p_a]$. Thus, $my_j \leq [a]$ in $\text{Cu}(\mathcal{M}(A))$.

Finally, note that, since A is an ideal of $\mathcal{M}(A)$, a pair of elements in A are Cuntz subequivalent in $\mathcal{M}(A)$ if and only if they are Cuntz subequivalent in A ; see, for example, [Thi17, Proposition 2.18].

Consequently, we have

$$[a] \leq y_1 + \dots + y_n, \quad \text{and} \quad my_j \leq [a]$$

in $\text{Cu}(A)$, as desired. \square

Remark 5.19. Note that there exist stable, nowhere scattered C^* -algebras with a multiplier algebra that is not nowhere scattered. Indeed, simply take A as in Example 3.6. Then, $A \otimes \mathcal{K}$ is a nowhere scattered C^* -algebra by [TV21b, Proposition 4.12] that contains a $(2, \omega)$ -divisible element of infinite divisibility.

It follows from Theorem 5.18 that $\mathcal{M}(A \otimes \mathcal{K})$ cannot be nowhere scattered.

In this paper we have studied conditions under which a multiplier algebra $\mathcal{M}(A)$ is nowhere scattered. A problem that seems to be much more involved is to determine when $\mathcal{M}(A)$ has the Global Glimm Property.

Question 5.20. Assume that A has the Global Glimm Property. When does $\mathcal{M}(A)$ also have the Global Glimm Property?

Note that this has a positive solution whenever $\sup_{a \in A_+} \text{Div}_{2m}(m[a]) < \infty$ and $\mathcal{M}(A)/A$ satisfies a condition that ensures that the Global Glimm Problem can be answered affirmatively. For example, Lin and Ng show in [LN16] that $\mathcal{M}(\mathcal{Z} \otimes$

$\mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ has real rank zero. Thus, since $\mathcal{Z} \otimes \mathcal{K}$ is covered by Theorem 5.11, we get that $\mathcal{M}(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ is nowhere scattered. By [ER06] (see also [TV22b, Proposition 7.4]), $\mathcal{M}(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ has the Global Glimm Property. Consequently, $\mathcal{M}(\mathcal{Z} \otimes \mathcal{K})$ also has the Global Glimm Property by [TV22b, Theorem 3.10].

REFERENCES

- [APRT21] R. ANTOINE, F. PERERA, L. ROBERT, and H. THIEL, Edwards' condition for quasitraces on C^* -algebras, *Proc. Roy. Soc. Edinburgh Sect. A* **151** (2021), 525–547.
- [APRT22] R. ANTOINE, F. PERERA, L. ROBERT, and H. THIEL, C^* -algebras of stable rank one and their Cuntz semigroups, *Duke Math. J.* **171** (2022), 33–99.
- [APT18] R. ANTOINE, F. PERERA, and H. THIEL, Tensor products and regularity properties of Cuntz semigroups, *Mem. Amer. Math. Soc.* **251** (2018), viii+191.
- [APT20] R. ANTOINE, F. PERERA, and H. THIEL, Cuntz semigroups of ultraproduct C^* -algebras, *J. Lond. Math. Soc. (2)* **102** (2020), 994–1029.
- [APT11] P. ARA, F. PERERA, and A. S. TOMS, K -theory for operator algebras. Classification of C^* -algebras, in *Aspects of operator algebras and applications*, *Contemp. Math.* **534**, Amer. Math. Soc., Providence, RI, 2011, pp. 1–71.
- [AVTV23] A. ASADI-VASFI, H. THIEL, and E. VILALTA, Dimension and radius of comparison of soft Cuntz semigroups, in preparation, 2023.
- [Bos22] J. BOSA, Stable elements and property (S), *J. Math. Anal. Appl.* **512** (2022), Paper No. 126171, 16.
- [BP09] L. G. BROWN and G. K. PEDERSEN, Limits and C^* -algebras of low rank or dimension, *J. Operator Theory* **61** (2009), 381–417.
- [BC09] N. P. BROWN and A. CIUPERCA, Isomorphism of Hilbert modules over stably finite C^* -algebras, *J. Funct. Anal.* **257** (2009), 332–339.
- [CR21] A. CHAND and L. ROBERT, Simplicity, bounded normal generation, and automatic continuity of groups of unitaries, *Adv. Math.* (to appear), preprint (arXiv:2108.09796 [math.OA]), 2021.
- [Chr17] M. S. CHRISTENSEN, *Regularity of C^* -algebras and central sequence algebras*, 2017, Thesis (Ph.D.)—University of Copenhagen (Denmark).
- [CRS10] A. CIUPERCA, L. ROBERT, and L. SANTIAGO, The Cuntz semigroup of ideals and quotients and a generalized Kasparov stabilization theorem, *J. Operator Theory* **64** (2010), 155–169.
- [CEI08] K. T. COWARD, G. A. ELLIOTT, and C. IVANESCU, The Cuntz semigroup as an invariant for C^* -algebras, *J. Reine Angew. Math.* **623** (2008), 161–193.
- [Cun78] J. CUNTZ, Dimension functions on simple C^* -algebras, *Math. Ann.* **233** (1978), 145–153.
- [Ell74] G. A. ELLIOTT, Derivations of matroid C^* -algebras. II, *Ann. of Math. (2)* **100** (1974), 407–422.
- [ERS11] G. A. ELLIOTT, L. ROBERT, and L. SANTIAGO, The cone of lower semicontinuous traces on a C^* -algebra, *Amer. J. Math.* **133** (2011), 969–1005.
- [ER06] G. A. ELLIOTT and M. RØRDAM, Perturbation of Hausdorff moment sequences, and an application to the theory of C^* -algebras of real rank zero, in *Operator Algebras: The Abel Symposium 2004*, *Abel Symp.* **1**, Springer, Berlin, 2006, pp. 97–115.
- [Eng14] M. ENGBERS, Decomposition of simple Cuntz semigroups, Ph.D Thesis, WWU Münster, 2014.
- [FHL⁺21] I. FARAH, B. HART, M. LUPINI, L. ROBERT, A. TIKUISIS, A. VIGNATI, and W. WINTER, Model theory of C^* -algebras, *Mem. Amer. Math. Soc.* **271** (2021), viii+127.
- [GP22] E. GARDELLA and F. PERERA, The modern theory of cuntz semigroups of C^* -algebras, preprint (arXiv:2212.02290 [math.OA]), 2022.
- [KNZ17] V. KAFTAL, P. W. NG, and S. ZHANG, Strict comparison of positive elements in multiplier algebras, *Canad. J. Math.* **69** (2017), 373–407.
- [KNZ19] V. KAFTAL, P. W. NG, and S. ZHANG, Purely infinite corona algebras, *J. Operator Theory* **82** (2019), 307–355.
- [Kir06] E. KIRCHBERG, Central sequences in C^* -algebras and strongly purely infinite algebras, in *Operator Algebras: The Abel Symposium 2004*, *Abel Symp.* **1**, Springer, Berlin, 2006, pp. 175–231.
- [KR02] E. KIRCHBERG and M. RØRDAM, Infinite non-simple C^* -algebras: absorbing the Cuntz algebra \mathcal{O}_∞ , *Adv. Math.* **167** (2002), 195–264.
- [KR15] E. KIRCHBERG and M. RØRDAM, When central sequence C^* -algebras have characters, *Internat. J. Math.* **26** (2015), 1550049, 32.

- [KNP10] D. KUCEROVSKY, P. W. NG, and F. PERERA, Purely infinite corona algebras of simple C^* -algebras, *Math. Ann.* **346** (2010), 23–40.
- [Lin91] H. X. LIN, Simple C^* -algebras with continuous scales and simple corona algebras, *Proc. Amer. Math. Soc.* **112** (1991), 871–880.
- [Lin04] H. LIN, Simple corona C^* -algebras, *Proc. Amer. Math. Soc.* **132** (2004), 3215–3224.
- [LN16] H. LIN and P. W. NG, The corona algebra of the stabilized Jiang-Su algebra, *J. Funct. Anal.* **270** (2016), 1220–1267.
- [NW10] P. W. NG and W. WINTER, Nuclear dimension and the corona factorization property, *Int. Math. Res. Not.* **2010** (2010), 261–278.
- [Ped72] G. K. PEDERSEN, Applications of weak* semicontinuity in C^* -algebra theory, *Duke Math. J.* **39** (1972), 431–450.
- [PR04] F. PERERA and M. RØRDAM, AF-embeddings into C^* -algebras of real rank zero, *J. Funct. Anal.* **217** (2004), 142–170.
- [Rob13] L. ROBERT, The cone of functionals on the Cuntz semigroup, *Math. Scand.* **113** (2013), 161–186.
- [Rob16] L. ROBERT, On the Lie ideals of C^* -algebras, *J. Operator Theory* **75** (2016), 387–408.
- [RR13] L. ROBERT and M. RØRDAM, Divisibility properties for C^* -algebras, *Proc. Lond. Math. Soc.* (3) **106** (2013), 1330–1370.
- [RT17] L. ROBERT and A. TIKUISIS, Nuclear dimension and \mathcal{Z} -stability of non-simple C^* -algebras, *Trans. Amer. Math. Soc.* **369** (2017), 4631–4670.
- [Rør91] M. RØRDAM, Ideals in the multiplier algebra of a stable C^* -algebra, *J. Operator Theory* **25** (1991), 283–298.
- [RW10] M. RØRDAM and W. WINTER, The Jiang-Su algebra revisited, *J. Reine Angew. Math.* **642** (2010), 129–155.
- [Sak71] S. SAKAI, Derivations of simple C^* -algebras. III, *Tohoku Math. J.* (2) **23** (1971), 559–564.
- [Thi17] H. THIEL, The Cuntz semigroup, lecture notes available at www.math.uni-muenster.de/u/hannes.thiel/, 2017.
- [Thi20] H. THIEL, Ranks of operators in simple C^* -algebras with stable rank one, *Comm. Math. Phys.* **377** (2020), 37–76.
- [TV21a] H. THIEL and E. VILALTA, Covering dimension of Cuntz semigroups II, *Internat. J. Math.* **32** (2021), 27 p., Paper No. 2150100.
- [TV21b] H. THIEL and E. VILALTA, Nowhere scattered C^* -algebras, *J. Noncommut. Geom.* (to appear), preprint (arXiv:2112.09877 [math.OA]), 2021.
- [TV22a] H. THIEL and E. VILALTA, Covering dimension of Cuntz semigroups, *Adv. Math.* **394** (2022), 44 p., Article No. 108016.
- [TV22b] H. THIEL and E. VILALTA, The Global Glimm Property, *Trans. Amer. Math. Soc.* **376**, (2023), 4713–4744. DOI: 10.1090/tran/8880, 2022.
- [TV23] H. THIEL and E. VILALTA, Soft C^* -algebras, preprint (arXiv:2304.11644 [math.OA]), 2023.
- [Vil22] E. VILALTA, A local characterization for the Cuntz semigroup of AI-algebras, *J. Math. Anal. Appl.* **506** (2022), Paper No. 125612, 47.
- [Win12] W. WINTER, Nuclear dimension and \mathcal{Z} -stability of pure C^* -algebras, *Invent. Math.* **187** (2012), 259–342.
- [WZ10] W. WINTER and J. ZACHARIAS, The nuclear dimension of C^* -algebras, *Adv. Math.* **224** (2010), 461–498.
- [Zha90] S. ZHANG, A Riesz decomposition property and ideal structure of multiplier algebras, *J. Operator Theory* **24** (1990), 209–225.

EDUARD VILALTA, DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, BARCELONA, SPAIN.

Email address: evilalta@mat.uab.cat

URL: www.eduardvilalta.com