

Numerical approximation of invariant objects: Fourier *vs* Wavelets

Treball de Recerca realitzat per:

David Romero i Sànchez

Sota la direcció de

Dr. Lluís Alsedà i Soler

Dr. Josep Maria Mondelo González

Contents

Part I Introduction to wavelets

1	Introduction	5
1.1	General definitions	5
1.2	Motivating example	6
1.3	Haar basis on $\mathcal{L}^2([0, 1])$	7
1.4	A comparison	8
2	Orthonormal Wavelet Basis	11
2.1	MRA and an orthonormal basis	11
2.2	Determining the scaling function: the <i>Low-Pass</i> Filter	17
2.3	Determining the Wavelet: the <i>High-Pass</i> Filter	20
2.4	Fast Orthonormal Wavelet Transform	24
2.5	How do we choose an orthonormal wavelet?	29
2.6	Drawing wavelets	31

Part II Wavelets applied to Dynamical Systems

3	A comparison between FWT and FFT	39
3.1	The CRTBP	40
3.1.1	The test for the CRTBP	44
3.2	The keller-GOPY attractor	46
3.2.1	The test for the keller-GOPY attractor	50
3.3	Conclusions	51
3.4	Future work	52
4	A numerical estimate of the Hölder exponent	53
4.1	Introduction	53

VIII Contents

4.2	The space of $\Lambda_\alpha(\mathbb{R})$	54
4.3	The Besov spaces $\mathcal{B}_{p,q}^s$	58
4.4	Daubechies wavelets in $\mathcal{B}_{\infty,\infty}^s(\mathbb{R})$	60
4.5	The algorithm	62
4.6	The algorithm on the Keller-GOPY SNA	64
4.7	Conclusions	69
4.8	Future work	69
Appendix A		71
Appendix B		75
6.1	Fourier Polynomials	75
6.2	Fast Fourier Transform	76
6.3	Using the FFT	77
References		79