

A NOTE ON THE ITERATION OF EXPONENTIALS

Carles Simó

Secció de Matemàtiques

Universitat Autònoma de Barcelona

We consider the sequence  $y_{i+1} = z^{y_i}$ ,  $i \geq 0$ ,  $z \in \mathbb{R}_+$ , with  $y_0 = z$ . In [1], the question of the behavior of such sequence is posed. Subsequently, many references to solutions are given (see [2]), for instance [3]. In this paper we obtain a full description of these iterates as functions of the parameter  $z$ , for every value of  $y$ . Our technique just uses the discrete dynamical system in  $\mathbb{R}_+$  defined by  $f_z(x) = z^x$ . The properties of the curves of fixed points and of two-periodic points are also given.

§ 1.- Fixed points.

a) If  $z > 1$ ,  $f_z(x)$  is concave;  $z^x = x$  has solution iff  $z \leq b$ , where  $b$  must satisfy  $x = b^x$ ,  $1 = b^x \ln b \implies b = e^{1/e}$ . If  $z \in (1, b)$  there are two fixed points  $x_1(z) < x_2(z)$ . They coincide for  $z = b$ .

For  $x_1(z)$  we have  $0 < f'_z(x_1(z)) < 1$ . Then it is stable. Instability occurs for  $x_2(z)$ .

b) If  $z < 1$ ,  $f_z(x)$  is monotonically decreasing. Then  $x = z^x$  has only one solution  $x_1(z)$ .

Stability:  $f'_z(x) < 0$  implies  $x_1(z)$  stable if  $f'_z(x_1) > -1$ . The limit of stability is found at  $x \ln z = -1 \implies x = 1/e$  and  $z_{\lim} = a = e^{-e}$ .

For  $z = 1 - \varepsilon$ ,  $\varepsilon$  small enough:  $z^{x_1} > 1 - \varepsilon \implies f'_z(x_1) = 0(\varepsilon)$ . Then the fixed point is stable for  $z \in [a, 1)$ . The negative character of  $f'$  implies that the iterates alternate around the fixed point.

c) Curve of fixed points: Consider the curve  $x = x(z)$ ,  $z \in (0, b]$  given by  $x = z^x$  (two branches if  $z > 1$ );  $x' = \frac{dx}{dz} = \frac{x}{z(1 - \ln x)} = \frac{\ln x}{z \ln z(1 - \ln x)}$ . One has  $x' = \infty$  at  $z = b$ . The upper branch has  $x'_2 < 0$ ,  $z \in (1, b)$ , and lower one gives  $x'_1(z) > 0$  in  $(0, b)$ . We get as limiting values:  $\lim_{z \rightarrow 0^+} x'_1 = -\lim_{z \rightarrow 0^+} \frac{1}{z \ln z} = \infty$ ;

$\lim_{z \rightarrow 1^+} x'_2 = -\infty$ ;  $\lim_{z \rightarrow 1} x'_1 = 1$ . We obtain for the second derivative

$$x'' = \frac{x \ln x + x / (1 - \ln x)}{z^2 (1 - \ln x)^2} \quad \text{zero values iff } \ln x = (1 \pm \sqrt{5})/2. \text{ Then}$$

there are only two turning points: one,  $x_2^i$ , in  $x_2(z)$  and the other,  $x_1^i$ , in  $x_1(z)$  for some  $z < 1$ . With this information we can plot  $x(z)$ . This is done in fig.1.

## § 2.- Periodic points.

a) Being  $f_z(x)$  increasing if  $z > 1$ , there are no periodic points. For  $z < 1$ ,  $f_z^2(x)$  is also increasing. Then there are only fixed points under  $f_z$  (studied in §1) or 2-periodic points  $x_3(z)$ ,  $x_4(z)$ .

b) We consider the function  $g_z(x) = z^x - \log_z x$  for  $z < a$ . We have  $g_z(x_1) = 0$ ,  $g'_z(x_1) < 0$  and  $g_z(1) > 0$ . Then there are points  $y \in (x_1, 1)$  fixed under  $g_z$ . Let us now show their uniqueness.

It is enough to prove that there is a unique point  $x$  such that  $g'_z(x) = 0$ . Then  $x z^x \ln^2 z = 1$ . We define  $\Psi(x) = x z^x$ . As  $\Psi'(x) = (1 + \ln x) z^x$ , we have for  $x > x_1$ :

$$|x \ln z| > |x_1 \ln z| = |\ln x_1| > 1 \Rightarrow \psi'(x) < 0 \text{ for } x \in (x_1, 1).$$

$$\text{But } f'_z(x_1) = z^{x_1} \ln z, \quad (f'_z(x_1))^2 = x_1 z^{x_1} \ln^2 z \Rightarrow \psi'(x_1) =$$

$$= (f'_z(x_1))^2 / \ln^2 z. \text{ Then } g'_z \text{ has a zero in } (x_1, 1) \text{ iff } x_1 \text{ is un-}$$

stable for  $f_z$ , i.e., iff  $z \in (0, a)$ . So there is only one 2-pe-  
riodic point in  $(x_1, 1)$  which is  $x_4(z)$ . The image under  $f_z$ ,

$x_3(z)$ , is also 2-periodic and belongs to  $(0, x_1)$ . The stabili-  
ty of 2-periodic points is guaranteed because  $g'_z(x_i) > 0$ ,  $i=3,4$ .

Furthermore, if  $h_z(x) = z^{z^x}$  we have  $h'_z(x_i) > 0$ ,  $i=3,4$ .

c) Curve of two-periodic points: There are two branches  
for  $z \in (0, a]$  which coincide if  $z=a$ . From  $z^x = \frac{\ln x}{\ln z}$  we derive  
 $z \ln z (x \ln x \ln z - 1) x' = -x \ln x (1 + x \ln z)$ . For  $x \ln x \ln z = 1$  we get  
 $x' = \infty$ . This happens if  $x = e^{-1}$ ,  $z = a$ . The signs of the factors  
allow us to state that  $x'_3 > 0$ ,  $x'_4 < 0$ . Indeed, we begin by proving  
that  $1 + x \ln z$  has only one zero:  $z^x = e^{-1} = \ln x / \ln z$  and  $x \ln z = -1$   
imply  $x \ln x = -e^{-1}$ , i.e.,  $x = e^{-1}$ .

The same happens for  $x \ln x \ln z - 1$ , but the proof is more  
tedious:  $z^x = e^{1/\ln x} = \ln x / \ln z$  and  $x \ln x \ln z = 1$  give us  
 $x \ln^2 x = e^{1/\ln x}$ . The change  $t = 1/\ln x$  transforms the above given  
condition to  $\xi(t) = \xi(t^{-1})$ , where  $\xi(t) = t e^t$ ,  $t < 0$ . We must veri-  
fy that  $t = -1$  is the unique solution. This is equivalent to find  
the positive solutions of  $\varphi(t) = t$ , where  $\varphi(t) = \exp(-\frac{1}{2}(t - 1/t))$ .  
Obviously 0, 1 are solutions. But  $\varphi'(t) = (t^2 + 1)\varphi(t)/(2t^2)$ ;  $\varphi''(t) =$   
 $= (t^4 + 2t^2 - 4t + 1)\varphi(t)/(4t^4)$ ;  $\varphi'''(t) = (t^6 + 3t^4 - 12t^3 + 27t^2 - 12t + 1)\varphi(t)/(8t^6)$ .  
Then,  $\varphi'(0) < 1$ ,  $\varphi'(1) = 1$ ,  $\varphi''(1) = 0$ ,  $\varphi'''(1) > 0$  implies that the  
number of zeros of  $\varphi(t) = t$  in  $(0, 1)$  counted with their multiplici-  
ties is even. If that number is positive,  $\varphi''(t)$  must have at least  
two zeros in  $(0, 1)$ , but such zeros satisfy  $t^4 + 2t + 1 = 4t$ . Since  
 $(t^2 + 1)^2$  is concave, there are exactly 2 solutions and one of them  
is 1. Then there are no solutions of  $\varphi(t) = t$  in  $(0, 1)$ . On the

other side  $\varphi''' > 0$  if  $t > 1$ , implies  $\varphi(t) > t, \forall t > 1$ . This ends the proof.

The behavior of the two branches near  $z=0$  is found by asymptotic expansions: Let  $x_3 = z(1 + \alpha(z))$ ,  $\alpha(z) = o(1)$ . We try to satisfy  $z^x = \ln x / \ln z$ . Then  $\alpha(z) = z \ln^2 z + o(z^2 \ln^2 z)$ . The image under  $f_z$  gives  $x_4 = 1 + z \ln z + o(z^2 \ln^2 z)$ . This allows us to plot  $x_i(z)$ ,  $i=3,4$ . See fig.1.

§ 3.- Behavior of the iterates.

Let be  $y_0, y_1, y_2, y_3, \dots$  the successive iterates.

a) If  $z > b$  one has  $y_n \uparrow \infty$ .

b) For  $z = b$  and  $y_0 \leq e$ , we have  $y_n \uparrow e$ .

For  $y_0 > e \Rightarrow y_n \uparrow \infty$ .

c) If  $z \in (1, b)$  and  $y_0 \leq x_1(z)$  we get

$y_n \uparrow x_1(z)$ ;  $y_0 \in (x_1, x_2) \Rightarrow y_n \uparrow x_1$ ;  $y_0 > x_2$

$\Rightarrow y_n \uparrow \infty$ .

d) When  $z \in [a, 1)$ , for every initial value  $y_0$  we have  $y_n \rightarrow x_1(z)$ , but the iterates alternate in  $(0, x_1), (x_1, \infty)$ .

So,  $y_0 \in (0, x_1) \Rightarrow y_{2k} \uparrow x_1, y_{2k+1} \downarrow x_1$ .

For the critical value  $z = a$  we have a bifurcation:  $x_1(z)$  loses the stability and a two-point stable cycle appears.

e) If  $z \in (0, a)$  we have also the fixed point  $x_1(z)$ , but any  $y_0 \neq x_1(z)$  gives iterates converging to the cycle

$x_{3,4}(z)$  (and then they do not properly converge).




Fig.1

$$y_0 \in (x_3, x_1) \Rightarrow y_{2k+1} \uparrow x_4, y_{2k} \downarrow x_3; y_0 \in (0, x_3) \Rightarrow y_{2k+1} \downarrow x_4, y_{2k} \uparrow x_3.$$

Similar results are obtained for  $y_0 \in (x_1, x_4)$  or  $y_0 \in (x_4, \infty)$ .

In particular, if  $y_0 = z$  the iterates converge to  $x_1$  iff  $z \in [a, b]$  and to the cycle  $\{x_3, x_4\}$  iff  $z < a$ .

### References

- [1] Notices Amer. Math. Soc. 25(1978), 197.
- [2] Notices Amer. Math. Soc. 25(1978), 253, 335.
- [3] Bromwich, T.J.I'A.: "An Introduction to the Theory of Infinite Series", MacMillan, 1965, p. 23.