

EQUIVARIANT MAPS UP TO HOMOTOPY
AND BOREL SPACES

Martin Fuchs

Equivariant maps between G -spaces induce fiber preserving maps between the associated Borel spaces. We will show that not all fiber preserving maps between Borel spaces are induced that way, not even all fiber homotopy classes of such maps. However there is a one-to-one correspondence between homotopy classes of G_∞ -maps (i.e. maps equivariant up to homotopy in a way, see section 1 for definitions) between G -spaces and fiber homotopy classes of maps between Borel spaces. This one-to-one correspondence is obtained by a functor equivalence between the respective categories (Theorem 1 and 2 in section 4). As a result equivariant homotopy theory (in a modified sense) is equivalent to the theory of homotopy fibrations.

To prove these theorems we have to include H -spaces into our discussion: In fact, the functor equivalence mentioned above is an extension of the equivalence between the categories of H -spaces and classifying spaces presented in [2]. Therefore we need the notion of a Borel space for H -spaces.

The Borel space we use, is associated with the modified Dold-Lashof construction in [3].

In section seven we present a number of examples of G-spaces with differing fix point sets, such that these differences cannot be detected by studying the cohomology of their Borel spaces, nor by studying the Borel space itself. The groups in most examples are \mathbb{Z}_p or S^1 , but the G-spaces are not all of finite dimension. Thus we illustrate the limits of theorems like the localization theorem by Hsiang ([5], p. 47). All the examples arise from the fact that if $h = \{h_n\}$ $n = 0, 1, \dots$ is a G_∞ -map between the G-spaces X_1 and X_2 and h is an ordinary homotopy equivalence, then the fiber map induced between the Borel spaces is a fiber homotopy equivalence.

1. Definitions

1.1. The H-spaces H we are using are supposed to be strictly associative and to have a strict unit element e . Furthermore we assume H has a homotopy inverse v (such that $H \xrightarrow{\Delta} H \times H \xrightarrow{1 \times v} H \times H \xrightarrow{\mu} H$ is homotopic to id_H).

1.2. We say that a topological space X is a G-space, if an H-space H acts on X from the left continuously and in a strictly associative manner. We assume that $ex = x$ for all $x \in X$.

1.3. As usual, an H_∞ -map h from H_1 to H_2 (of length r) is a sequence of continuous maps $h_n : (H_1 \times I_r)^n \times H_1 \rightarrow H_2$ ($n = 0, 1, 2, \dots$) such that

$$h_n(g_0, t_1, \dots, t_n, g_n)$$

$$= \begin{cases} h_{n-1}(g_0, t_1, \dots, g_{i-1}g_i, \dots, t_n, g_n) & t_i = r \\ h_{i-1}(g_0, t_1, \dots, g_{i-1})h_{n-i}(g_i, \dots, g_n) & t_i = 0 \end{cases}$$

for $n > 0$, $g_0, \dots, g_n \in H_1$, and $t_1, \dots, t_n \in I_r = [0, r] \subseteq \mathbb{R}$. If $r = 0$, the map h_0 is a homomorphism in the usual sense.

1.4. If H_1 acts on X_1 and H_2 acts on X_2 from the left, and if h is an H_∞ -map from H_1 to H_2 of length r , then we define a G_∞ -map f from X_1 to X_2 of length r associated with h to be a sequence of maps

$$f_n : (H_1 \times I_r)^n \times X_1 \rightarrow X_2 \quad (n = 0, 1, 2, \dots)$$

such that for $n > 0$

$$f_n(g_0, t_1, \dots, g_{n-1}, t_n, x) = \begin{cases} f_{n-1}(g_0, t_1, \dots, g_{i-1}g_i, \dots, g_{n-1}, t_n, x) & t_i = r \\ h_{i-1}(g_0, \dots, g_{i-1})f_{n-i}(g_i, \dots, g_{n-1}, t_n, x) & t_i = 0 \end{cases}$$

Composition of H_∞ -maps and G_∞ -maps is defined as in [3].

1.5. If f is a G_∞ -map from X_1 to X_2 associated to the H_∞ -map h from H_1 to H_2 , then f is called a G_∞ -homotopy equivalence if there exists an H_∞ -map k from H_2 to H_1 and a G_∞ -map g from X_2 to X_1

associated with k such that $g \circ f$ and $f \circ g$ are G_∞ -homotopic to id_{X_1} and id_{X_2} respectively (associated to the H_∞ -homotopies between $k \circ h$ respectively $h \circ k$ and id_{H_1} respectively id_{H_2}).

We are going to use the theorem from [4]:

Theorem. If H_1 acts on X_1 and H_2 acts on X_2 and if $h: H_1 \rightarrow H_2$ is an H_∞ -map such that h_0 is an ordinary homotopy equivalence, and if $f: X_1 \rightarrow X_2$ is a G_∞ -map associated with h such that f_0 is an ordinary homotopy equivalence, then h is an H_∞ -homotopy equivalence and f is a G_∞ -homotopy equivalence associated to h .

1.6. H -spaces and H_∞ -maps form the category \mathcal{K} and G -spaces and G_∞ -maps form the category \mathcal{L} . The associated homotopy categories are denoted by $\underline{\mathcal{K}}$ and $\underline{\mathcal{L}}$.

2. Construction of the Borel Space

In this section we rely heavily on [3], where many additional details can be found.

2.1. Let (p, r) be an H -principal fibration

$$\begin{array}{ccc} E \times H & \xrightarrow{r} & E \\ \text{pr}_1 \downarrow & & \downarrow p \\ E & \xrightarrow{p} & B \end{array}$$

as described in [3] and let X be a G -space with respect to H with action $s : H \times X \rightarrow X$. Assume that $p_X : EX \rightarrow B$ is a fibration with fiber X associated to $p : E \rightarrow B$ in the following sense: 1) The two fibrations are fiber homotopy trivial with respect to the same numerable covering \mathcal{U} of B and every $U \in \mathcal{U}$ is contractible in B . 2) There is a map $r_X : E \times X \rightarrow EX$ such that for each $U \in \mathcal{U}$ the diagram

$$(1) \quad \begin{array}{ccc} U \times H \times X & \xrightarrow{1 \times s} & U \times X \\ \alpha \uparrow & & \alpha_X \uparrow \\ p^{-1}(U) \times X & \xrightarrow{r_X} & p_X^{-1}(U) \\ & & \downarrow \beta_X \\ & & X \end{array}$$

is commutative ($(\alpha, \beta, \alpha_X, \beta_X)$ are the obvious coordinate maps). In addition we want

$$(2) \quad \begin{array}{ccc} E \times H \times X & \xrightarrow{1 \times s} & E \times X \\ \downarrow r_X \circ 1 & & \downarrow r_X \\ E \times X & \xrightarrow{r_X} & EX \end{array}$$

to be commutative.

2.2. For the general step of the Borel space construction we look at the H -principal fibration (\tilde{p}, \tilde{r}) as described in [3], p. 329-331.

The base space \tilde{B} of the new fibration is the mapping cone of $p : E \rightarrow B$ with the coordinate topology. We consider the covering of \tilde{B} consisting of

$$B_1 = \{y \perp t \mid t > \frac{1}{3}\} \text{ and } B_2 = \{y \perp t \mid t < \frac{2}{3}\}.$$

Let $p_1 : E_1 \rightarrow B_1$ respectively $p_{1X} : E_1X \rightarrow B_1$ be the fibrations induced by $f(y \perp t) = p(y)$, the map collapsing B_1 to the range space B of the mapping cone \tilde{B} . p_{1X} is associated to p_1 if we define $r_{1X} : E_1 \times X \rightarrow E_1X$ by

$$r_1(y \perp t, y_1, x) = (y \perp t, r_X(y_1, x)) .$$

Furthermore let $E_2 = B_2 \times H$ and $E_2X = B_2 \times X$. Define $r_{2X}(y \perp t, h, x) = (y \perp t, hx)$. Obviously these fibrations are associated.

We recall from [3], p. 330, that the map $F : p_2^{-1}(B_1 \cap B_2) \rightarrow p_1^{-1}(B_1 \cap B_2)$ defined by

$$F(y \perp t, h) = (y \perp t, yh)$$

is a strictly equivariant fiber homotopy equivalence. We define the associated map $F_X : p_{2X}^{-1}(B_1 \cap B_2) \rightarrow p_{1X}^{-1}(B_1 \cap B_2)$ by

$$F_X(y \perp t, x) = (y \perp t, r_X(y, x)) .$$

F_X is a map over $B_1 \cap B_2$ and a homotopy equivalence on each fiber (this follows from diagram (1) and the fact that H has a homotopy inverse) and hence is a fiber homotopy equivalence according to Theorem 6.3 in [1].

2.3. As in [3], p. 330 we now form the mapping cylinder of F and of F_X and construct the H -principal fibration $\tilde{p} : \tilde{E} \rightarrow \tilde{B}$ and similarly the associated fibration

$\tilde{p}_X : \tilde{E}X \rightarrow \tilde{B}X$. With the help of r_{X1} and r_{X2} we construct $\tilde{r}_X : \tilde{E} \times X \rightarrow \tilde{E}X$ in the obvious manner. No problem arises since the diagram

$$\begin{array}{ccc}
 (y \perp t, h, x) & \xrightarrow{r_{2X}} & (y \perp t, hx) \\
 \downarrow F_X \text{id} & & \downarrow F_X \\
 (y \perp t, yh, x) & \xrightarrow{r_{1X}} & (y \perp t, r_X(yh, x))
 \end{array} \quad \frac{1}{3} < t < \frac{2}{3}$$

commutes as a consequence of diagram (2). So it is easy to see that \tilde{E} and $\tilde{E}X$ are associated.

2.4. To construct the Borel space of X we start out with $p_0 : E_0 \rightarrow B_0$, where $E_0 = H$ and $B_0 = \{*\} = \text{point}$, and with $p_{0X} : E_0X \rightarrow B_0$, where $E_0X = X$. From p_n and p_{nX} we construct p_{n+1} and $p_{n+1,X}$ by letting $E_{n+1} = \tilde{E}_n$, $B_{n+1} = \tilde{B}_n$ and $E_{n+1}X = \tilde{E}_nX$. Obviously $p_{n+1} = \tilde{p}_n$ and $p_{n+1,X} = \tilde{p}_{nX}$ are associated. As on p. 333 in [3] we use telescopes to finally get the universal H -principal fibration $p_H : EH \rightarrow BH$ and the associated fibration $p_X : EX \rightarrow BH$. We call EX the Borel space of X and p_X the Borel fibration of X . Notice that p_X is a numerable, locally fiber homotopy trivial fibration with fiber X associated with p_H through the map $r_X : EH \times X \rightarrow EX$. r_X is essentially the direct limit of the maps $r_{n,X}$, and it is continuous because we used the telescope construction. (Compare the continuity of r_H in [3], p. 333).

3. Induced Maps Between Borel-Spaces

3.1. Before we can discuss G -spaces, we have to know more about H -spaces. So let $h: H_1 \rightarrow H_2$ be an H_∞ -map between the H -spaces H_1 and H_2 . We define a G_∞ -map $E_0 h: E_0 H_1 \rightarrow E_0 H_2$ as $E_0 h = h$. (Note that all the spaces $E_n H$ have a right action, so the notion of G_∞ -map has to be modified accordingly). Also we let $B_0 h: B_0 H_1 \rightarrow B_0 H_2$ be the trivial map.

Assume that $E_0 h$ has been extended to a G_∞ -map $E_n h: E_n H_1 \rightarrow E_n H_2$ associated with h and $B_0 h$ has been extended to $B_n h$ such that

$$p_{n2} \circ E_n h_k(y, t_1, g_1, \dots, t_k, g_k) = B_n h \circ p_{n1}(y).$$

(We will call a G_∞ -map with this property fiber preserving).

First we extend $B_n h$ from $B_n H_1$ to $\tilde{B}_n H_1$ by defining

$$\tilde{B}_n h(y \downarrow t) = (E_n h_0(y) \downarrow t)$$

On $E_{n1} H_1$ we define

$$E_{n1} h_0(y \downarrow t, y_0) = (E_n h_0(y) \downarrow t, E_n h_0(y_0))$$

and

$$\begin{aligned} E_{n1} h_k(y \downarrow t, y_0, t_1, g_1, \dots, t_k, g_k) \\ = (E_n h_0(y) \downarrow t, E_n h_k(y_0, t_1, g_1, \dots, t_k, g_k)) \end{aligned}$$

for $k = 1, 2, \dots$

Recall (from [3], p. 330) that $E_{n2}H'_1 = (B_{n2}H_1 \times H_1) \cup (B_{n1} \cap B_{n2} \times I \times H_1)$ and define

$$E'_{n2}h_k(y \perp t, \tau, g_0, t_1, \dots, t_k, g_k) = \begin{cases} (E_n h_0(y) \perp t, h_k(g_0, t_1, \dots, t_k, g_k)) \\ \quad \tau = 0, \quad 0 \leq t \leq \frac{1}{3} \\ \\ (E_n h_0(y) \perp t, 2\tau, h_k(g_0, t_1, \dots, t_k, g_k)) \\ \quad \text{when } 0 \leq \tau \leq \frac{1}{2} \text{ and } \frac{1}{3} < t < \frac{2}{3} \\ \\ (E_n h_0(y) \perp t, E_{n-k+1} h_k(y, 2\tau - 1, g_0, t_1, \dots, t_k, g_k)) \\ \quad \text{when } \frac{1}{2} \leq \tau \leq 1 \text{ and } \frac{1}{3} < \tau < \frac{2}{3}. \end{cases}$$

(When $\tau = 1$ we use that $E_{n-k+1} h_k(y, 1, g_0, t_1, \dots) = E_n h_k(yg_0, t_1, \dots)$). Hence $E'_{n2}h_k$ and $E_{n1}h$ together induce a G_ω -map $\tilde{E}_n h$ from $\tilde{E}_n H_1$ to $\tilde{E}_n H_2$ which satisfies all the conditions mentioned before and hence we get $E_{n+1} h : E_{n+1} H_1 \rightarrow E_{n+1} H_2$ together with $B_{n+1} h$. In the obvious manner we obtain the G_ω -map $Eh : EH_1 \rightarrow EH_2$ associated with h .

Because of our definition of $E'_{n2}h_k$ on the mapping cylinder part of $\tilde{E}_n H$, we only get $E(h \circ h')$ is G_ω -homotopic to $Eh \circ Eh'$ and similarly $B(h \circ h') \simeq Bh \circ Bh'$. In fact the G_ω -homotopy mentioned is fiber preserving. We get the

Theorem. The construction of universal fibrations described in [3] induces a functor $(\underline{E}, \underline{B})$ from the category \mathcal{K} as described in 1.6 to the category \mathcal{U} of universal fibrations and fiber homotopy classes of G_∞ -maps (with distinguished fiber).

3.2. Now let X be a topological space on which the H -space H acts from the left. The map $r_X : EH \times X \rightarrow EX$ discussed in section 2 is part of the structure of EX . A map between two Borel spaces has to preserve this structure at least up to homotopy. This leads to the following.

Definition. Let Y_1 and Y_2 be topological spaces on which H_1 and H_2 respectively act from the right, let X_1 and X_2 be topological spaces on which H_1 and H_2 respectively act from the left, and let $r_1 : Y_1 \times X_1 \rightarrow Z_1$ and $r_2 : Y_2 \times X_2 \rightarrow Z_2$ be maps (Z_1 and Z_2 are topological spaces) such that

$$\begin{array}{ccc}
 Y_i \times H_i \times X_i & \xrightarrow{1 \times \mu_{ri}} & Y_i \times X_i \\
 \mu_{li} \times 1 \downarrow & & \downarrow r_i \\
 Y_i \times X_i & \xrightarrow{r_i} & Z_i
 \end{array}$$

are commutative ($i = 1, 2$). Assume $h : H_1 \rightarrow H_2$ is a G_∞ -map and $k : Y_1 \rightarrow Y_2$ and $f : X_1 \rightarrow X_2$ are G_∞ -maps associated with h , then a G_∞ -map associated with

h, k , and f is a sequence of maps F_0, F_1, \dots such that

$$F_0 : Z_1 \rightarrow Z_2$$

and

$$F_k : Y_1 \times I \times (H_1 \times I)^{k-1} \times X_1 \rightarrow Z_2 \quad k = 1, 2, \dots$$

with

$$F_k(y, t_1, g_1, \dots, g_{k-1}, t_k, x) = \begin{cases} r_2(k_{i-1}(y, t_1, \dots, g_{i-1}), f_{k-i}(g_i, \dots, t_k, x)) & t_i = 0 \\ F_{k-1}(y, t_1, \dots, g_{i-1}g_i, \dots, t_k, x) & t_i = 1 \end{cases}$$

and appropriate modifications in special cases (like $k = 1$ or $i = 0$ and $i = k$).

3.3. Now we are ready to discuss Borel fibrations. Let

X_1 and X_2 be topological spaces on which H_1 and H_2 respectively act from the left. Assume $f : X_1 \rightarrow X_2$ is a G_∞ -map associated with the H_∞ -map $h : H_1 \rightarrow H_2$.

Again we define the G_∞ -map $E_0 f : E_0 X_1 \rightarrow E_0 X_2$ by $E_0 f = f$.

Assume we defined a G_∞ -map $E_n f : E_n H_1 \times X_1 \rightarrow E_n X_2$ in the sense of 3.2, associated with $E_n h, f$, and h . Furthermore we assume that all maps in $E_n f$ are "fiber-maps" over $B_n h$ in the obvious manner. Let us extend $E_n f$ to $\tilde{E}_n f : \tilde{E}_n H_1 \times X_1 \rightarrow \tilde{E}_n X_2$. We define $\tilde{E}_n f_0 : \tilde{E}_n X_1 \rightarrow \tilde{E}_n X_2$ first on

$$E_{n1}X_1 = \{(y \perp t, x_n) \mid (y \perp t \in B_n H_1, x_n \in E_n X, p(y) = p_X(y_n))\}$$

as

$$E_{n1}f_0(y \perp t, x_n) = (E_n h_0(y) \perp t, E_n f_0(x_n))$$

Then we define for $k = 1, 2, \dots$

$$\begin{aligned} E_{n1}f_k(y \perp t, y_0, t_1, \dots, g_{k-1}, t_k, x) \\ = (E_n h_0(y) \perp t, E_n f_k(y_0, t_1, \dots, g_{k-1}, t_k, x)) \end{aligned}$$

where $(y \perp t, y_0) \in E_{n1}H_1$, $x \in X_1$, $g_i \in H_1$ and $t_i \in I$.

On $E_{n2}X'_1$ we define for $k = 0$

$$\begin{aligned} E'_{n2}f_0(y \perp t, \tau, x) \\ = \begin{cases} (E_n h_0(y) \perp t, f_0(x)) & 0 \leq t \leq \frac{1}{3}, \quad \tau = 0 \\ (E_n h_0(y) \perp t, 2\tau, f_0(x)) & \frac{1}{3} < t < \frac{2}{3}, \quad 0 \leq \tau \leq \frac{1}{2} \\ (E_n h_0(y) \perp t, E_n f_1(y, 2\tau-1, x)) & \frac{1}{3} < t < \frac{2}{3}, \quad \frac{1}{2} \leq \tau \leq 1 \end{cases} \end{aligned}$$

and for $k = 1, 2, \dots$ we define $E'_{n2}f_k$ just like $E'_{n2}h_k$ with the following changes: replace h_k and h_{k+1} by f_k and f_{k+1} respectively and g_k by x . $E'_{n2}f_k$ and $E_{n1}f_k$ can be pieced together to obtain $\tilde{E}_n f_k$ for $k = 0, 1, 2, \dots$. Ultimately we get the G_∞ -map

$\{Ef\}: EH_1 \times X_1 \rightarrow EX_2$ "over" $Bh: BH_1 \rightarrow BH_2$ associated with Eh, f and h .

3.3. We point out that if $h, k: H_1 \rightarrow H_2$ are H_∞ -maps

which are H_∞ -homotopic, then Bh is homotopic to Bk

leaving the base point fixed, and Eh is G_ω -fiber homotopic to Ek over the homotopy between Bh and Bk .

Furthermore if $f, g: X_1 \rightarrow X_2$ are G_ω -maps associated to h and k , and if f, g are G_ω -homotopic associated to the H_ω -homotopy between h and k , then Ef and Eg are fiber homotopic associated with the G_ω -fiber homotopy between Eh and Ek etc. and over the homotopy between Bh and Bk .

Definition. Let \mathcal{F} be the category whose objects are fibrations $p: E \rightarrow B$ which are locally fiber homotopy trivial with respect to a numerable covering of sets contractible in B , and whose morphisms are fiber homotopy classes of fiber preserving maps. Let \mathcal{F}_* be the associated category of fibrations with a distinguished fiber over a basepoint $*$, and let $\underline{\mathcal{F}}$ and $\underline{\mathcal{F}}_*$ be the associated homotopy categories.

Theorem. The constructions EH , BH , and EX define a functor $B: \underline{\mathcal{F}} \rightarrow \underline{\mathcal{F}}_*$, the Borel functor.

4. The Inverse Functor of B

For every topological space X and subsets $A, B \subseteq X$ we recall that

$$L(X; A, B) = \{(\omega, r) \mid \omega: \mathbb{R}^+ \rightarrow X, \omega(0) \in A, \\ \omega(t) = \omega(r) \in B \text{ for } t \geq r\}$$

Often we omit r in our notation for the sake of simplicity.

Definition. For every fibration $p: E \rightarrow B$ with distinguished fiber $F_* = p^{-1}(*)$ we define

$$\bar{E} = \{(\omega, y) \mid y \in E, \omega \in L(B; B, B), \omega(r) = p(y)\}$$

and $\bar{p}: \bar{E} \rightarrow \bar{B}$ as $\bar{p}(\omega, y) = \omega(0)$.

If the fibration $p: E \rightarrow B$ is an object in \mathcal{F}_* then the fiber map $\tau: E \rightarrow \bar{E}$ defined by $\tau(y) = (\omega_y, y)$ is a fiber homotopy equivalence, see [1], Theorem 6.3 ($\omega_y: \mathbb{R}^+ \rightarrow E$ is defined as $\omega_y(t) = y$ for all $t \in \mathbb{R}^+, r = 0$).

Let $WE = \bar{p}^{-1}(*)$ be the distinguished fiber of \bar{p} , then $\tau|_{F_*}$ is a homotopy equivalence between F_* and WE . We observe that the loop space of B , $\Omega(B, *)$, acts on WE from the left ($\Omega(B, *) = L(B; *, *)$ is an H-space). Furthermore if p, p' are two fibrations in \mathcal{F}_* and if (F, f) is a based fiber map from p to p' , then $Wf: WE \rightarrow WE'$ defined by $Wf(\omega, y) = (Lf(\omega), F(y))$ is an equivariant map associated with the induced homomorphism $\Omega f: \Omega(B, *) \rightarrow \Omega(B', *)$. We summarize this observation in the

Definition. W induces a functor

$$\underline{W}: \underline{\mathcal{F}}_* \rightarrow \underline{\mathcal{L}},$$

the inverse functor to \underline{B} , as we shall see in the following

Theorem 1. \underline{WB} is equivalent to $\underline{1}_{\underline{\mathcal{L}}}$

and

Theorem 2. \underline{BW} is equivalent to $\underline{1_x}$.

5. Proof of Theorem 1

To prove Theorem 1 we have to review the natural transformation $S : H \rightarrow \Omega BH$.

5.1. We need from [3], p. 333 the

Theorem. EH is contractible.

Let $k : EH \times I \rightarrow EH$ be a contraction with $k(y, 0) = y$ and $k(y, 1) = * = k(*, t)$. (For this it is necessary that $* \in H$ is a nondegenerate base point. If necessary one can switch to $H \vee I$, see [2], p. 215).

Associated with the contraction k is the map $K : EH \rightarrow L(EH; EH, *)$ defined by $K(y) = (k(y, t), 1)$.

5.2. Define $S_0 : H \rightarrow \Omega(BH, *)$ as

$$S_0(y) = Lp_H \circ K|_{E_0 H}$$

with $Lp_H : L(EH; EH, *) \rightarrow L(BH; BH, *)$ induced by p_H .

Lemma 1. S_0 is a homotopy equivalence.

Proof: $L(BH; BH, *)$ is the total space of a numerable fibration over BH , and so is EH . Both total spaces are contractible. S_0 is the restriction of $Lp_H \circ K$, which is a fiber map over id_{BH} and which is also a homotopy equivalence. Theorem 6.1 in [1]

implies that $Lp_H \circ K$ is a fiber homotopy equivalence and hence S_0 is a homotopy equivalence.

Lemma 2. S_0 can be extended to an H_∞ -map.

Proof: Let $K|_{E_0H} = K|_H = K_0$. Then we have to find maps S_1, S_2, \dots which make $S_0 = Lp_H \circ K_0 : H \rightarrow \Omega BH$ into an H_∞ -map. Assume we already constructed $S_i = Lp_H \circ K_i$ ($i = 0, 1, \dots, n$). Then S_{n+1} and hence K_{n+1} is defined on $\partial H(n+1)$ through the maps S_i and K_i respectively ($i = 0, \dots, n$).

Associated with K_i are the maps

$$k_i : H(i) \times \mathbb{R}^+ \rightarrow EH$$

and

$$r_i : H(i) \rightarrow \mathbb{R}^+$$

with $k_i(g_0, t_1, \dots, t_i, g_i, 0) = *$ and $k_i(g_0, t_1, \dots, t_i, g_i, \tau) = g_0 \dots g_i$ for $\tau \geq r_i(g_0, t_1, \dots, t_i, g_i)$. These maps define k_{n+1} and r_{n+1} respectively on $\partial H(n+1)$. Since \mathbb{R}^+ is contractible we can extend r_{n+1} to all of $H(n+1)$. Then we can extend k_{n+1} to all of $H(n+1)$ such that $k_{n+1}(g_0, t_1, \dots, t_{n+1}, g_{n+1}, 0) = *$ and $k_{n+1}(g_0, t_1, \dots, t_{n+1}, g_{n+1}, r_{n+1}(\dots)) = g_0 \dots g_{n+1}$, since EH is contractible.

Define

$$K_{n+1} = (k_{n+1}, r_{n+1}) \quad \text{and} \quad S_{n+1} = Lp_H \circ K_{n+1}.$$

For further details compare [2], p. 214-215. (Note the addition of paths on p. 213 should be reversed.)

5.2. Proposition. S is a natural transformation between

$$L_{\mathbb{H}} \text{ and } \Omega B.$$

Proof: In the diagram

$$\begin{array}{ccccc}
 & H & \xrightarrow{h} & H' & \\
 & K \downarrow & & \downarrow K & \\
 S \swarrow & L(EH; EH, *) & \xrightarrow{LEh} & L(EH'; EH', *) & \\
 & Lp_H \downarrow & & \downarrow Lp_{H'} & \\
 & \Omega(BH, *) & \xrightarrow{\Omega Bh} & \Omega(BH', *) &
 \end{array}$$

the lower portion commutes for all the maps of LEh .

To see that the upper portion commutes up to an H_∞ -homotopy, one has to look again at the associated maps into EH' . Since EH' is contractible, all extensions necessary to construct the H_∞ -homotopy between $LEh \circ K$ and $K \circ h$ can be carried out. Further details in [2].

(In [2] the G_∞ -map Eh was not discussed. Instead the notion of a "regular" H -homomorphism had to be used. Now EH provides the homotopy between formula 2 and 2a on p. 217 in 2, translated from right to left actions.)

5.3. With S out of the way we define for any G -space X :

$$T_0 : X \rightarrow WE \quad \text{as} \quad T_0 = \tau \mid X .$$

We already know that $T_0(Y) = (*, y)$ is a homotopy equivalence. We define $T_n : (H \times I)^n \times X \rightarrow WE$ as

$$\begin{aligned} T_n(g_0, t_1, \dots, t_n, x) \\ = (pK_{n-1}(g_0, \dots, t_{n-1}, g_{n-1})(t_n + \sigma), \\ r_X(K_{n-1}(g_0, \dots, t_{n-1}, g_{n-1})(t_n), x)) \end{aligned}$$

with $0 \leq t_n \leq r_{n-1}(g_0, \dots, t_{n-1}, g_{n-1})$ and $0 \leq \sigma \leq r_{n-1} - t_n$. Recall $r_X : EH \times X \rightarrow EX$. We have

$$\begin{aligned} T_n(g_0, t_1, \dots, g_{n-1}, t_n, x) \\ = \begin{cases} (S_{n-1}(g_0, t_1, \dots, g_{n-1}), x) & t_n = 0 \\ (*, g_0 g_1 \dots g_{n-1}, x) & t_n = r . \end{cases} \end{aligned}$$

The " G_∞ -homotopy" between $LEh \circ K$ and $K \circ h$ implies that T is a natural transformation between $\underline{l}_\mathcal{E}$ and \underline{WB} .

6. Proof of Theorem 2

6.1. Let \mathcal{J}_* be the category of based topological spaces X , which have a numerable covering \mathcal{U} such that every $U \in \mathcal{U}$ is contractible in X , and based continuous maps. Let $\underline{\mathcal{J}}_*$ be the associated homotopy category.

Remark. It is easy to see that for every H in \mathcal{K} the classifying space BH is in \mathcal{T}_* .

In preparation for the proof of Theorem 2 we list three universal fibrations with fiber $\Omega(X,*)$ for $X \in \mathcal{T}_*$.

a) Application of the modified Dold-Lashof construction to the trivial fibration $\Omega(X,*) \rightarrow *$ leads to

$$P_{\Omega X} : E\Omega X \rightarrow B\Omega X$$

b) It is well-known that

$$P_L : L(X;X,*) \rightarrow X$$

also classifies numerable $\Omega(X,*)$ -fibrations.

c) If we apply the modified Dold-Lashof construction to p_L of b), we get again a universal fibration

$$P_{EL} : ELX \rightarrow BLX .$$

All three constructions induce functors from \mathcal{T}_* to \mathcal{T}_* .

6.2. The inclusion of $\Omega(X,*)$ as distinguished fiber of $p_L : L(X;X,*) \rightarrow X$ can be interpreted as a principal map of principal fibrations and hence it induces the fiber map (f, \bar{f}) :

$$\begin{array}{ccc}
 E(\Omega X) & \xrightarrow{f} & E(LX) \\
 P_{\Omega X} \downarrow & & \downarrow P_{LX} \\
 B(\Omega X) & \xrightarrow{\bar{f}} & B(LX)
 \end{array}$$

which is a principal fiber homotopy equivalence; (f, \bar{f}) is an inclusion, hence $P_{\Omega X}$ is principal fiber homotopy equivalent to the pullback of P_{LX} . For universal fibrations this implies \bar{f} is a homotopy equivalence. Let \bar{g} be a homotopy inverse of \bar{f} .

As a result, (f, \bar{f}) represents a functor equivalence between the functors from \mathcal{J}_* to \mathcal{I}_* induced by a) and c).

6.3. The inclusion

$$\begin{array}{ccc}
 L(X; X, *) & \xrightarrow{k} & ELX \\
 P_L \downarrow & & \downarrow P_{LX} \\
 X & \xrightarrow{\bar{k}} & BLX
 \end{array}$$

is a fiber homotopy equivalence by the same reasoning as described in 6.2. So (k, \bar{k}) represents a functor equivalence between the functors arising from b) and c).

6.4. Now consider a fibration $p: E \rightarrow X$ from the category \mathcal{I}_* . The associated Hurewicz-fibration $\bar{p}: \bar{E} \rightarrow X$ admits a map

$$r_0: L(X; X, *) \times WE \rightarrow \bar{E}$$

defined through the addition of paths, which makes \bar{E} a look alike of a Borel space associated to WE .

Assigning to p the Hurewicz fibration \bar{p} induces a functor H_r on \mathcal{F}_* which is obviously equivalent to $\text{id}_{\mathcal{F}_*}$. We are now going to show $\underline{BW} \simeq \underline{Hr}$. Consider the diagram of Borel spaces:

$$\begin{array}{ccccc}
 L(X; X, *) \times WE & \xrightarrow{k \times 1} & ELX \times WE & \xrightarrow{g \times 1} & E \cap X \times WE \\
 \downarrow \bar{E} & \xrightarrow{K} & \downarrow & \xrightarrow{G} & \downarrow \\
 \bar{p} \downarrow & & EL(WE) & & E(WE) \\
 X & \xrightarrow{\bar{k}} & \downarrow & \xrightarrow{\bar{g}} & \downarrow \\
 & & BLX & & B \cap X
 \end{array}$$

K is induced by applying the Borel space construction to \bar{p} (an obvious modification) and G is induced by g , the homotopy inverse of f from 6.2.

(K, \bar{k}) and (G, \bar{g}) represent functor equivalences associated to the equivalences (k, \bar{k}) and (g, \bar{g}) discussed in 6.2 and 6.3. Since the right side of the diagram represents BW and the left side represents H_r , the proof is complete.

7. Two Applications

7.1. Let $G = \mathbb{R}^1$ and $X = \mathbb{R}^2$. Consider the two \mathbb{R}^1 -spaces X_1 and X_2 defined by the two actions

$$\mu_1 : \mathbb{R}^1 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad \mu_1(t, re^{i\varphi}) = re^{i(\varphi+t)},$$

$$\mu_2 : \mathbb{R}^1 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad \mu_2(t, re^{i\varphi}) = re^{i(\varphi+t(1-r))}$$

The fix point set of μ_1 is just the origin of \mathbb{R}^2 and the fix point set of μ_2 is the origin and the unit circle. Obviously we could define actions with more complicated fix point sets.

The constant map from one of these spaces to the origin of the other is an equivariant map which is also an ordinary homotopy equivalence. It induces (according to section four) a homotopy equivalence between the Borel spaces of the two spaces.

7.2. a) Let P be an acyclic finite polyhedron with nontrivial fundamental group. Then the suspension ΣP is a contractible \mathbb{Z}_2 -space with fix point set P , and the join $P * S^1$ is a contractible S^1 or \mathbb{Z}_p -space ($p \neq 2$) with fix point set P in the obvious manner (notice $P * S^1 \cong \Sigma^2 P$).

b) Let P be any finite polyhedron. The obvious \mathbb{Z}_2 -action on ΣP can be extended to $\Sigma^2 P$ etc. so that $\lim_{n \rightarrow \infty} \Sigma^n P$ is a contractible \mathbb{Z}_2 -space with fix point set P .

For $G = \mathbb{Z}_p$ ($p \neq 2$) and $G = S^1$ we can do the same by reiterating the join with S^1 .

7.3. Let G be either \mathbb{Z}_p or S^1 and let X be a G -space with fix points. Let Y be a contractible G -space with nonempty fix point set F , e.g. let Y be one of the spaces mentioned above. The one point union W of X and Y formed by identifying two

fix points is a new G -space in the obvious manner and the inclusion of X into W is an equivariant map and also an ordinary homotopy equivalence.

By the theorem in [4] the inclusion represents an isomorphism in \mathcal{Z} and induces a fiber homotopy equivalence between BX and BW by section 4. Hence the cohomology of these Borel spaces carries no information about F .

7.4. Assume G is either \mathbb{Z}_p^k or $(S^1)^k$ and X_1, X_2 are G -spaces which satisfy the assumptions for Borel's theorem as described in Proposition 1 of Chapter IV in [5], i.e., let X_1, X_2 be paracompact G -spaces with finite cohomology dimension. Let $f: X_1 \rightarrow X_2$ be an equivariant map which is also an ordinary homotopy equivalence. Again $Ef: EX_1 \rightarrow EX_2$ is a fiber homotopy equivalence between Borel spaces. Ef induces isomorphisms between $H_G^*(X_2)$ and $H_G^*(X_1)$ as $H^*(BG)$ modules. Hence Proposition 1 on p.45 in [5] tells us, that $f|_{F_1}: F_1 \rightarrow F_2$ induces an isomorphism of the cohomology rings $H^*(F_2) \otimes_k R_0$ and $H^*(F_1) \otimes_k R_0$ of the fix point sets F_1 and F_2 .

T. Petrie in [7] and elsewhere, Ch. N. Lee and A. Wasserman in [6] have constructed examples of such maps which do not have equivariant homotopy inverses. Hence the fiber homotopy inverse of Ef is not induced by an equivariant map from X_2 to X_1 . This answers the opening statement of the introduction of this paper.

References

1. Dold, A., Partitions of unity in the theory of fibrations, *Ann. of Math.* 78, 223-255 (1963).
2. Fuchs, M., Verallgemeinerte Homotopie-Homomorphismen und klassifizierende Räume, *Math. Ann.* 161, 197-230 (1965).
3. Fuchs, M., A modified Dold-Lashof construction that does classify H-principal fibrations, *Math. Ann.* 192, 328-340 (1971).
4. Fuchs, M., Homotopy equivalences in equivariant topology, *Proc. Amer. Math. Soc.* 58, 347-352 (1976).
5. Hsiang, W.Y., *Cohomology theory of topological transformation groups*, Springer-Verlag 1975.
6. Lee, Ch. N. and Wasserman, A.G., On the groups $JO(G)$, *Memoirs Amer. Math. Soc.* 159 (1975).
7. Petrie, T., Smooth S^1 actions and bilinear forms, *Bull. Amer. Math. Soc.* 79, 1056-1059 (1973).

Rebut el 15 de setembre del 1983

Department of Mathematics
Michigan State University
East Lansing, MI 48824
USA