Pub. Mat. UAB
Vol. 28 nº 2-3 Set. 1984

POLYNOMIAL RINGS OVER NON-COMMUTATIVE RINGS*

Poobhalan Pillay

Let R be an associative, not necessarily commutative ring with identity. It is shown that the polynomial ring $R[x_\alpha]$, where the x_α's constitute an arbitrary set of central indeterminates, has a semisimple (resp. right Artinian, Quasi-Frobenius) classical right quotient ring if and only if, R has the same property.

In [10], Lance Small showed that if an associative ring R with identity element is a right order in a semisimple ring $Q_{c\ell}(R)$, then so is the polynomial ring $R[x_1,x_2,\ldots x_n]$ where the x's are central indeterminates. In [11] he showed that a similar conclusion holds if R is a Noetherian order in a right Artinian ring. Shock [9] extended these results to the case where the indeterminates form a countably infinite set. In continuing these investigations we prove in this paper that if I is any index set, a ring R is a right order in a P-ring if and only if $S = R[x_\alpha]_{\alpha\varepsilon I}$ is a right order in a P-ring where P is any one of P_1: semisimple, P_2: right Artinian, P_3: Quasi-Frobenius.

For the case $P = P_1$, a connection between the associated matrix of rings of $Q_{c\ell}(S)$ and $Q_{c\ell}(R)$ is established. (Theorem 2.5).

Sufficiency for the cases $P = P_2$ and $P = P_3$ exploit the fact that any polynomial ring over R has among its ring of quotients, a group ring RG where G is a free abelian group, the rank of which can be chosen to be the cardinality of I. In establishing necessity for

*This paper represents a portion of the author's Ph.D. thesis written under the direction of Dr. R. W. Wilkerson at the University of the Witwatersrand, Johannesburg. Research supported by CSIR (South Africa). AMS Subject Classification (1970), 16A08

P_3, it is first observed that if $Q_{max}(S)$ is self-injective, so is $Q_{max}(R)$.

The set of all polynomials which are not zero-divisors has not yet been successfully described in terms of the coefficient ring R, even when R is commutative or I finite. However, in each of the cases P_1, P_2, P_3 we exhibit a well described right Ore semigroup M(P) of S such that

$$Q_{c\ell}(S) = \{fg^{-1} | f \in S,\ g \in M(P)\}$$

§1. Preliminaries: Throughout this paper, a ring R is associative with identity element. Objects of mod-R will be denoted by M_R, and will be unitary. If S is a subset of R, the right and left annihilators of S in R will be denoted by $r_R(S)$ and $\ell_R(S)$ respectively. An element a of R is regular if and only if $r_R(a) = \ell_R(a) = 0$. The set of all regular elements of R/A, where A is a two sided ideal of R, will be denoted by $C_R(A)$. In particular, $C_R(0)$ is the set of all regular elements of R.

Defn. 1.1: Let $\{x_\alpha\}_{\alpha \in I}$ be any set indexed by a set I of arbitrary cardinality, and let $<$ be a (fixed) well ordering on I. Let G be the (multiplicatively written) free abelian group on $\{x_\alpha\}$, and

$$RG = \{f: G \to R | f \text{ has finite support}\},$$

the associated group ring. Let

$$M = \{x_{\alpha_1}^{n_1} x_{\alpha_2}^{n_2} \cdots x_{\alpha_k}^{n_k} \in G | n_i \geq 0 \text{ for each } i\}$$

Then M is a submonoid of G and the subring $RM = \{f \in RG | f(g) = 0$ for every $g \notin M\}$ is called the polynomial ring in $\{x_\alpha\}_{\alpha \in I}$ over R. Henceforth, we shall denote RM by S or $R[x_\alpha]$.

If $\phi: R \to RG$ and $\psi: G \to RG$ are defined by

$$\phi(a)(g) = a \text{ if } g = 1, \text{ and } 0 \text{ otherwise}$$

$$\psi(g)(g') = 1 \text{ if } g' = g, \text{ and } 0 \text{ otherwise}$$

then ϕ (resp. ψ) is a ring (resp. monoid) monomorphism. For $a \in R$ and $g \in G$, we write ag to mean $\phi(a)\psi(g) \in RG$. Then each $f \in R[x_\alpha]$ has a decomposition

$$f = \sum_{i=1}^{n} f_i m_i$$

where $f_i \in R$, $m_i \in M$ for each i, and $f = 0$ if and only if $f_i = 0$ for every i. The set of <u>coefficients</u> of f will be denoted by $\{f_i\}_{i=1}^{n}$.

Our definition of a polynomial ring is easily seen to be consistent with the "usual" definition for a polynomial ring. In fact for any $n \geq 2$

$$R[x_1, x_2, \ldots, x_n] \simeq R[x_1, x_2, \ldots, x_{n-1}][x_n]$$

canonically as rings.

If $f \in R[x_\alpha]$, the <u>leading coefficient</u> $L(f) \in R$ of f is defined as follows. If $f = 0$, let $L(f) = 0$. Suppose $f \neq 0$. If $|I| = 1$, $L(f)$ is defined as usual. In general, $\exists$ a finite subset $\{\alpha_1, \alpha_2, \ldots, \alpha_n\}$ of I such that $\alpha_1 < \alpha_2 < \ldots < \alpha_n$ and $f \in R[x_{\alpha_1}, x_{\alpha_2}, \ldots, x_{\alpha_n}]$.

Since $R[x_{\alpha_1}, x_{\alpha_2} \ldots, x_{\alpha_n}] \simeq R[x_{\alpha_1}, x_{\alpha_2}, \ldots, x_{\alpha_{n-1}}][x_{\alpha_n}]$ we may regard f as a polynomial in x_{α_n} and proceed by induction to obtain $0 \neq L(f) \in R$.

The following lemma follows from the definitions.

<u>Lemma 1.2</u>: For any f, g in $R[x_\alpha]$,

$$L(f)L(g) \neq 0 \Rightarrow L(fg) = L(f)L(g) \Rightarrow L(fg) \neq 0$$

Hence if $fg = 0$, then $L(f)L(g) = 0$.

Next we examine some connections between the lattice of right ideals of a polynomial ring and that of its coefficient ring. Let $S = R[x_\alpha]$. By $L_r(R)$ (resp. $L_r(S)$ we shall mean the lattice of right ideals of R (resp.S). If $K \in L_r(R)$, let

$$K[x_\alpha] = \{f \in S \mid f_i \in K \text{ for all } i\}$$

Then $K[x_\alpha] = KS$ is the right ideal in S generated by K.

Defn. 1.3: Define $\omega : L_r(R) \to L_r(S)$ and

$$\Omega : L_r(S) \to L_r(R)$$

by $\omega(K) = K[x_\alpha]$ and

$\Omega(J) = J \cap R$

The next lemma requires a routine verification.

Lemma 1.4: (i) For all $K \in L_r(R)$, $(\Omega\omega)(K) = K$, so Ω is surjective.

(ii) ω is 1-1, preserves inclusions, arbitrary and direct sums, arbitrary intersections and finite products.

(iii) If A is a subset of R and

$$A' = \{f \in S \mid f_i \in A \text{ for every } i\}$$

then $\omega(r_R(A)) = r_S(A')$ so takes right annihilators into right annihilators. Moreover, if A is an ideal,

$$\omega(\ell_R(A)) = \ell_S(\omega(A))$$

(iv) Ω preserves inclusions and arbitrary intersections.

A ring is right Goldie finite dimensional (abbreviated right f.d.) if any direct sum of right ideals of R has at most finitely many non-zero summands. A right Goldie ring is a right f.d. ring which satisfies the maximum condition on annihilator right ideals.

Lemma 1.5: If R is right f.d., so is $S = R[x_\alpha]_{\alpha \in I}$. If S is right Goldie, so is R.

Proof: The first part is due to Shock (Theorem 2.6 of [9]) where the statement is proved for countable I. The same proof adapts readily to the general case.

Let S be right Goldie. If $\{K_\beta\}_{\beta \in B}$ is an independent family of right ideals in R with

$$K = \bigoplus_{\beta \in B} K_\beta$$

then by 1.4 (ii)

$$\omega(K) = \bigoplus_{\beta \in B} \omega(K_\beta)$$

so $\omega(K_\beta) = 0$ (hence $K_\beta = 0$) for all but finitely many β, proving R is right f.d.

Let $K_1 \subseteq K_2 \subseteq \ldots \subseteq K_n \subseteq \ldots$ (*)

be any ascending sequence of right annihilators in R. By 1.4,

$$\omega(K_1) \subseteq \omega(K_2) \subseteq \ldots \subseteq \omega(K_n) \subseteq \ldots \quad (**)$$

is an ascending chain annihilator right ideals of S so that (**), hence (*), becomes eventually constant.

The prime radical rad R is the meet of all the prime ideals of R. R is semiprime if and only if rad R = 0. The right singular ideal of R is the ideal

$$Z_r(R) = \{a \in R \mid r_R(a) \text{ is an essential right ideal of } R\}.$$

Lemma 1.6: Let $S = R[x_\alpha]$. Then

(i) rad $S = \omega(\text{rad } R)$

(ii) $Z_r(S) = \omega(Z_r(R))$

Proof: (i) is due to Amitsur [1] and (ii) to Shock [9].

Lemma 1.7: Let A be an ideal of R. Then

$$R[x_\alpha]/A[x_\alpha] = S/\omega(A) \cong R/A[x_\alpha].$$

Proof: Let $\pi: R \to R/A$ be the natural ring epimorphism and define

$$\bar{\pi}: S \to R/A[x_\alpha] \text{ by}$$

$$\bar{\pi}(f) = \sum_{i=1}^{n} \pi(f_i)m_i \text{ whenever } f = \sum_{i=1}^{n} f_i m_i \in S.$$

Then $\bar{\pi}$ is a ring epimorphism with kernel $A[x_\alpha] = \omega(A)$.

§2. ORDERS IN SEMISIMPLE RINGS: The maximal ring of right quotients of R is denoted by $Q_{max}(R)$. Asano [2] has shown that if N is a multiplicatively closed subset of $C_R(0)$ and R satisfies the right Ore condition with respect to N i.e. (for all $a \in R$, $n \in N$, $\exists\ a_1 \in R$, $n_1 \in N$ such that $an_1 = na_1$) then there is a ring R_N containing R, unique up to an isomorphism that extends the identity map of R satisfying

(i) if $n \in N$, n is a unit in R_N

(ii) if $x \in R_N$, $a \in R$, $n \in N$ such that $x = an^{-1}$.

Such an N is called a right Ore semigroup and R_N is called a partial ring of quotients of R with respect to N. R_N is a ring of quotients of R in the sense of Utumi [13] wo we may suppose $R_N \subseteq Q_{max}(R)$. If $N = C_R(0)$ we say R has a classical right quotient ring and write $R_N = Q_{cl}(R)$ in this case. If $Q_{cl}(R)$ exists, we say that R is a right order in $Q_{cl}(R)$.

It is well known that the following are equivalent (see for example 3):

(a) $Q_{c\ell}(R)$ exists and is semisimple

(b) R is semiprime right Goldie

(c) R is semiprime, right finite dimensional and $Z_r(R) = 0$.

Then $Q_{c\ell}(R) = Q_{max}(R)$.

The next theorem is a direct consequence of 1.5 and 1.6 and the equivalence (a) $\Leftrightarrow$ (c) above.

Theorem 2.1: Let $S = R[x_\alpha]$. Then R is a right order in a semisimple ring if and only if S is such an order.

The Artin-Wedderburn Theorems now guarantee that under the hypotheses of 2.1, $Q_{c\ell}(R)$ and $Q_{c\ell}(S)$ are finite direct sums of matrix rings over division rings. We proceed to investigate the connections between these matrix rings.

Lemma 2.2: Let D be a division ring. Then

$$Q_{max}(D[x_\alpha]) = Q_{c\ell}(D[x_\alpha]) \text{ is a division ring.}$$

Proof: By Lemma 1.2, $S = D[x_\alpha]$ is a domain, and is right Ore with respect to $C_S(0)$, by 2.1. The assertion now follows.

Lemma 2.3: (i) If R and T are rings with $R \simeq T$, then $R[x_\alpha] \simeq T[x_\alpha]$.

(ii) If $\{R_i\}$ $1 \leq i \leq n$, is a family of rings

(a) $\left(\bigoplus_{i=1}^{n} R_i\right)[x_\alpha] \simeq \bigoplus_{i=1}^{n} (R_i[x_\alpha])$ and

(b) $Q_{max}(\bigoplus_{i=1}^{n} R_i) = \bigoplus_{i=1}^{n} Q_{max}(R_i)$

(iii) If R_n is the nxn matrix ring over R,

(a) $R_n[x_\alpha] \simeq (R[x_\alpha])_n$

(b) $Q_{max}(R_n) = (Q_{max}(R))_n$

Proof: The isomorphisms in (i), (ii) and (iii) are all natural. (ii)(b) and (iii)(b) are due to Utumi (Theorems 2.1 and 2.3 of [13]).

Lemma 2.4: Let N be a right Ore semigroup of R. Then N is a right Ore semigroup of S and

$$S_N = R_N[x_\alpha]$$

Proof: Since $C_R(0) \subseteq C_S(0)$, N is a multiplicatively closed subset of S. Each $n \in N$ is a unit in $R_N \subseteq R_N[x_\alpha]$.

If $\tilde{f} \in R_N[x_\alpha]$, then there exists $f \in S$ and $n \in N$ such that $\tilde{f} = fn^{-1}$. To see this let $\tilde{f} = \sum_{i=1}^{n} \tilde{f}_i m_i$ where $\tilde{f}_i \in R_N$ for each i.

For $1 \leq i \leq n$, $\exists\ f_i \in R$, $n_i \in N$ such that $\tilde{f}_i = f_i n_i^{-1}$.

By the common multiple property (see for example, Lemma 1.4 of [10]), $\exists\ n \in N$ such that $n_i^{-1} n \in R$ for every i. So $\tilde{f}n = f \in S$ and $\tilde{f} = fn^{-1}$ where $f \in S$ and $n \in N$. It readily follows that N is a right Ore semigroup of S and $S_N = R_N[x_\alpha]$.

Theorem 2.5: Let R be a semiprime right Goldie ring. If

$$Q_{c\ell}(R) = \bigoplus_{i=1}^{k} D_{n_i}^{(i)}$$

where $D^{(i)}$ is a division ring for all i, then

$$Q_{c\ell}(S) \simeq \bigoplus_{i=1}^{k} (Q_{c\ell}(D^{(i)}[x_\alpha]))_{n_i} .$$

Proof: Let $T^{(i)} = Q_{c\ell}(D^{(i)}[x_\alpha]) = Q_{max}(D^{(i)}[x_\alpha])$, $1 \leq i \leq k$. By 2.2, $T^{(i)}$ is a division ring, for each i.

Let $N = C_R(0)$. Then

$$R_N = Q_{c\ell}(R) = \bigoplus_{i=1}^{k} D_{n_i}^{(i)} \text{ exists so by 2.4}$$

$$S_N = R_N[x_\alpha] = (\bigoplus_{i=1}^{k} D_{n_i}^{(i)})[x_\alpha] .$$

By 2.3, $S_N \simeq \bigoplus_{i=1}^{k} (D^{(i)}[x_\alpha])_{n_i}$.

Again by 2.3

$$Q_{max}(S_N) \simeq \bigoplus_{i=1}^{k} (Q_{max}(D^{(i)}[x_\alpha]))_{n_i} = \bigoplus_{i=1}^{k} T_{n_i}^{(i)}$$

Since S_N is a ring of quotients of S,

$$Q_{c\ell}(S) = Q_{max}(S) = Q_{max}(S_N) \simeq \bigoplus_{i=1}^{k} T_{n_i}^{(i)}$$

as asserted.

Next we determine whether

$$Q_{c\ell}(S) = \{ fg^{-1} \mid f \in S,\ g \in C_S(0) \}$$

can be described completely in terms of R. Evidently this can be done if all the regular elements of $S = R[x_\alpha]$ can be found, given R, and we have not been able to do so in general. However, $Q_{c\ell}(S)$ can still be obtained by localizing at a known set of regular elements.

Let $M = \{g \in S \mid L(g) \in C_R(0)\}$.

By 1.2, M is a multiplicatively closed subset of $C_S(0)$.

Lemma 2.6: If R is a semiprime right Goldie ring and $g \in C_S(0)$, then $\exists\ g' \in S$ such that $gg' \in M$.

Proof: For any right ideal B' of S, let

$$B = \{L(f) \mid f \in B'\}$$

Then B is a right ideal of R. If B' is an essential right ideal of S, then B is an essential right ideal of R. To see this, let $0 \neq A \in L_r(R)$. Then $0 \neq \omega(A) \in L_r(S)$ so $\omega(A) \cap B' \neq 0$. If $0 \neq f \in \omega(A) \cap B'$ we have $0 \neq L(f) \in A \cap B$.

Now let $B' = gS$ where $g \in C_S(0)$. Since S is right f.d., gS is an essential right ideal of S so that

$$B = \{L(gg') \mid g' \in S\}$$

is an essential right ideal of R. Since R is semiprime right Goldie, B contains a regular element. Hence $\exists\ g' \in S$ such that $gg' \in M$.

Theorem 2.7: Let R be a semiprime right Goldie ring. Then

$$Q_{c\ell}(S) = \{fg^{-1} \mid f \in S,\ g \in M\} = S_M.$$

Proof: Let $q \in Q_{c\ell}(S)$. Then $\exists\ g_1 \in C_S(0)$, $f_1 \in S$ such that $qg_1 = f_1 \in S$. By 2.6, $\exists\ g_1' \in S$ such that $g_1 g_1' \in M$. Let $g = g_1 g_1'$, $f = f_1 g_1'$. Then $q = fg^{-1}$ where $f \in S$, $g \in M$.

§3. ORDERS IN ARTINIAN RINGS: In this section, we prove the analog of Theorem 2.1 for orfers in Artinian rings (Theorem 3.6). Again we show that $Q_{c\ell}(S)$ can be obtained by localizing at a well described set of regular elements of S.

The rather cumbersome definition of a polynomial ring in §1 will be put to use in the next lemma, where it is observed that a polynomial ring has, among its rings of quotients, a group ring.

Lemma 3.1: If G and M are as in Defn. 1.1, then M is a right Ore semigroup of $S = R[x_\alpha]$. Moreover, $S_M = RG$, where S_M is the partial ring of quotients of S with respect to M.

Proof: For any $f \in M$, $L(f) = 1 \in C_R(0)$ so $f \in C_S(0)$ and $M \subseteq C_S(0)$. Clearly M is multiplicatively closed. Each $m \in M$ is a unit in G, hence in RG. Next we show that

$$\tilde{f} \in RG \Rightarrow \exists\ f \in S,\ m \in M \text{ such that } \tilde{f} = fm^{-1}.$$

For each $\alpha \in I$, define $\tilde{\alpha}: G \to Z$ (the integers) by $\tilde{\alpha}(g) = n \Leftrightarrow g = x_\alpha^n x_{\alpha_1}^{n_1}, \ldots, x_{\alpha_k}^{n_k}$. Let $\tilde{f} \in RG$ be any element.

Then $\exists\ f_i \in R,\ g_i \in G,\ 1 \leq i \leq n$ such that

$$\tilde{f} = \sum_{i=1}^{n} f_i g_i$$

For each $\alpha \in I$ let

$$n_\alpha = \max\{|\alpha(g_i)| \mid 1 \leq i \leq n\}.$$

Then $n_\alpha \geq 0$ for every α and $n_\alpha = 0$ for all but finitely many α. Let

$$m = \prod_{\alpha \in I} x_\alpha^{n_\alpha}.$$

Then $m \in M$ and $g_i m \in M$ for every i, $1 \leq i \leq n$. Hence

$$\tilde{fm} = \sum_{i=1}^{n} f_i g_i m = f \in S$$

and $\tilde{f} = fm^{-1}$, as required. It now follows that M is a right Ore semigroup of S and $RG = S_M$.

The next Lemma is due to Small (Lemma 1.7 of [9]).

Lemma 3.2: If N is a right Ore semigroup of a ring T and T_N is a right order in a right Artinian ring, then T is a right order in a right Artinian ring. In fact $Q_{c\ell}(T) = Q_{c\ell}(T_N)$.

A Quasi-Forbenius (QF) ring is a right Artinian right self-injective ring. For other characterizations of QF-rings, see [4].

Defn. 3.3: For the sake of brevity, say a ring T has P_2 (resp. has P_3) if T is a right order in a right Artinian ring (resp. QF-ring).

Lemma 3.4: Let G be a group which has a collection of subgroups $\{G_\beta \mid \beta$ is an ordinal$\}$ such that for all β

(a) G_β is a normal subgroup of $G_{\beta+1}$

(b) $G_{\beta+1}/G_\beta$ is either finite or cyclic, of which at most finitely many are finite, and

(c) $G_\eta = G$ for some ordinal η.

Then RG has P_2 (resp. P_3) if R has P_2 (resp. P_3).

Proof: For P_2 see Hughes [7] and for P_3, see Horn [6].

Theorem 3.5: If R has P_2 (resp. P_3), then $S = R[x_\alpha]$ has P_2 (resp. P_3).

Proof: Evidently, any free abelian group satisfies the conditions of Lemma 3.4. Hence if G is as in definition 1.1, then by 3.1, 3.2 and 3.4,

R has P_2 (resp. P_3) $\Rightarrow$ RG has P_2 (resp. P_3)
$\Rightarrow$ S_M has P_2 (resp. P_3)
$\Rightarrow$ S has P_2 (resp. P_3)

An ideal K of a ring T is said to be locally nilpotent if for every finite subset F of K, there is an integer $k = k(F)$ such that every product involving k elements from F, vanishes. The Levitzki radical of T, L(T), is the sum of all the locally nilpotent ideals of T, and is itself a locally nilpotent ideal. For the moment, let S be any ring, and let rad S = N(S). Small [11] has shown that S has P_2 if, and only if,

(i) L(S) is nilpotent

(ii) For each integer $k \geq 0$, $S_k = S/J_k$ is a right Goldie ring, where

$J_k = \ell_S(L(S)^k) \cap L(S)$ and $L(S)^0 = S$.

(iii) a is regular in S whenever a+N(S) is regular in $S/N(S)$.

Theorem 3.6: A ring R is a right order in the right Artinian ring if, and only if, $S = R[x_\alpha]_{\alpha \in I}$ is a right order in a right Artinian ring, for any index set I.

Proof: Sufficienty has been shown in 3.5. Suppose then that S has P_2. We show that R satisfies conditions (i), (ii) and (iii) above.

(i) L(R) is nilpotent: We first observe that

$\omega(L(R)) = L(R)[x_\alpha]$

is a locally nilpotent ideal of S. For let

$F = \{f^{(1)}, f^{(2)}, \ldots, f^{(n)}\} \subseteq \omega(L(R))$

For each i, $1 \leq i \leq n$, let

$$f^{(i)} = \sum_{j=1}^{n(i)} f_j^{(i)} m_{ij}$$

Then $F' = \{f_j^{(i)}\}_{i,j}$ is a finite subset of $L(R)$, which is locally nilpotent. Hence there is an integer k such that any product of k elements from F' vanishes. So any product of k elements from F vanishes, proving that $\omega(L(R)) \subseteq L(S)$. Let n be the index of nilpotency of $L(S)$. Then by 1.4,

$$0 = (\omega(L(R)))^n = \omega((L(R)^n) \text{ so } (L(R))^n = 0$$

(ii) $J_0 = \ell_S(L(S)^0) \cap L(S) = 0$ so $S_0 = S$ is a right Goldie ring. Since S satisfies the maximum condition on annihilator right ideals, we have by a result of Herstein and Small [5] that $L(S)$ contails all nil ideals. In particular $L(S) \supseteq N(S)$. Since $L(S)$ is nilpotent $L(S) = N(S)$. Since S is right Goldie, so is R by 1.5. By the same reasoning, $L(R) = N(R)$.

Let $I_K = \ell_R(N(R)^k) \cap N(R)$, $k \geq 0$.

Since $\omega(N(R)) = \omega(\text{rad } R) = \text{rad } S = N(S)$, by 1.6, we obtain by repeated use of 1.4 that

$$\omega(I_k) = J_k \text{ for every } k \geq 0$$

Let $R_k = R/I_k$. Then by 1.7,

$$R_k[x_\alpha] \cong S/J_k = S_k$$

Hence by 1.6, R_k is right Goldie for each $k \geq 0$-

(iii) Let $a+N(R)$ be regular in $R/N(R)$. Then $a+N(R)$ is regular $R/N(R)[x_\alpha]$. The map

$$\phi : S \to R/N(R)[x_\alpha]$$

defined by $\phi(\sum_{i=1}^{n} f_i m_i) = \sum_{i=1}^{n} (f_i+N(R))m_i$

induces an isomorphism between ${}^{S}/_{N(S)}$ and ${}^{R}/_{N(R)}[x_\alpha]$ in which $a+N(R)$ corresponds to $a+N(S)$. Hence $a+N(S)$ is regular in ${}^{S}/_{N(S)}$. Since S satisfies condition (iii), a is regular in S. In particular $a \in C_R(0)$.

This proves the theorem.

Theorem 3.7: Let R be a right order in a right Artinian ring, and let $S = R[x_\alpha]$. If

$$M = \bigcup_{h \in S} \{h+N(S) \mid L(h) \in C_R(0)\}$$

then M is a right Ore semigroup of S and $Q_{c\ell}(S) = S_M$.

Proof: It is well known that if a ring T is a right order in a right Artinian (indeed semiprimary) ring, then ${}^{T}/_{N(T)}$ is semiprime Goldie and $a \in C_T(0)$ if and only if $a+N(T)$ is regular in ${}^{T}/_{N(T)}$ (See for example, Prop. 3.1, p. 286 of [12]).

(i) $M \subseteq C_S(0)$: Let $f \in M$. Then $\exists\ h \in C_S(0)$ such that $f+N(S) = h+N(S)$. Since S is a right order in a right Artinian ring, $f \in C_S(0)$.

Clearly, M is multiplicatively closed.

(ii) $f \in C_S(0) \Rightarrow f' \in S$ such that $ff' \in M$: Let $\tilde{\pi}: S \to {}^{R}/_{N(R)}[x_\alpha]$ be as in Lemma 1.7 and let $\tilde{\pi}(g) = \bar{g}$ for each $g \in S$.

Let $f \in C_S(0)$. Then $f+N(S)$ is regular in ${}^{S}/_{N(S)}$. In the induced isomorphism ${}^{S}/_{N(S)} \cong {}^{R}/_{N(R)}[x_\alpha]$ the image $\bar{f}$ of $f+n(S)$ is regular in ${}^{R}/_{n(R)}[x_\alpha]$. By 2.6 and the fact that ${}^{R}/_{N(R)}$ is a semiprime right Goldie ring, $\exists\ f' \in S$ such that $L(\bar{f}\bar{f}')$ is regular in ${}^{R}/_{N(R)}$.

Let $\bar{f}\bar{f}' = \overline{ff'} = \sum_{i=1}^{n} (a_i + N(R))m_i$

where $a_n + N(R) = L(\overline{ff'})$.

Let $h = \sum_{i=1}^{n} a_i m_i \in S$.

Since $a_n \neq 0$, $L(h) = a_n$. But $a_n \in C_R(0)$ since $a_n + N(R)$ is regular in $R/N(R)$. By defn. of M, $h + N(S) \subseteq M$. But $\bar{h} = \overline{ff'}$ so

$$ff' \in h + N(R)[x_\alpha] = h + N(S),$$ proving $ff' \in M$ for some $f' \in S$.

We proceed as in 2.7 to complete the proof.

Corollary 3.8: If R is a right order in a right Artinian ring and $f \in C_S(0)$, then $\exists$ $f' \in S$ such that some coefficient of ff' is regular in R.

Remark: Theorem 3.5 cannot be extended to orders in Noetherian rings, even when R is commutative. The following example was kindly provided by Prof. Vasconcelos:

Let k be a field and let $k[[x,y,z]]$ be the power series ring in the indeterminates x,y,and z. Let $R = k[[x,y,z]]/(x^2,xy,xz)$ Then R is a commutative Noetherian ring so that its classical quotient ring is Noetherian (Theorem 1.9 of [2]). Let $Q = Q_{c\ell}(R[T_1,T_2,....])$. If f goes to $\bar{f}$ in the natural map

$$k[[x,y,z]] \to k[[x,y,z]]/(x^2,xy,xz),$$

the ideal in Q generated by

$$\{\bar{x}, \bar{y}T_1 - \bar{x}T_2, \bar{y}T_2 - \bar{z}T_3, ...\}$$

is not finitely generated, so Q is not Noetherian.

§4. ORDERS IN QUASI-FROBENIUS RINGS: Let M_R be an object of mod-R. If N_R is a submodule of M_R, we write $N_R \triangleleft M_R$ (resp. $N_R \blacktriangleleft M_R$) whenever M_R is an essential (resp. rational) extension of N_R. Thus

$N_R \triangleleft M_R \Leftrightarrow$ whenever $0 \neq m \in M$, $\exists\ r \in R$ such that $0 \neq mr \in N$ and

$N_R \blacktriangleleft M_R \Leftrightarrow$ whenever $0 \neq m_1 \in M$, $m_2 \in M$, $\exists\ r \in R$ such that

$$0 \neq m_1 r,\ m_2 r \in N.$$

The injective hull of M_R is designated by $E(M_R)$. If R is a subring of a ring T, then T is a ring of right quotients of R provided $R_R \blacktriangleleft T_R$. The maximal ring of right quotients $Q_{max}(R)$ of R is the biendomorphism ring of $E(R_R)$.

A characterization of $Q_{max}(R)$ we shall often use is

$$Q_{max}(R) = \{x \in E(R_R) \mid x^{-1}R \blacktriangleleft R_R\}$$

where $x^{-1}R$ is the right ideal

$$\{r \in R \mid xr \in R\}.$$

$Q_{max}(R)$ is a ring of right quotients of R containing every ring T of right quotients of R satisfying

$$R \subseteq T \subseteq E(R_R).$$

For details, refer to either [3] or [8].

In this section we show that a ring R is a right order in a QF-ring if and only if $S = R[x_\alpha]_{\alpha \in I}$ is such an order, for any index set I. (Theorem 4.5). To this end, we observe that if $Q_{max}(R[x_\alpha])$ is right self-injective, so is $Q_{max}(R)$ (Theorem 4.3).

Any object of mod-S is an object of mod-R via the inclusion map $R \to S$.

Lemma 4.1:

$$K_R \lhd R_R \Leftrightarrow (\omega(K))_R \lhd S_R \Leftrightarrow (\omega(K))_S \lhd S_S.$$

Proof: Suppose $K_R \lhd R_R$. If $0 \neq f \in S$, $g \in S$, then $\exists\ 0 \neq f_i \in R$ for some coefficient f_i of f. Let $g = \sum_{j=1}^{n} g_j m_j$. There is an $r \in R$ such that $0 \neq f_i r$ and $g_j r \in K$ for every j. Hence $0 \neq fr$ and $gr \in \omega(K)$, proving $(\omega(K))_R \lhd S_R$. Since $R \subset S$, it follows that $(\omega(K))_S \lhd S_S$. The implication $(\omega(K))_S \lhd S_S \Rightarrow K_R \lhd R_R$ is trivial.

Lemma 4.2: Let A be an object of mod-S. If A_S is injective, then so is A_R.

Proof: Let A_S be injective and let K be any right ideal of R. Let $\phi \in \mathrm{Hom}_R(K_R, A_R)$. Define $\phi^* : \omega(K) \to A_S$ by

$$\phi^*\left(\sum_{i=1}^{n} f_i m_i\right) = \sum_{i=1}^{n} \phi(f_i) m_i.$$

Then $\phi^* \in \mathrm{Hom}_S(\omega(K)_S, A_S)$.

Since A_S is injective, (by Baer's criterion for injective modules) $\exists$ an $a \in A$ such that $\phi^*(f) = af$ for every $f \in \omega(K)$. Hence for $k \in K$, $\phi(k) = \phi^*(k) = ak$, proving that A_R is injective.

Theorem 4.3: If $Q_{max}(S)$ is right self-injective, so is $Q_{max}(R)$.

Proof: Let $Q' = Q_{max}(S)$. Now Q' is right self-injective if, and only if, Q'_S is injective (Prop. 4.3.3 of [8]). By Lemma 4.2, Q'_R is injective. Since $R_R \subset S_R \subset Q'_R$, we may assume that $R_R \subseteq E(R_R) \subseteq Q'_R$. The assertion will follow if we can show that $Q_{max}(R) = E(R_R)$.

We first claim that:

"If $a_1, a_2, \ldots, a_n$ in $E(R_R)$ and $m_1, m_2, \ldots m_n$ in M satisfy $\sum_{i=1}^{n} a_i m_i = 0$, then $a_i = 0$ for every i." (*)

Let $\sum_{i=1}^{n} a_i m_i = 0$. Note that $E(R_R)$ is a subset of Q', which is an object of mod-S, so that $a_i m_i$ is defined for each $i, 1 \leq i \leq n$. Suppose that some $a_i \neq 0$. Since $E(R_R)$ is an essential extension of R_R, $\exists\ r \in R$ such that $f_i = a_i r \in R$ for each i, not all $f_i = 0$. But then $f = \sum_{i=1}^{n} f_i m_i = (\sum_{i=1}^{n} a_i m_i)r = 0 \in S$ so that $f_i = 0$ for every i, a contradiction.

Let $a \in E(R_R) \subseteq Q_{max}(S)$ Then

$$f = \sum_{i=1}^{n} f_i m_i \in \omega(a^{-1}R) \Rightarrow f_i \in a^{-1}R \text{ for every } i$$

$$\Leftrightarrow \quad af_i \in R \text{ for every } i$$

$$\Rightarrow \quad af = \sum_{i=1}^{n} (af_i)m_i \in S$$

$$\Leftrightarrow \quad af \in a^{-1}S.$$

Further if $af = \sum_{i=1}^{n} (af_i)m_i \in S$ we conclude from (*) and the fact that $af_i \in E(R_R)$ for each i, that $af_i \in R$ for each i. Hence $a^{-1}S = \omega(a^{-1}R)$. Since $a \in Q_{max}(S)$, $a^{-1}S \blacktriangleleft S_S$. By Lemma 4.1, $a^{-1}R \blacktriangleleft R_R$ proving $a \in Q_{max}(R)$. Hence $Q_{max}(R) = E(R_R)$ and $Q_{max}(R)$ is right self-injective.

Lemma 4.4: If a ring T of right quotients of R is right self-injective, that $T = Q_{max}(R)$ so that $Q_{max}(R)$ is itself right self-injective.

Proof: Let $E(R_R)$ be the injective hull of R_R. We may assume

$$R \subseteq T \subseteq U = Q_{max}(R) \subseteq E(R_R).$$

Since T_T is injective, there exists a submodule K_T of U_T such that

$$U_T = T_T \oplus K_T.$$

Since $R \subseteq T$, K is an R-submodule of $E(R_R)$ and $K \cap R \subseteq K \cap T = 0$. But $R_R \lhd E(R_R)$. Hence $K = 0$ and $T = U = Q_{max}(R)$.

Theorem 4.5: A ring R is a right order in QF-ring $\Leftrightarrow$ $S = R[x_\alpha]$ is such an order.

Proof: Sufficienty has been proved in 3.5. Let S be a right order in a QF-ring. Then $Q_{c\ell}(S)$ is a right self-injective right Artinian ring. Since $Q_{c\ell}(S)$ is a ring of right quotients of S, it follows from 4.4 that $Q_{c\ell}(S) = Q_{max}(S)$ is right self-injective. By Theorem 4.3 and its proof we may assume that

$$R \subseteq Q_{max}(R) = E(R_R) \subset Q_{max}(S) = Q_{c\ell}(S).$$

By 3.6, $Q_{c\ell}(R)$ exists and is right Artinian. We need only show that $Q_{c\ell}(R) = Q_{max}(R)$ to complete the proof. Let then $a \in Q_{max}(R)$. Since $a \in Q_{c\ell}(S)$, $\exists\ g \in C_S(0)$ such that $ag = f \in S$. By Corollary 3.8, $\exists\ g' \in S$ such that some coefficient d of gg' is in $C_R(0)$. But $g \in a^{-1}S$ so that $gg' \in a^{-1}S = \omega(a^{-1}R)$. Hence $d \in a^{-1}R \cap C_R(0)$ and $ad = r \in R$. Since $d^{-1} \in Q_{c\ell}(R)$ exists, $a = rd^{-1} \in Q_{c\ell}(R)$, proving that $Q_{c\ell}(R) = Q_{max}(R)$ is a QF-ring.

ACKNOWLEDGEMENTS: The author is deeply indebted to his thesis advisor Dr. R.W. Wilkerson for his constant advice and encouragement, and to Prof. Carl Faith for several stimulating conversations during the author's sabbatical year at Rutgers.

References

[1] Amitsur, S.A., Radicals of Polynomial Rings, Can. J. Math. 8 (1956), 355-361.

[2] Asano, K., Quotientbildung und Schiefringe, J.Math.Soc. Japan 1 (1949), 73-78.

[3] Faith, C., Lectures on Injective modules and Quotient Rings, Lecture Notes in Mathematics, vol. 49, Springer (1967).

[4] Faith, C., Algebra II, Ring Theory, Springer (1976).

[5] Herstein, I.N. and Small, L.W. Nil rings satisfying certain chain conditions, Canad. J. Math. 16 (1964), 771-776.

[6] Horn, A., Quasi-Frobenius quotient rings of Group Rings, J. Aust. Math. Soc. 20 (1975), 394-397.

[7] Hughes, I., Artinian Quotient rings of Group Rings, J. Aust. Math. Soc. 16 (1973), 379-384.

[8] Lambek, J., Lectures on Rings and Modules, Waltham, Mass., Blaisdell Publishing Co., (1966).

[9] Shock, R.C., Polynomial Rings over finite dimensional rings, Pac. J. Math. 42 (1972), 251-258.

[10] Small, L.W., Orders in Artinian rings, J. Alg. 4 (1966), 13-41.

[11] Small, L.W., Orders in Artinian rings, II, J. Alg. 9 (1968), 266-273.

[12] Stenstrom, B., Rings of Quotients, Springer (1975).

[13] Utumi, Y., On Quotient Rings, Osaka Math. J. 8 (1956), 1-18.

Rebut el 10 de novembre del 1983

Department of Mathematics
University of Durban-Westville
Private Bag X54001 Durban 4001
SOUTH AFRICA