

A SMALL CLOSED CONVEX PROJECTIVE 4-MANIFOLD VIA DEHN FILLING

GYE-SEON LEE, LUDOVIC MARQUIS, AND STEFANO RIOLO

Abstract: In order to obtain a closed orientable convex projective 4-manifold with small positive Euler characteristic, we build an explicit example of convex projective Dehn filling of a cusped hyperbolic 4-manifold through a continuous path of projective cone-manifolds.

2010 Mathematics Subject Classification: 22E40, 53A20, 53C15, 57M50, 57N16, 57S30.

Key words: real projective structure, hyperbolic 4-manifold, Dehn filling, Euler characteristic, cone-manifold, Hilbert geometry.

1. Introduction

Convex projective manifolds form an interesting class of aspherical manifolds, including complete hyperbolic manifolds. We refer to [10, 48, 20] and [51] for surveys on convex projective manifolds and hyperbolic 4-manifolds, respectively. This class of geometric manifolds has been studied notably in the context of deformations of geometric structures on manifolds or orbifolds (see the survey [20] and the references therein), or for its link to dynamical systems through the notion of Anosov representation [7, 30, 29] (see [43, 38] for the notion of Anosov representation).

A *convex projective n -manifold* is the quotient Ω/Γ of a properly convex¹ domain Ω in the real projective space \mathbb{RP}^n by a subgroup Γ of the projective linear group $\mathrm{PGL}_{n+1} \mathbb{R}$ acting freely and properly discontinuously on Ω . A *convex projective n -orbifold* is defined similarly without requiring the action of Γ to be free. If Ω is endowed with its Hilbert metric, then Γ acts on Ω by isometry, and the manifold (or orbifold) Ω/Γ inherits a complete Finsler metric (see [63] for an introduction to Hilbert geometry). In the case where Ω is an open ellipsoid, it is isometric to the hyperbolic space \mathbb{H}^n ,² and the quotient Ω/Γ is a complete hyperbolic

¹A subset Ω of \mathbb{RP}^n is *properly convex* if its closure $\overline{\Omega}$ is contained and convex in some affine chart.

²This is in fact the projective model of the hyperbolic space, also known as the Beltrami–Cayley–Klein model.

manifold (or orbifold). A complete hyperbolic manifold is *cusped* if it is non-compact and of finite volume. A *convex projective* (resp. *complete hyperbolic*) *structure* on a manifold M is a diffeomorphism between M and a convex projective (resp. hyperbolic) manifold Ω/Γ .

The goal of this paper is to prove the following:

Theorem A. *There exists a closed orientable convex projective 4-manifold X containing ten disjoint totally geodesic 2-tori $\Sigma = T_1 \sqcup \cdots \sqcup T_{10} \subset X$ such that:*

- (1) *The complement $M = X \setminus \Sigma$ admits a complete finite-volume hyperbolic structure.*
- (2) *The Euler characteristic of X (and of M) is 12.*
- (3) *The hyperbolic manifold M has a maximal cusp section in which each filling curve has length 6.*
- (4) *The fundamental group $\pi_1 X$ is relatively hyperbolic with respect to the collection of rank-2 abelian subgroups $\{\pi_1 T_i, \pi_1 T'_i\}_i$, where $\{T'_1, \dots, T'_{10}\}$ is another collection of disjoint, totally geodesic, 2-tori such that each T_i is transverse to each T'_j .*
- (5) *The hyperbolic structure σ_0 on M and the convex projective structure $\sigma_{2\pi}$ on X arise as limits of an analytic path $\theta \mapsto \sigma_\theta$ of projective cone-manifold structures on X , singular along Σ with cone angle $\theta \in (0, 2\pi)$.*
- (6) *For each integer $m \geq 1$, the structure $\sigma_{2\pi/m}$ is the underlying cone-manifold structure of a convex projective orbifold Ω_m/Γ_m . For $m \geq 2$, the group Γ_m is relatively hyperbolic with respect to the collection of rank-2 abelian subgroups $\{\pi_1 T_i\}_i$.*

We refer the reader to Remark 2.2 for the meaning of totally geodesic submanifold in the real projective setting and to Subsection 2.5 for the definition of relative hyperbolicity and some facts about it.

Note that the manifold X does not admit a hyperbolic structure because $\pi_1 X$ contains \mathbb{Z}^2 (the tori T_i and T'_i are indeed π_1 -injective). The hyperbolic manifold M has ten cusps, each with section diffeomorphic to the 3-torus. A *filling curve* is a closed geodesic in a cusp section of M (with respect to the induced flat metric) that bounds a disc in X .

Cone-manifolds and Dehn filling. Projective cone-manifolds are singular projective manifolds generalising the more familiar hyperbolic cone-manifolds (see Definition 2.1). The convex projective 4-manifold X of Theorem A is obtained from the cusped hyperbolic 4-manifold M by “projective Dehn filling”. This is in analogy with Thurston’s hyperbolic Dehn filling [61], where $\theta \mapsto \sigma_\theta$ is a path of hyperbolic cone-manifold structures on a 3-manifold X , singular along a link $\Sigma \subset X$. In both

hyperbolic and projective cases, as the cone angle θ approaches 2π , the projective cone-manifold structure becomes non-singular, and we get a convex projective structure on X . At the other extreme of the path, as θ tends to 0, the singular locus Σ is drilled away, giving rise to the cusps of the hyperbolic manifold M .

The projective cone-manifold structures σ_θ of Theorem A are singular along the tori Σ , and induce (non-singular) projective structures³ on both $M = X \setminus \Sigma$ and Σ . The path $\theta \mapsto \sigma_\theta$ is *analytic*, meaning that for each $\theta \in (0, 2\pi)$ it is possible to choose a holonomy representation $\rho_\theta \in \text{Hom}(\pi_1 M, \text{PGL}_5 \mathbb{R})$ of the projective structure $\sigma_\theta|_M$ so that the function $\theta \mapsto \rho_\theta(\gamma)$ is analytic for all $\gamma \in \pi_1 M$. Each torus $T_i \subset \Sigma$ has a meridian $\gamma_i \in \pi_1 M$ whose holonomy $\rho_\theta(\gamma_i) \in \text{PGL}_5 \mathbb{R}$ is conjugate to a (projective) rotation of angle θ . In addition, the sequence of representations $\{\rho_{2\pi/m}\}_m$ converges algebraically to ρ_0 as $m \rightarrow \infty$, and the sequence of convex sets $\{\overline{\Omega_m}\}_m$ converges to $\overline{\mathbb{H}^4} \subset \mathbb{R}\mathbb{P}^4$ in the Hausdorff topology.⁴

New features of the result. Theorem A is shown by an explicit construction. Since the convex projective manifold X has non-zero Euler characteristic, it is indecomposable⁵ (see Fact 2.15). It seems that, at the time of writing this paper, the literature misses concrete examples of closed indecomposable convex projective n -manifolds, $n \geq 4$, which do not admit a hyperbolic structure. For the moment, we know only two techniques to obtain such manifolds:

- (1) torsion-free subgroups of some discrete projective reflection groups using Vinberg's theory [65], as shown by Benoist [8, 9] and Choi and the first two authors [21, 22];
- (2) some Gromov–Thurston manifolds, as shown by Kapovich [40].

In contrast with Theorem A, the Selberg lemma has an important role to guarantee the existence of such manifolds in all these cases. In particular, very little is known about the topology of closed convex projective manifolds. Note that the techniques involved in the construction of our X are in the spirit of (1) rather than (2).

Remark 1.1. There is a clear distinction between the manifolds constructed in [9, 40] and the ones in [8, 21, 22], including our X : the fundamental groups of the former are Gromov-hyperbolic, but those of the latter are not.

³A (real) projective structure on an n -manifold is a $(\text{PGL}_{n+1} \mathbb{R}, \mathbb{R}\mathbb{P}^n)$ -structure.

⁴These additional facts can be proved as in [21, §12].

⁵A properly convex domain Ω of $\mathbb{R}\mathbb{P}^n$ is *indecomposable* if it is not a convex hull of lower-dimensional domains. A convex projective manifold or orbifold Ω/Γ is *indecomposable* if Ω is indecomposable.

The Euler characteristic of a closed even-dimensional manifold can be seen as a rough measure of its topological complexity. Note that a well-known conjecture states that closed aspherical 4-manifolds have Euler characteristic $\chi \geq 0$, and this is certainly true in the hyperbolic case by the Gauß–Bonnet theorem. Our manifold X has $\chi(X) = 12$, and appears to be the closed orientable indecomposable convex projective 4-manifold with the smallest known Euler characteristic (to the best of our knowledge). In the hyperbolic case, the smallest known value of χ is 16 [23, 46].⁶

Theorem A is an effective version, in dimension 4, of a result by Choi and the first two authors [21, Theorem B]. Let us first recall their construction, called “convex projective generalised Dehn filling”. They build a sequence of discrete projective reflection groups $\{\Gamma_m\}_{m \geq m_0}$ of $\mathrm{PGL}_{n+1} \mathbb{R}$, each acting cocompactly on a properly convex domain $\Omega_m \subset \mathbb{R}\mathbb{P}^n$ of dimension $n = 4, 5$, or 6, whose limit as $m \rightarrow \infty$ is a discrete hyperbolic reflection group $\Gamma_\infty < \mathrm{Isom}(\mathbb{H}^n)$ of finite covolume. A fundamental domain of the group Γ_m is a compact Coxeter polytope P_m in Ω_m whose combinatorics does not depend on m . The hyperbolic Coxeter polytope P_∞ , instead, is combinatorially obtained from P_m by substituting a ridge with an ideal vertex. In other words, the cusp of the hyperbolic orbifold $\mathbb{H}^n/\Gamma_\infty$ is “projectively filled”. By applying a refined version of Selberg’s lemma to Γ_∞ , they get a statement similar to Theorem A(6). The difference is that $X = \Omega_{m_0}/\Gamma'_{m_0}$ is “only” an orbifold (where Γ'_{m_0} is a finite-index subgroup of Γ_{m_0}). To promote X to a manifold, one should then apply the Selberg lemma again, this time to Γ'_{m_0} . Thus, our improvement is twofold: we found an X with small Euler characteristic, and a continuous (rather than discrete) family of cone-manifolds.

Another interesting feature of X is the relative hyperbolicity of $\pi_1 X$. Indeed, Gromov and Thurston ([12, 2])⁷ have shown that the fundamental group of a Dehn filling of a hyperbolic manifold with torus cusps is relatively hyperbolic with respect to the subgroups associated with the inserted tori, provided that the filling curves are longer than 2π .⁸ The

⁶Similarly to the orientable hyperbolic 4-manifolds that one gets from [23, 46], our manifold X is built as the orientable double cover of a non-orientable convex projective manifold. So the smallest known value of χ for a closed indecomposable convex projective (resp. hyperbolic) 4-manifold is currently 6 (resp. 8).

⁷A proof of the Gromov–Thurston 2π theorem was given by Bleiler and Hodgson [12] in the context of 3-manifolds, and the same proof holds in any dimension, as explained in [2, Section 2.1].

⁸See also [56, 37, 32] for the geometric group theoretic generalisation of that statement.

fundamental group of X is not relatively hyperbolic with respect to the subgroups associated with the inserted tori by Theorem A(4) (see Subsection 2.5).⁹ There is no contradiction between the Gromov–Thurston 2π theorem and Theorem A(3) because $2\pi > 6$.

Divisible convex domains. Recall that a properly convex domain Ω of \mathbb{RP}^n is *divisible* (by Γ) if there exists a discrete subgroup Γ of $\mathrm{PGL}_{n+1}\mathbb{R}$ acting cocompactly on Ω . A theorem of Benoist ([7]) implies that the indecomposable divisible convex domains $\Omega_m \subset \mathbb{RP}^4$ of Theorem A(6) are not strictly convex¹⁰ because the groups Γ_m of Theorem A(6) are not Gromov-hyperbolic.

Currently there are very few known constructions of inhomogeneous indecomposable divisible non-strictly convex domains. For a complete historical account, we refer the reader to the introduction of [21]. Here we mention only its essentials.

The first construction of such domains is due to Benoist [8], and has been extended in [47, 4, 22]. In those constructions the compact quotient Ω/Γ is homeomorphic to the union along the boundaries of finitely many submanifolds, each admitting a complete finite-volume hyperbolic structure in its interior. As a result, if Ω is of dimension n , then Γ is relatively hyperbolic with respect to a collection of virtually abelian subgroups of rank $n - 1$.

In [21], a different construction of inhomogeneous indecomposable divisible non-strictly convex domains is given by convex projective generalised Dehn filling. In contrast with the previous examples, these are relatively hyperbolic with respect to a collection of virtually abelian subgroups of rank $n - 2$. The divisible (by Γ_m) domains $\Omega_m \subset \mathbb{RP}^4$ of Theorem A(6) are new examples of this kind.

We point out that at the time of writing there is no example of inhomogeneous indecomposable divisible non-strictly convex domain of dimension n , for any $n \geq 9$.

We stress that such domains, to all appearances, are linked to the geometrisation problem, i.e. putting a (G, X) -structure on a manifold. So far, almost all manifolds geometrised through this process are either obtained by gluing cusped hyperbolic manifolds, or by Dehn filling of a cusped hyperbolic manifold. Here, the goal is to do so with a small manifold and using cone-manifolds. It is especially important that we do not use Selberg’s lemma.

⁹It is relatively hyperbolic with respect to a larger family of abelian groups.

¹⁰A subset Ω of \mathbb{RP}^n is *strictly convex* if it is properly convex and its boundary does not contain any non-trivial projective line segment.

Dehn fillings of hyperbolic manifolds. Let us say that a closed manifold X is a *filling* of a manifold M if there exists a codimension-2 submanifold $\Sigma \subset X$ such that the complement $X \setminus \Sigma$ is diffeomorphic to M . Note that the manifold M is diffeomorphic to the interior of a compact manifold \overline{M} whose boundary $\partial\overline{M}$ fibres in circles over Σ . Given M as above, we obtain a filling by attaching to \overline{M} the total space of a D^2 -bundle $E \rightarrow \Sigma$ through a diffeomorphism $\partial\overline{M} \rightarrow \partial E$. This operation is commonly called a *Dehn filling* of M . Any cusped hyperbolic manifold M has a finite covering M' with torus cusps (see e.g. [54, Theorem 3.1]). In other words, $\partial\overline{M}'$ consists of $(n-1)$ -tori. Typically the manifold M' has infinitely many fillings up to diffeomorphism.

Thurston's hyperbolic Dehn filling theorem states that every filling of a cusped hyperbolic 3-manifold with torus cusps, except for finitely many fillings on each cusp, admits a hyperbolic structure. In dimension $n \geq 4$, except for finitely many fillings on each cusp, the fundamental group of a filling of a cusped hyperbolic n -manifold is relatively hyperbolic with respect to a collection of subgroups virtually isomorphic to \mathbb{Z}^{n-2} , by [56, Theorem 1.1]. Since $n \geq 4$, those groups contain \mathbb{Z}^2 and so those fillings do not admit any hyperbolic structure. The geometry of the remaining fillings is rather unpredictable, but it is expected that they also do not carry a hyperbolic structure. But Theorem A(1) and [21, Theorem B] show that some fillings of some cusped hyperbolic n -manifolds admit a convex projective structure. This leads to the following:

Question 1.2. *Which filling of a cusped hyperbolic manifold of dimension $n \geq 4$ (with torus cusps) admits a convex projective structure?*

It is worth mentioning that almost all fillings of any cusped hyperbolic manifold with torus cusps admit a complete Riemannian metric of non-positive sectional curvature by the Gromov–Thurston 2π theorem [12, 2], and an Einstein metric of negative scalar curvature by work of Anderson [2] and Bamler [3]. Both, in some sense, extend Thurston's 3-dimensional theorem to higher dimension (compare also with [59, 32, 33]). But those theorems cannot be applied to the manifold X of Theorem A, since the filling curves are too short.

Let us also note that there is an opportune version of hyperbolic Dehn filling in dimension 4: one can sometimes fill some cusps of a hyperbolic 4-manifold and get another cusped hyperbolic 4-manifold, at the expense of drilling some totally geodesic surfaces [52, 45].

Projective flexibility. Thurston's hyperbolic Dehn filling theorem essentially relies on the flexibility of the complete hyperbolic structure of cusped hyperbolic 3-manifolds. The local rigidity theorem of Garland

and Raghunathan [34] (see also [11]), on the other hand, says that the holonomy representation ρ of any cusped hyperbolic manifold M of dimension $n \geq 4$ has a neighbourhood in $\text{Hom}(\pi_1 M, \text{Isom}(\mathbb{H}^n))$ consisting of conjugates of ρ . Now, Theorem A(5) shows that the hyperbolic structure σ_0 on the 4-manifold M is *projectively flexible*, i.e. the conjugacy class of the holonomy representation ρ_0 of σ_0

$$[\rho_0] \in \text{Hom}(\pi_1 M, \text{PGL}_5 \mathbb{R}) / \text{PGL}_5 \mathbb{R}$$

is not an isolated point. To avoid confusion with the terminology, we mention that in this definition of flexibility there is no restriction on the holonomy of the peripheral subgroups of $\pi_1 M$. For instance, all cusped hyperbolic 3-manifolds are projectively flexible by the hyperbolic Dehn filling theorem. It is thus natural to ask the following question, which is a priori different from Question 1.2.

Question 1.3. *Which cusped hyperbolic 4-manifold is projectively flexible? Is every cusped hyperbolic 4-manifold finitely covered by a projectively flexible one?*¹¹

We note that some cusped hyperbolic 4-orbifolds, for example the Coxeter pyramid [6, 3, 3, 3, ∞], are not projectively flexible [65, Proposition 20].



FIGURE 1. The Coxeter diagram of the pyramid [6, 3, 3, 3, ∞] (see Subsection 2.3 for the basic terminology on Coxeter groups). The associated cusped hyperbolic 4-orbifold is projectively rigid.

On the proof. As already said, the proof of Theorem A is constructive. We begin with the ideal hyperbolic rectified 4-simplex $R \subset \mathbb{H}^4$, which is a Coxeter polytope. By applying the techniques introduced in [21], we perform a “convex projective generalised Dehn filling” on R : the hyperbolic structure on the orbifold R is deformed to projective structures which extend to “mirror polytope” structures (see Subsection 2.2) on the bitruncated 4-simplex Q (see Subsection 2.6). Note that Q minus some ridges is stratum-preserving homeomorphic to R . This will be translated into the fact that X minus some tori is homeomorphic to M .

We build the hyperbolic manifold M as an orbifold covering of R , by exploiting a construction by Kolpakov and Slavich [42]. By lifting the deformation from R to M , we get the path $\theta \mapsto \sigma_\theta$. The manifold X covers a Coxeter orbifold based on the bitruncated 4-simplex Q .

¹¹Similar considerations were made by Cooper, Long, and Thistlethwaite [25] for closed hyperbolic 3-manifolds.

Since there exist cusped orientable hyperbolic 4-manifolds M_0 tessellated by copies of R with $\chi(M_0) < 12$ [42, 60, 58, 41], one could wonder why a smaller convex projective manifold X should not be built by Dehn filling such an M_0 . A first obstruction is topological: a cusp section of those M_0 does not always fibre in circles. Even when all cusp sections of such an M_0 do fibre in circles, M_0 does not cover R but covers the quotient of R by its symmetry group. The latter is the Coxeter pyramid $[6, 3, 3, 3, \infty]$ [58, Lemma 2.2], which is projectively rigid, so our technique does not apply in these cases.

Structure of the paper. In Section 2 we introduce some basic concepts of projective cone-manifolds, mirror polytopes, and the truncation process of the 4-simplex, and in Section 3 we prove Theorem A.

Acknowledgements. We would like to thank Caterina Campagnolo, Roberto Frigerio, Jason Manning, Marco Moraschini, and Andrea Seppi for interesting discussions. Part of this work was done when the second and third authors visited the mathematics department of Heidelberg University, which we thank for the hospitality. We also thank the referees for their time and their valuable advice and comments.

G.-S. Lee was supported by the European Research Council under ERC-Consolidator Grant 614733 and by DFG grant LE 3901/1-1 within the Priority Programme SPP 2026 “Geometry at Infinity”, and he acknowledges support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network). L. Marquis acknowledges the Centre Henri Lebesgue ANR-11-LABX-0020 LEBESGUE for its support. S. Riolo was supported by the mathematics department of the University of Pisa (research fellowship “Deformazioni di strutture iperboliche in dimensione quattro”), and by the Swiss National Science Foundation (project no. PP00P2-170560).

2. Preliminaries

In this section we introduce some preliminary notions and fix some notation.

2.1. Projective cone-manifolds. Riemannian (G, X) cone-manifolds were introduced by Thurston [62] (see also [53]). If the geometry (G, X) locally embeds in real projective geometry $(\mathrm{PGL}_{n+1} \mathbb{R}, \mathbb{RP}^n)$ (such as the constant-curvature geometries), a (G, X) cone-manifold can be thought of as a projective cone-manifold. Hyperbolic cone-manifolds of dimension 3 appear in the proofs of Thurston’s hyperbolic Dehn filling theorem [61] (see also [50, Chapter 15]) and of the orbifold theorem [24,

13]. Projective cone-manifolds were introduced by Danciger [27, 28] in the context of geometric transition from hyperbolic to anti-de Sitter 3-dimensional structures. Quite recently, some higher-dimensional cone-manifolds have been used, in particular, in dimension 4: for hyperbolic Dehn filling or degeneration [52, 45], and in the projective context of AdS-hyperbolic transition [57].

We now define projective cone-manifolds “with cone angles along link singularities”. Our definition is in the spirit of Barbot–Bonsante–Schlenker [5].

Let $\mathbb{S}^n = (\mathbb{R}^{n+1} \setminus \{0\})/\mathbb{R}_{>0}$ be the projective sphere and $\hat{\mathbb{S}}: \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{S}^n$ the canonical projection. For every subset U of \mathbb{R}^{n+1} , let $\mathbb{S}(U)$ denote $\hat{\mathbb{S}}(U \setminus \{0\})$. With a little abuse of notation, we embed the projective spheres \mathbb{S}^{n-2} and \mathbb{S}^1 into \mathbb{S}^n , $n \geq 2$, as follows:

$$\begin{aligned} \mathbb{S}^{n-2} &= \mathbb{S}(\{(x_1, \dots, x_{n-1}, 0, 0)\}) \subset \mathbb{S}^n \quad \text{and} \\ \mathbb{S}^1 &= \mathbb{S}(\{(0, \dots, 0, x_n, x_{n+1})\}) \subset \mathbb{S}^n. \end{aligned}$$

Given an open subset $U \subset \mathbb{S}^1$, we define

$$\mathbb{S}^{n-2} * U = \bigcup_{p \in \mathbb{S}^{n-2}} \bigcup_{q \in U} [p, -p]_q \subset \mathbb{S}^n,$$

where $[p, -p]_q \subset \mathbb{S}^n$ denotes the half circle containing q with endpoints p and its antipode $-p$. For example, $\mathbb{S}^{n-2} * \mathbb{S}^1 = \mathbb{S}^n$.

A *projective circle* is a closed connected $(\mathrm{SL}_2 \mathbb{R}, \mathbb{S}^1)$ -manifold. Let C be a projective circle. We may think of C as

$$C = \left(\prod_{\alpha \in A} U_\alpha \right) / \sim$$

for a collection $\{U_\alpha\}_{\alpha \in A}$ of open subsets of \mathbb{S}^1 , a collection $\{U_{\alpha\beta}\}_{\alpha, \beta \in A}$ of open subsets $U_{\alpha\beta} \subset U_\alpha$, a collection $\{g_{\alpha\beta}\}_{\alpha, \beta \in A}$ of diffeomorphisms $g_{\alpha\beta}: U_{\alpha\beta} \rightarrow U_{\beta\alpha}$ which are restrictions of elements of $\mathrm{SL}_2 \mathbb{R}$, and the relation $x \sim g_{\alpha\beta}(x)$ for all $x \in U_{\alpha\beta}$.

We now add the extra requirement that $C = C_\theta$ is an *elliptic circle*, i.e. that the holonomy representation ρ of C_θ sends a generator γ of $\pi_1 C_\theta$ to an elliptic element $\rho(\gamma) \in \mathrm{SL}_2 \mathbb{R}$. Going on to the universal covering $\widetilde{\mathbb{S}^1}$ of \mathbb{S}^1 and the covering group $\widetilde{\mathrm{SL}_2 \mathbb{R}}$ of $\mathrm{SL}_2 \mathbb{R}$ which acts on $\widetilde{\mathbb{S}^1}$, we lift ρ to a representation $\tilde{\rho}: \pi_1 C_\theta \rightarrow \widetilde{\mathrm{SL}_2 \mathbb{R}}$. To the element $\tilde{\rho}(\gamma) \in \widetilde{\mathrm{SL}_2 \mathbb{R}}$ we naturally associate a unique real number $\theta > 0$ which characterises the elliptic circle C_θ . Note that $\rho(\gamma)$ is conjugate to $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ (see [35, Section 5.4]).

By extending $g_{\alpha\beta} \in \mathrm{SL}_2 \mathbb{R}$ to $\hat{g}_{\alpha\beta} = \begin{pmatrix} \mathrm{Id} & 0 \\ 0 & g_{\alpha\beta} \end{pmatrix} \in \mathrm{SL}_{n+1} \mathbb{R}$ (thus fixing $\mathbb{S}^{n-2} \subset \mathbb{S}^n$ pointwise), we can define

$$\mathbb{S}^{n-2} * C_\theta = \left(\coprod_{\alpha \in A} \mathbb{S}^{n-2} * U_\alpha \right) / \sim$$

with the relation $x \sim \hat{g}_{\alpha\beta}(x)$ for all $x \in \mathbb{S}^{n-2} * U_{\alpha\beta}$.

This space will be the local model for our cone-manifolds. By canonically embedding \mathbb{S}^{n-2} and C_θ into $\mathbb{S}^{n-2} * C_\theta$, we have that the couple $(\mathbb{S}^{n-2} * C_\theta, \mathbb{S}^{n-2})$ is homeomorphic to $(\mathbb{S}^n, \mathbb{S}^{n-2})$. Moreover, the sphere $\mathbb{S}^{n-2} * C_\theta$ is stratified in two projective manifolds:

- the *singular locus* \mathbb{S}^{n-2} , and
- the *regular locus* $(\mathbb{S}^{n-2} * C_\theta) \setminus \mathbb{S}^{n-2}$.

The holonomy of a meridian¹² of the singular locus in the regular locus is the holonomy of a generator of $\pi_1 C_\theta$.

Definition 2.1. Let X be an n -manifold and $\Sigma \subset X$ a codimension-2 submanifold. A *projective cone-manifold structure* on X , singular along Σ with cone angles, is an atlas for X , each chart of which has values in some $\mathbb{S}^{n-2} * C_\theta$ and sends the points of Σ to the singular locus \mathbb{S}^{n-2} , and whose transition functions restrict to isomorphisms of projective manifolds on each stratum.

The *regular locus* of the cone-manifold X is the complement $X \setminus \Sigma$, while Σ is the *singular locus*. Both are (non-singular) projective manifolds. To each connected component Σ_i of Σ we associate a cone angle $\theta_i > 0$.

Remark 2.2. A projective cone-manifold X with all cone angles $\theta = 2\pi$ is simply a projective manifold with a totally geodesic codimension-2 submanifold $\Sigma \subset X$. Here, by *totally geodesic* we mean that the preimage of Σ under the universal covering of X is locally mapped to codimension-2 projective subspaces of \mathbb{S}^n . If moreover X is a convex projective (and so Finsler) manifold, then Σ is totally geodesic in the usual sense.

2.2. Mirror polytopes. We now introduce the main objects for our proof of Theorem A: mirror polytopes. Roughly speaking, a mirror polytope is a polytope in the projective sphere, together with a choice of a projective reflection along each of the supporting hyperplanes of the facets (satisfying some extra conditions). We refer to [21, 65] for further details.

¹²Let N be a manifold and $S \subset N$ a connected submanifold of codimension 2. We call a *meridian* of S an element $\gamma \in \pi_1(N \setminus S)$ that is represented by a curve which is freely homotopic in $N \setminus S$ to the boundary of a fibre of a tubular neighbourhood of S in N .

A subset P of \mathbb{S}^n is *convex* if there exists a convex cone¹³ U of \mathbb{R}^{n+1} such that $P = \mathbb{S}(U)$, and moreover a convex subset P is *properly convex* if its closure \overline{P} does not contain a pair of antipodal points. A *projective n -polytope* is a properly convex subset P of \mathbb{S}^n such that P has a non-empty interior and

$$P = \bigcap_{i=1}^N \mathbb{S}(\{v \in \mathbb{R}^{n+1} \mid \alpha_i(v) \leq 0\}),$$

where $\alpha_i, i = 1, \dots, N$, are linear forms on \mathbb{R}^{n+1} . We always assume that the set of linear forms is *minimal*, i.e. none of the half-spaces $\mathbb{S}(\{v \in \mathbb{R}^{n+1} \mid \alpha_i(v) \leq 0\})$ contains the intersection of all the others, hence the polytope P has N facets. A *facet* (resp. *ridge*) of a polytope is a face of codimension 1 (resp. 2).

A *projective reflection* is an element of $\mathrm{SL}_{n+1}^{\pm} \mathbb{R}$ of order 2 which is the identity on a projective hyperplane. Each projective reflection σ can be written as:

$$\sigma = \mathrm{Id} - \alpha \otimes b, \quad \text{in other words} \quad \sigma(v) = v - \alpha(v)b \quad \forall v \in \mathbb{R}^{n+1},$$

where α is a linear form on \mathbb{R}^{n+1} and b is a vector in \mathbb{R}^{n+1} such that $\alpha(b) = 2$. The projective hyperplane $\mathbb{S}(\ker(\alpha))$ is called the *support* of σ . A *projective rotation* is an element of $\mathrm{SL}_{n+1} \mathbb{R}$ which is the identity on a subspace $H \subset \mathbb{R}^{n+1}$ of codimension 2 and whose induced linear map from \mathbb{R}^{n+1}/H to itself is conjugate to a matrix $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ with $0 < \theta < 2\pi$. The real number θ is called the *angle* of rotation.

Definition 2.3. A *mirror polytope* is a pair consisting of a projective polytope P of \mathbb{S}^n and a finite collection of projective reflections $\{\sigma_s = \mathrm{Id} - \alpha_s \otimes b_s\}_{s \in S}$ with $\alpha_s(b_s) = 2$ such that:

- The index set S consists of all the facets of P .
- For each facet $s \in S$, the support of σ_s is the supporting hyperplane of s .
- For every pair of distinct facets s, t of P , $\alpha_s(b_t)$ and $\alpha_t(b_s)$ are either both negative or both zero.

Remark 2.4. The third item of Definition 2.3 may seem a bit awkward at first glance. In fact, [65, Proposition 6] shows that the third item holds when the group Γ generated by $\{\sigma_s\}_{s \in S}$ satisfies the condition:

$$\gamma \mathrm{Int}(P) \cap \mathrm{Int}(P) = \emptyset, \quad \forall \gamma \in \Gamma \setminus \{\mathrm{Id}\},$$

where $\mathrm{Int}(P)$ denotes the interior of P .

¹³By a *cone* we mean a subset of \mathbb{R}^{n+1} which is invariant under multiplication by positive scalars.

Note that the reflections σ_s of P determine the couples $\{(\alpha_s, b_s)\}_{s \in S}$ up to $\alpha_s \mapsto \lambda_s \alpha_s$ and $b_s \mapsto \lambda_s^{-1} b_s$ for $\lambda_s > 0$, because P is defined by $\alpha_i \leq 0$. Moreover, $\alpha_s(b_t) \alpha_t(b_s) = 4 \cos^2 \theta$ for some $\theta \in (0, \pi/2]$ if and only if the product $\sigma_s \sigma_t$ is a rotation of angle 2θ . The *dihedral angle* of a ridge $s \cap t$ of a mirror polytope is θ if $\sigma_s \sigma_t$ is a rotation of angle 2θ , and 0 otherwise. A *Coxeter polytope* is a mirror polytope whose dihedral angles are submultiples of π , i.e. each dihedral angle is π/m for some integer $m \geq 2$ or $m = \infty$.

For any number $N \in \mathbb{N}$, we set $\llbracket N \rrbracket := \{1, \dots, N\}$. An $N \times N$ matrix $A = (A_{ij})_{i,j \in \llbracket N \rrbracket}$ is *Cartan* if: (i) $A_{ii} = 2$ for all $i \in \llbracket N \rrbracket$, (ii) $A_{ij} \leq 0$ for all $i \neq j$, and (iii) $A_{ij} = 0 \Leftrightarrow A_{ji} = 0$ for all $i \neq j$. A Cartan matrix A is *irreducible* if it is not the direct sum of smaller matrices (after any re-ordering of the indices). Every Cartan matrix A is obviously the direct sum of irreducible Cartan matrices $A'_1 \oplus \dots \oplus A'_k$. Each irreducible Cartan matrix A'_i , $i = 1, \dots, k$, is called a *component* of A . If $x = (x_1, \dots, x_N)$ and $y = (y_1, \dots, y_N) \in \mathbb{R}^N$, we write $x > y$ if $x_i > y_i$ for all $i \in \llbracket N \rrbracket$, and $x \geq y$ if $x_i \geq y_i$ for all $i \in \llbracket N \rrbracket$.

Proposition 2.5 (Vinberg [65, Theorem 3]). *If A is an irreducible Cartan matrix of size $N \times N$, then exactly one of the following holds:*

- (P) (i) *The matrix A is non-singular, and (ii) for every $x \in \mathbb{R}^N$, if $Ax \geq 0$, then $x > 0$ or $x = 0$.*
- (Z) (i) *The rank of A is $N - 1$, (ii) there exists a vector $y \in \mathbb{R}^N$ such that $y > 0$ and $Ay = 0$, and (iii) for every $x \in \mathbb{R}^N$, if $Ax \geq 0$, then $Ax = 0$.*
- (N) (i) *There exists a vector $y \in \mathbb{R}^N$ such that $y > 0$ and $Ay < 0$, and (ii) for every $x \in \mathbb{R}^N$, if $Ax \geq 0$ and $x \geq 0$, then $x = 0$.*

We say that A is of positive, zero, or negative type if (P), (Z), or (N) holds, respectively.

A Cartan matrix A is of *positive* (resp. *zero*) *type* if every component of A is of positive (resp. zero) type. The *Cartan matrix* of a mirror polytope P is the matrix $A_P = (\alpha_s(b_t))_{s,t \in S}$. Note that the Cartan matrix of P is well defined up to conjugation by a positive diagonal matrix because the reflections σ_s of P determine the couples $\{(\alpha_s, b_s)\}_{s \in S}$ up to $\alpha_s \mapsto \lambda_s \alpha_s$ and $b_s \mapsto \lambda_s^{-1} b_s$ for $\lambda_s > 0$. Two Cartan matrices A and B are *equivalent* if $A = DBD^{-1}$ for some positive diagonal matrix D . We will make essential use of the following:

Theorem 2.6 ([65, Corollary 1]). *Let A be a Cartan matrix of size $N \times N$. If A is irreducible, of negative type, and of rank $n + 1$, then there exists a mirror polytope P of \mathbb{S}^n with N facets (unique up to the action of $SL_{n+1}^{\pm}(\mathbb{R})$) such that $A_P = A$.*

Remark 2.7. Theorem 2.6 is not explicitly stated in [65, Corollary 1] for non-Coxeter polytopes, but it follows from Propositions 13 and 15 of [65] that the consequent Corollary 1 of [65] is valid not only for Coxeter polytopes, but also for mirror polytopes.

To understand the combinatorics¹⁴ of a mirror polytope P with facets $\{s\}_{s \in S}$, we introduce the poset $\sigma(P) \subset 2^S$ partially ordered by inclusion, which is dual¹⁵ to the face poset of P :

$$\sigma(P) := \{T \subset S \mid T = \sigma(f) \text{ for some face } f \text{ of } P\},$$

where $\sigma(f) := \{s \in S \mid f \subset s\}$. For any subset $T \subset S$, we denote by A_T the restriction of the Cartan matrix A_P of P to $T \times T$.

Theorem 2.8 ([65, Theorems 4 and 7]). *Let P be a mirror n -polytope with facets $\{s\}_{s \in S}$ and with irreducible Cartan matrix A_P of negative type. Let T be a proper subset of S (i.e. $T \neq \emptyset, S$). Then:*

- (1) *If A_T is of positive type and $\sharp T = k$, then $T \in \sigma(P)$ and its corresponding face $\cap_{s \in T} s$ is of dimension $n - k$.*
- (2) *If A_T is of zero type and of rank $n - 1$, then $T \in \sigma(P)$ and the face $\cap_{s \in T} s$ is of dimension 0, i.e. a vertex of P .*

Remark 2.9. Theorem 2.8 is not explicitly stated in [65, Theorems 4 and 7], but it is obtained by applying [65, Theorem 4] as in the proof of [65, Theorem 7].

Remark 2.10. Theorem 2.8(1) tells us that for any mirror polytope P with facets $\{s\}_{s \in S}$ and reflections $\{\sigma_s = \text{Id} - \alpha_s \otimes b_s\}_{s \in S}$, if $\alpha_s(b_t)\alpha_t(b_s) < 4$, then the intersection $s \cap t$ is a face of codimension 2, i.e. a ridge of P .

2.3. Coxeter groups. A *Coxeter matrix* $M = (M_{st})_{s,t \in S}$ on a finite set S is a symmetric matrix with the entries $M_{st} \in \{1, 2, \dots, m, \dots, \infty\}$ such that the diagonal entries $M_{ss} = 1$ and the others $M_{st} \neq 1$. To any Coxeter matrix $M = (M_{st})_{s,t \in S}$ we associate a *Coxeter group* $W_{S,M}$ given by a presentation $\langle S \mid (st)^{M_{st}} = 1 \text{ for } M_{st} \neq \infty \rangle$. We denote the Coxeter group $W_{S,M}$ also simply by W , W_S , or W_M . The *rank* of W_S is the cardinality $\sharp S$ of S .

The *Coxeter diagram* of $W_{S,M}$ is a labelled graph \mathcal{G}_W such that: (i) the set of nodes (i.e. vertices) of \mathcal{G}_W is the set S , (ii) two nodes $s, t \in S$ are connected by an edge \overline{st} of \mathcal{G}_W if and only if $M_{st} \in \{3, \dots, m, \dots, \infty\}$, and (iii) the edge \overline{st} is labelled by M_{st} if and only if $M_{st} > 3$. A Coxeter group W is *irreducible* if the Coxeter diagram \mathcal{G}_W is connected.

¹⁴The *combinatorics* (or *face poset*) of a polytope is the poset of its faces partially ordered by inclusion.

¹⁵Two posets \mathcal{P}_1 and \mathcal{P}_2 are *dual* to each other provided there exists an order-reversing isomorphism between \mathcal{P}_1 and \mathcal{P}_2 .

An irreducible Coxeter group W is *spherical* (resp. *affine*) if it is finite (resp. infinite and virtually abelian). For a Coxeter group W (not necessarily irreducible), each connected component of the Coxeter diagram \mathcal{G}_W corresponds to a Coxeter group, called a *component* of W . A Coxeter group W is *spherical* (resp. *affine*) if each component of W is spherical (resp. affine). We sometimes refer to Appendix A for the list of all the irreducible spherical and irreducible affine Coxeter diagrams.

For each $T \subset S$, the subgroup W' of W generated by T is called a *standard subgroup* of W . It is well known that W' identifies with the Coxeter group W_{T, M_T} , where M_T is the restriction of M to $T \times T$. A subset $T \subset S$ is said to be “*something*” if the Coxeter group W_T is “something”. For example, the word “something” can be replaced by “spherical”, “affine”, and so on. Two subsets $T, U \subset S$ are *orthogonal* if $M_{tu} = 2$ for every $t \in T$ and every $u \in U$. This relationship is denoted by $T \perp U$.

2.4. Coxeter polytopes. Recall that a *Coxeter polytope* is a mirror polytope whose dihedral angles are submultiples of π , i.e. each dihedral angle is π/m for some integer $m \geq 2$ or $m = \infty$.

Let P be a Coxeter polytope with the set of facets S and the set of reflections $\{\sigma_s = \text{Id} - \alpha_s \otimes b_s\}_{s \in S}$. The *Coxeter group* W_P of P is the Coxeter group $W_{S, M}$ associated with a Coxeter matrix $M = (M_{st})_{s, t \in S}$ satisfying that $M_{st} = m_{st}$ if $\alpha_s(b_t)\alpha_t(b_s) = 4 \cos^2(\pi/m_{st})$ and $M_{st} = \infty$ if $\alpha_s(b_t)\alpha_t(b_s) \geq 4$. For a proper face f of P (i.e. $f \neq \emptyset, P$), we write $\sigma(f) = \{s \in S \mid f \subset s\}$ and $W_f := W_{\sigma(f)}$.

Theorem 2.11 (Tits [14, Chapter V] for Tits simplex, and Vinberg [65]). *Let P be a Coxeter polytope of \mathbb{S}^n with Coxeter group W_P and let Γ_P be the subgroup of $\text{SL}_{n+1}^{\pm} \mathbb{R}$ generated by the reflections $\{\sigma_s\}_{s \in S}$. Then:*

- (1) *The homomorphism $\sigma : W_P \rightarrow \Gamma_P \subset \text{SL}_{n+1}^{\pm} \mathbb{R}$ defined by $\sigma(s) = \sigma_s$ is an isomorphism.*
- (2) *The Γ_P -orbit of P is a convex subset \mathcal{C}_P of \mathbb{S}^n , and $\gamma \text{Int}(P) \cap \text{Int}(P) = \emptyset$ for all non-trivial $\gamma \in \Gamma_P$.*
- (3) *The group Γ_P acts properly discontinuously on the interior Ω_P of \mathcal{C}_P .*
- (4) *An open proper face f of P lies in Ω_P if and only if the Coxeter group W_f is spherical.*

Theorem 2.11 tells us that Ω_P is a convex domain of \mathbb{S}^n , and that if Ω_P is properly convex then Ω_P/Γ_P is a convex projective Coxeter orbifold.

2.5. Relative hyperbolicity. Let Y be a proper Gromov-hyperbolic space (see e.g. [39, Section 2] for a quick review and [16, Part III.3] for

details on Gromov-hyperbolic spaces). We recall that for every isometry γ of Y exactly one of the following holds:

- (1) γ fixes a point of Y .
- (2) γ fixes exactly one point of the Gromov boundary ∂Y of Y .
- (3) γ fixes two points of ∂Y .

We say that γ is *parabolic* (resp. *hyperbolic*) if (2) (resp. (3)) holds. Let Γ be a subgroup of isometries of Y that acts properly discontinuously. A subgroup of Γ is *parabolic* if it is infinite and contains no hyperbolic element. A parabolic subgroup fixes a unique point of ∂Y , called a *parabolic point*. The stabiliser of a parabolic point is a maximal parabolic subgroup.

Relative hyperbolicity has many equivalent definitions, see e.g. [39, Section 3]. We recall one of them, named *cusp uniform action*. A group Γ is *relatively hyperbolic with respect to a collection \mathcal{P} of subgroups* if there exist a proper Gromov-hyperbolic metric space Y and a properly discontinuous effective action of Γ on Y by isometry such that:

- the collection \mathcal{P} is a set of representatives of the conjugacy classes of maximal parabolic subgroups of Γ ,
- there exists a Γ -equivariant collection \mathcal{H} of disjoint open horoballs¹⁶ centered at the parabolic points of Γ ,
- the action of Γ on $Y \setminus U$ is cocompact, where U denotes the union of the horoballs in \mathcal{H} .

For example, the fundamental group of a cusped hyperbolic n -manifold (or n -orbifold) is relatively hyperbolic with respect to its cusp subgroups, which are virtually \mathbb{Z}^{n-1} [15, 31].

For $k \geq 2$, any \mathbb{Z}^k -subgroup Λ of a relatively hyperbolic group Γ with respect to a collection \mathcal{P} of subgroups must lie in a conjugate of a subgroup $P \in \mathcal{P}$. Indeed, the centraliser of a hyperbolic element in a discrete subgroup of isometries of Y is virtually \mathbb{Z} . Thus Λ must contain a parabolic isometry δ with a unique fixed point $p \in \partial Y$, and any other element $\gamma \in \Lambda$ also has to fix p since it commutes with δ . So Λ lies in the stabiliser of p .

In particular, the fundamental group $\pi_1 X$ of Theorem A, which is relatively hyperbolic with respect to the collection $\mathcal{P} = \{\pi_1 T_i, \pi_1 T'_i\}_i$ of rank-2 abelian subgroups, is not relatively hyperbolic with respect to any proper subcollection of \mathcal{P} .

We end this section by giving a criterion to determine when a Coxeter group is relatively hyperbolic with respect to a collection of standard subgroups. We will use this criterion in Subsection 3.2.

¹⁶We refer to [39, Section 2] for the notion of horoballs in Gromov-hyperbolic space.

Theorem 2.12 (Moussong [55] and Caprace [18, 19]). *Let W_S be a Coxeter group, and let \mathcal{T} be a collection of subsets of S . Then the group W_S is relatively hyperbolic with respect to $\{W_T \mid T \in \mathcal{T}\}$ if and only if the following hold:*

- (1) *For every irreducible affine subset $U \subset S$ of rank ≥ 3 , there exists $T \in \mathcal{T}$ such that $U \subset T$.*
- (2) *For every pair of irreducible non-spherical subsets S_1, S_2 of S with $S_1 \perp S_2$, there exists $T \in \mathcal{T}$ such that $S_1 \cup S_2 \subset T$.*
- (3) *For every pair $T, T' \in \mathcal{T}$ with $T \neq T'$, the intersection $T \cap T'$ is spherical.*
- (4) *For every $T \in \mathcal{T}$ and every irreducible non-spherical subset $U \subset T$, we have $U^\perp \subset T$, where $U^\perp := \{s \in S \mid s \perp U\}$.*

2.6. Operation on a simplex. Here we introduce three uniform¹⁷ Euclidean 4-polytopes via truncation, rectification, and bitruncation of the 4-simplex.

Roughly speaking, by truncation, rectification, and bitruncation of a regular polytope $P \subset \mathbb{R}^n$ we mean cutting P uniformly at every vertex with a hyperplane orthogonal to the line joining the vertex to the barycentre. This operation is nicely described in Coxeter's classic book [26, Section 8.1]. For example, a truncated (resp. rectified, resp. bitruncated) 3-simplex is a truncated tetrahedron (resp. an octahedron, resp. a truncated tetrahedron) in Figure 2. Combinatorially, by collapsing some ridges of the bitruncated P to vertices, one gets the rectified P .




FIGURE 2. The truncated (left), rectified (middle), and bitruncated (right) 3-simplex.

¹⁷A polytope P of dimension $n \geq 3$ (in the Euclidean space) is *uniform* if it is a vertex-transitive polytope with uniform facets. A *uniform polygon* is a regular polygon. By *vertex-transitive* we mean that the symmetry group of P acts transitively on the set of vertices of P .

We now explain this operation in detail for the 4-simplex. Consider a regular 4-simplex $\Delta \subset \mathbb{R}^4$ with barycentre at the origin and vertex v_1, \dots, v_5 . We denote by F_i the facet of Δ opposite to v_i and by H_i the closed half-space containing the origin with $F_i \subset \partial H_i$. Then $\Delta = H_1 \cap \dots \cap H_5$.

Let $c := |v_i|$ and fix a positive parameter $s \leq c$. We denote by H'_i the closed half-space (depending on s) containing the origin such that $\partial H'_i$ is orthogonal to v_i and $\frac{s}{c}v_i \in \partial H'_i$, and we set

$$Q_s = H_1 \cap \dots \cap H_5 \cap H'_1 \cap \dots \cap H'_5.$$

Note that $Q_c = \Delta$ is the original simplex. There exist some numbers $0 < a < b < c$ such that the combinatorics of the 4-polytope Q_s is constant for s in (a, b) and (b, c) , and changes at $s = a, b$, and c . The polytope Q_s (depicted in Figure 3) is called:

- a *truncated 4-simplex* for $s \in (b, c)$,
- a *rectified 4-simplex* for $s = b$,
- a *bitruncated 4-simplex* for $s \in (a, b)$.




FIGURE 3. The Schlegel diagrams of the truncated (left), rectified (middle), and bitruncated (right) 4-simplex. The facets F'_i are coloured darker than F_i (cf. Table 1 below).

The rectified simplex Q_b is in fact the convex hull of the midpoints of the edges of the regular simplex $\Delta = Q_c$, while Q_a is another rectified simplex. For all $s \in [a, c]$, the polytope Q_s is uniform.

2.7. The combinatorics of the rectified and bitruncated 4-simplices. We describe the combinatorics of Q_s for $s \in (a, b]$.

The link of a vertex of the rectified (resp. bitruncated) 4-simplex is a triangular prism (resp. a tetrahedron) as in Figure 4. Each vertex of Q_s is the intersection of facets $F'_{i_1} \cap F'_{i_2} \cap F'_{i_3} \cap F'_{i_4} \cap F_{i_5}$ for all distinct i_1, \dots, i_5 (resp. $F'_{i_1} \cap F'_{i_2} \cap F'_{i_3} \cap F_{i_4}$ for all distinct i_1, \dots, i_4), where F_i and F'_i denote the facets of Q_s whose supporting hyperplanes are ∂H_i and $\partial H'_i$, respectively. The ten facets of Q_s are divided into

five octahedra (resp. truncated tetrahedra) $F_i \subset \partial H_i$, and five tetrahedra (resp. truncated tetrahedra) $F'_i \subset \partial H'_i$. For all $i \neq j$, the ridge $F_i \cap F_j$ is a triangle, $F'_i \cap F'_j$ is a vertex (resp. triangle), and $F_i \cap F'_j$ is a triangle (resp. hexagon), while $F_i \cap F'_i = \emptyset$ (see Table 1).

	Truncated 4-simplex	Rectified 4-simplex	Bitruncated 4-simplex
F_i	Truncated tetrahedron	Octahedron	Truncated tetrahedron
F'_j	Tetrahedron	Tetrahedron	Truncated tetrahedron
$F_i \cap F_j$	Hexagon	Triangle	Triangle
$F_i \cap F'_i$	\emptyset	\emptyset	\emptyset
$F_i \cap F'_j$	Triangle	Triangle	Hexagon
$F'_i \cap F'_j$	\emptyset	Vertex	Triangle

TABLE 1. Some information on the faces of the truncated, rectified and bitruncated 4-simplex. The symbols i, j are two distinct indices in $\{1, \dots, 5\}$.




FIGURE 4. The vertex link of the rectified (left) and bitruncated (right) 4-simplex Q_s . A facet of the link labelled by i (resp. i') corresponds to the facet F_i (resp. F'_i) of the 4-polytope Q_s .

The following is a simple observation, but it will be useful later to prove Theorem A (more precisely, Proposition 3.1).

Lemma 2.13. *Let Q_s be the rectified or bitruncated 4-simplex for $s \in (a, b]$. Relabel each facet F'_i of Q_s with $F_{i'}$, and let $S := \{1', \dots, 5', 1, \dots, 5\}$.*

- (1) *In the case where Q_s is the rectified 4-simplex, i.e. $s = b$, each vertex of Q_s corresponds to a subset $\{i', j', k, l, m\} \subset S$ with $\#\{i, j, k, l, m\} = 5$ and each edge of Q_s corresponds to a subset $\{i', j, k\} \subset S$ with $\#\{i, j, k\} = 3$.*
- (2) *In the case where Q_s is the bitruncated 4-simplex, i.e. $a < s < b$, each vertex of Q_s corresponds to a subset $\{i', j', k, l\} \subset S$ with $\#\{i, j, k, l\} = 4$.*

2.8. The ideal hyperbolic rectified 4-simplex. Every vertex-transitive polytope $Q \subset \mathbb{R}^n$ can be realised as an ideal hyperbolic n -polytope, obtained by interpreting the ball in which Q is inscribed as a projective model of the hyperbolic n -space \mathbb{H}^n . What is nice about the rectified

simplex of dimension $n \leq 4$ is that its regular ideal hyperbolic realisation $R \subset \mathbb{H}^n$ is a Coxeter polytope.¹⁸ For instance, the polytope R of dimension 3 is a right-angled hyperbolic octahedron.

Let $R \subset \mathbb{H}^4$ be the ideal rectified 4-simplex. The facets of R are regular ideal Coxeter 3-polytopes: five right-angled octahedra F_i and five $\pi/3$ -angled tetrahedra F'_i . The horospherical link of any (ideal) vertex of R is a Euclidean right triangular prism with equilateral bases (see the left of Figure 4). Thus, for all $i \neq j$ the dihedral angle at a ridge $F_i \cap F_j$ (resp. $F'_i \cap F'_j$) is $\pi/3$ (resp. $\pi/2$), while F'_i and F'_j are parallel, i.e. the facet F'_i is tangent to F'_j at infinity.

Remark 2.14. The bitruncated 4-simplex is “combinatorially” a “filling” of the ideal rectified 4-simplex in the sense that the latter is obtained from the former by collapsing each triangular ridge $F'_i \cap F'_j$ for $i \neq j$ to a point and removing it. We call such triangles the *filling ridges* of the bitruncated 4-simplex.

2.9. Decomposability and Euler characteristics. We conclude the section with a remark on convex projective manifolds that has probably been noticed by many experts on the subject, but whose explicit statement seems to be missing in the literature. This remark is a simple consequence of works of Vey, Benoist, and Gottlieb.

Fact 2.15. *If a convex projective manifold Ω/Γ is decomposable, then $\chi(\Omega/\Gamma) = 0$.*

Proof: Assume by contradiction that Ω/Γ is decomposable but $\chi(\Omega/\Gamma) \neq 0$. By a theorem of Gottlieb [36, Corollary IV.3], the fundamental group of a finite, aspherical polyhedron with non-zero Euler characteristic has trivial centre. But since Ω/Γ is decomposable, by Proposition 4.4 of Benoist [6] (see also [64]), the centre of Γ contains a non-trivial free abelian subgroup. □

In particular, since $\chi(X) \neq 0$, the convex projective manifold X of Theorem A is indecomposable.

3. The proof of Theorem A

In this section, we prove Theorem A. In Subsection 3.1, we perform convex projective generalised Dehn filling on the ideal hyperbolic rectified 4-simplex $R \subset \mathbb{H}^4$, and build mirror polytopes P_α combinatorially

¹⁸It is well known that the link of the regular ideal hyperbolic rectified n -simplex is a Euclidean right simplicial $(n - 1)$ -prism with regular $(n - 2)$ -simplicial bases, and the dihedral angle of the Euclidean regular $(n - 2)$ -simplex is $\arccos(1/(n - 2))$. The latter is π/m for some integer $m \geq 2$ if and only if $n \leq 4$. Hence, R is a Coxeter polytope if and only if $n \leq 4$.

equivalent¹⁹ to the bitruncated 4-simplex. In Subsection 3.2, we show that the Coxeter group $W_p = W_{P_{\pi/p}}$ of the Coxeter polytope $P_{\pi/p}$ is relatively hyperbolic. In Subsections 3.3 and 3.4, we construct the cusped hyperbolic 4-manifold M , by gluing some copies of R , and a filling X of M , respectively. Finally, in Subsection 3.5, we give X a projective cone-manifold structure induced by P_α , and finish the proof of Theorem A.

3.1. Deforming the rectified 4-simplex. In this subsection, we obtain a family of projective structures on the orbifold \mathcal{O}_R associated with the ideal hyperbolic rectified 4-simplex R by deforming its original complete hyperbolic structure.

Accordingly, the composition of two projective reflections along the facets F'_i and F'_j of \mathcal{O}_R , $i \neq j$ (recall the notation from Subsection 2.6), will deform to be conjugate to a projective rotation of angle $2\alpha > 0$. At the original hyperbolic structure on \mathcal{O}_R , the facets F'_i and F'_j are parallel, i.e. $\alpha = 0$. But, at the deformed projective structure on \mathcal{O}_R , new ridges $F'_i \cap F'_j$ can be added to \mathcal{O}_R to form a mirror polytope P_α that is combinatorially equivalent to a bitruncated 4-simplex, where α is the dihedral angle of the ridge $F'_i \cap F'_j$. So, the goal is to prove the following:

Proposition 3.1. *There exists a path $(0, \pi/3] \ni \alpha \mapsto P_\alpha$ of mirror polytopes with the combinatorics of a bitruncated 4-simplex and dihedral angles:*

- α at the filling ridges $F'_i \cap F'_j$,
- $\pi/2$ at the ridges $F_i \cap F'_j$, and
- $\pi/3$ at the ridges $F_i \cap F_j$.

Moreover, the limit P_0 is the ideal hyperbolic rectified 4-simplex R .

Before proving Proposition 3.1, we begin with some auxiliary lemmas. First, let

$$t_3 = \frac{1}{2} \left(11 + 9\sqrt{2} - 3\sqrt{31 + 22\sqrt{2}} \right) \approx 0.0422$$

and for $t \in [t_3, 1]$,

$$f(t) = \frac{t(t+2)^3(2t+1)^3}{(t^2+t+1)^2(t^2+7t+1)^2}, \quad h(t) = 2 + \frac{1}{t} + \frac{t(t+2)^4}{(t^2+t+1)(t^2+7t+1)},$$

$$g_p(t) = \frac{2t^p(t+2)^p(2t+1)^{3-p}}{(t^2+t+1)(t^2+7t+1)}, \quad \text{and} \quad \bar{g}_p(t) = \frac{4f(t)}{g_p(t)},$$

¹⁹Two polytopes Q and Q' are *combinatorially equivalent* if the face poset of Q is isomorphic to the face poset of Q' .

where $p = 0, 1, 2, 3$. The number t_3 is the unique positive solution less than 1 satisfying the equation $f(t_3) = 1/4 = \cos^2(\pi/3)$. Since t_3 is positive, it is easy to check:

Lemma 3.2. *The functions f, h, g_p , and $\bar{g}_p : [t_3, 1] \rightarrow \mathbb{R}$ are well defined and positive.*

Given $t \in [t_3, 1]$, we now consider the matrix

$$C_t = \left(\begin{array}{ccccc|ccccc} 2 & -\bar{g}_0(t) & -\bar{g}_1(t) & -\bar{g}_2(t) & -\bar{g}_3(t) & -h(t) & 0 & 0 & 0 & 0 \\ -g_0(t) & 2 & -\bar{g}_0(t) & -\bar{g}_1(t) & -\bar{g}_2(t) & 0 & -h(t) & 0 & 0 & 0 \\ -g_1(t) & -g_0(t) & 2 & -\bar{g}_0(t) & -\bar{g}_1(t) & 0 & 0 & -h(t) & 0 & 0 \\ -g_2(t) & -g_1(t) & -g_0(t) & 2 & -\bar{g}_0(t) & 0 & 0 & 0 & -h(t) & 0 \\ -g_3(t) & -g_2(t) & -g_1(t) & -g_0(t) & 2 & 0 & 0 & 0 & 0 & -h(t) \\ \hline -2 & 0 & 0 & 0 & 0 & 2 & -1/t & -1/t & -1/t & -1/t \\ 0 & -2 & 0 & 0 & 0 & -t & 2 & -1/t & -1/t & -1/t \\ 0 & 0 & -2 & 0 & 0 & -t & -t & 2 & -1/t & -1/t \\ 0 & 0 & 0 & -2 & 0 & -t & -t & -t & 2 & -1/t \\ 0 & 0 & 0 & 0 & -2 & -t & -t & -t & -t & 2 \end{array} \right).$$

Lemma 3.3. *For every $t \in [t_3, 1]$, the matrix C_t is an irreducible Cartan matrix of negative type and of rank 5.*

Proof: It easily follows from Lemma 3.2 that C_t is an irreducible Cartan matrix. A simple but long computation shows that the rank of C_t is 5, which is less than $9 = 10 - 1$. Hence, the irreducible Cartan matrix C_t is of negative type by Proposition 2.5. \square

Lemma 3.4. *There exists a monotonically decreasing analytic function $\alpha : [t_3, 1] \rightarrow \mathbb{R}$ satisfying*

$$\cos^2 \alpha(t) = f(t), \quad \alpha(t_3) = \frac{\pi}{3}, \quad \alpha(1) = 0.$$




FIGURE 5. The graph of $f(t)$ over the interval $[0, 1]$.

Proof: The derivative f' of f satisfies:

$$f'(t) = -\frac{2(t-1)(t+1)(t+2)^2(2t+1)^2(t^4+2t^3+21t^2+2t+1)}{(t^2+t+1)^3(t^2+7t+1)^3}.$$

Since $f'(t)$ is positive for all $t \in [t_3, 1)$ and is zero for $t = 1$, the function $f: [t_3, 1] \rightarrow \mathbb{R}$ is monotonically increasing with $f(t_3) = 1/4$ and $f(1) = 1$ (see Figure 5). The lemma now follows easily. \square

We are finally ready to prove Proposition 3.1 by applying Theorems 2.6 and 2.8. Recall the combinatorics of the truncated, rectified, and bitruncated 4-simplex, described in Table 1. See Figure 6 for a geometric picture of the vertex link of the mirror polytope P_α .




FIGURE 6. The vertex links of a mirror polytope are also mirror polytopes. For P_α we have: (i) a right triangular prism with equilateral bases for the (ideal) rectified 4-simplex P_0 (left), and (ii) a tetrahedron for the bitruncated 4-simplex P_α , $\alpha \in (0, \frac{\pi}{3}]$ (right). The edges of dihedral angle $\pi/2$ (resp. $\pi/3$, resp. α) are drawn in black (resp. red, resp. green).

Proof of Proposition 3.1: Let us call and order the indices of C_t as $S := \{1', \dots, 5', 1, \dots, 5\}$. Lemmas 3.3 and 3.4 together with Theorem 2.6 imply that there exists a path $[0, \pi/3] \ni \alpha \mapsto P_\alpha$ of mirror polytopes in \mathbb{S}^4 with facets $\{F_s\}_{s \in S}$ and with Cartan matrix C_t . The facet $F_{i'}$ of P_α ($i = 1, \dots, 5$) is also denoted by F'_i .

We consider two cases separately: (i) $\alpha \in (0, \pi/3]$ and (ii) $\alpha = 0$.

In case (i), equivalently, $t \in [t_3, 1)$, by Remark 2.10 (or Theorem 2.8(1)), the intersections $F'_i \cap F'_j$, $F_i \cap F'_j$, and $F_i \cap F_j$, $i \neq j$, are ridges of P_α and their dihedral angles are α , $\pi/2$, and $\pi/3$, respectively, because for example

$$(C_t)_{i'j'}(C_t)_{j'i'} = g_p(t)\bar{g}_p(t) = 4f(t) = 4\cos^2 \alpha < 4.$$

We now claim that P_α is combinatorially equivalent to the bitruncated 4-simplex. For every subset $T = \{i', j', k, l\} \subset S$ with $\#\{i, j, k, l\} = 4$,

the submatrix $(C_t)_T$ is the direct sum $(C_t)_{\{i',j'\}} \oplus (C_t)_{\{k,l\}}$ of matrices of positive type, hence $\{F_s\}_{s \in T} \in \sigma(P_\alpha)$ ²⁰ and $\cap_{s \in T} F_s$ is a vertex of P_α by Theorem 2.8(1).

That is, if \mathcal{V} denotes the set of all subsets $\{F'_i, F'_j, F_k, F_l\}$ with $\#\{i, j, k, l\} = 4$, then $\mathcal{V} \subset \sigma(P_\alpha)$. Let $\hat{S} = \{F_s\}_{s \in S}$, and let \mathcal{F} be the subset of $2^{\hat{S}}$ defined by:

$$\mathcal{F} := \{\hat{T}' \in 2^{\hat{S}} \mid \hat{T}' \subset \hat{T} \text{ for some } \hat{T} \in \mathcal{V}\}.$$

As in the previous argument, Theorem 2.8(1) implies that \mathcal{F} is a subset of $\sigma(P_\alpha)$. We know, in addition, from Lemma 2.13(2) and Figure 4 that the poset \mathcal{F} is dual to the face poset of the bitruncated 4-simplex. It is a well-known fact (e.g. [17, Exercise 1.1.20]) that if two polytopes Q and Q' are of the same dimension and the face poset of Q' is a subset of the face poset of Q , then Q is combinatorially equivalent to Q' . As a result, the polytope P_α is combinatorially equivalent to the bitruncated 4-simplex, as claimed.

In case (ii), equivalently, $t = 1$, we claim that P_0 is combinatorially equivalent to the rectified 4-simplex. For every subset $U = \{i', j', k, l, m\} \subset S$ with $\#\{i, j, k, l, m\} = 5$, the submatrix $(C_1)_U$ is the direct sum $(C_1)_{\{i',j'\}} \oplus (C_1)_{\{k,l,m\}}$ of matrices of zero type and the rank of $(C_1)_U$ is 3. Hence $\{F_s\}_{s \in U} \in \sigma(P_0)$ and $\cap_{s \in U} F_s$ is a vertex of P_0 by Theorem 2.8(2). Furthermore, for every subset $U' = \{i', j, k\} \subset S$ with $\#\{i, j, k\} = 3$, the submatrix $(C_1)_{U'}$ is of positive type. Hence $\{F_s\}_{s \in U'} \in \sigma(P_0)$ and $\cap_{s \in U'} F_s$ is an edge of P_0 by Theorem 2.8(1). Then, as in the proof of case (i), using Lemma 2.13(1) and Figure 4, we may conclude that the polytope P_0 is combinatorially equivalent to the rectified 4-simplex, as claimed. Finally, a simple computation shows that the Cartan matrix C_1 is equivalent to a symmetric matrix of signature $(4, 1)$, and therefore the polytope P_0 (without vertices) may be identified with the ideal hyperbolic rectified 4-simplex R . □

Remark 3.5. The symmetry group of the mirror polytope P_α , $\alpha \in [0, \pi/3]$, is of order ≥ 5 . For, if we set $\hat{Q} = \begin{pmatrix} Q & 0 \\ 0 & Q \end{pmatrix}$ with

$$Q = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix},$$

²⁰See the end of Subsection 2.2 for the definition of the poset $\sigma(P)$.

then \hat{Q} is a permutation matrix of order 5 and $\hat{Q}C_t\hat{Q}^{-1}$ is equivalent to C_t .

Remark 3.6. A (slightly more complicated) computation reveals that the deformation space of projective structures on the orbifold \mathcal{O}_R associated with the ideal hyperbolic rectified 4-simplex R is 6-dimensional. In other words, one can find a 6-parameter family of Cartan matrices (of rank 5) which correspond to projective structures on \mathcal{O}_R . But, we choose a particular 1-parameter family of Cartan matrices having symmetry as described in Remark 3.5 in order to simplify the computation.

We end the subsection with the following.

Corollary 3.7. *For every integer $p \geq 3$, the mirror polytope $P_{\pi/p}$ is a Coxeter polytope. Moreover, if Γ_p denotes the subgroup of $SL_5^{\pm} \mathbb{R}$ generated by the associated reflections, then the Γ_p -orbit of $P_{\pi/p}$ is a properly convex domain Ω_p of \mathbb{S}^4 , i.e. it is divisible by Γ_p .*

Proof: It is obvious that the mirror polytope $P_{\pi/p}$ is a Coxeter polytope. For any subset $T = \{i', j', k, l\} \subset S$ with $\#\{i, j, k, l\} = 4$, the standard subgroup W_T of the Coxeter group W_S of $P_{\pi/p}$ is isomorphic to $D_p \times D_3$, where D_m is the dihedral group of order $2m$, hence it is a finite group. In the proof of Proposition 3.1, we show that every vertex of $P_{\pi/p}$ corresponds to a subset $\{i', j', k, l\} \subset S$ with $\#\{i, j, k, l\} = 4$, so by Theorem 2.11, the Γ_p -orbit of $P_{\pi/p}$ is a divisible convex domain of \mathbb{S}^4 . \square

3.2. The Coxeter group W_p . The goal of this subsection is to show that the Coxeter group W_p of $P_{\pi/p}$ is relatively hyperbolic with respect to a collection of virtually abelian subgroups of rank 2. To do so, we need to analyse the Coxeter diagram of W_p in Figure 7, and to use Theorem 2.12 together with the (complete) list of the irreducible spherical and affine Coxeter groups in Appendix A.

The following collection of subsets of $S = \{1', \dots, 5', 1, \dots, 5\}$ is denoted by \mathcal{T}_p :

- In the case $p \geq 4$, the collection \mathcal{T}_p consists of all subsets $\{i', j', k, l, m\}$ of S with $\#\{i, j, k, l, m\} = 5$, so the cardinality of \mathcal{T}_p , $p \geq 4$, is 10.
- In the case $p = 3$, the collection \mathcal{T}_3 consists of all subsets $\{i', j', k, l, m\}$ and $\{i', j', k', l, m\}$ of S with $\#\{i, j, k, l, m\} = 5$, so the cardinality of \mathcal{T}_3 is 20.

For each $U \in \mathcal{T}_p$, the standard subgroup $(W_p)_U$ of W_p is isomorphic to $\tilde{A}_2 \times I_2(p)$ (see Appendix A), hence it is virtually isomorphic to \mathbb{Z}^2 .




FIGURE 7. The Coxeter diagram of the Coxeter group W_p of $P_{\pi/p}$.

Proposition 3.8. *The Coxeter group W_p is relatively hyperbolic with respect to the collection of subgroups $\{(W_p)_U \mid U \in \mathcal{T}_p\}$, in particular, a collection of virtually abelian subgroups of rank 2.*

Proof: We only prove it for the case $p = 3$; the argument is similar for other cases where $p \geq 4$. Thanks to Theorem 2.12, we just need to carefully analyse the Coxeter diagram of W_p in Figure 7, using the list of irreducible spherical and affine Coxeter groups in Appendix A.

First, condition (1) holds because all irreducible affine subsets $U \subset S$ of rank ≥ 3 are $\{i', j', k'\}$ and $\{i, j, k\}$ with $\#\{i, j, k\} = 3$. Second, condition (2) holds because there does not exist a pair of irreducible non-spherical subsets S_1, S_2 of S with $S_1 \perp S_2$. Third, condition (3) holds because for every pair $T, T' \in \mathcal{T}_3$ with $T \neq T'$ the intersection $T \cap T'$ is a subset of $\{i', j', k, l\}$ with $\#\{i, j, k, l\} = 4$. Finally, condition (4) holds because for every $T \in \mathcal{T}_3$ there exists only one irreducible non-spherical subset $U \subset T$, which is either $\{i', j', k'\}$ or $\{i, j, k\}$ with $\#\{i, j, k\} = 3$, and $T = U \sqcup U^\perp$. \square

Remark 3.9. The Coxeter group W_p of $P_{\pi/p}$ is in fact a finite-index subgroup of the following Coxeter group \hat{W}_p :



It is easier to verify conditions (1)–(4) of Theorem 2.12 for the Coxeter group \hat{W}_p .

3.3. The cusped hyperbolic manifold. In this subsection, we build the cusped hyperbolic manifold M of Theorem A. Readers who are not familiar with hyperbolic manifolds (with totally geodesic boundary) can consult [50, Chapter 3].




FIGURE 8. The edge-labelled complete graph K_6 .

We first recall the construction of a building block B by Kolpakov and Slavich [42]. Consider the complete graph K_6 on six vertices with its edges labelled by numbers in $\{1, \dots, 5\}$ so that adjacent edges have distinct labels (see Figure 8). For each vertex of K_6 , take a copy of the ideal hyperbolic rectified 4-simplex $R \subset \mathbb{H}^4$, described in Subsection 2.8. If two vertices of K_6 are connected by an edge of label $i \in \{1, \dots, 5\}$, then we glue together the facets F_i of the two corresponding copies of R using the identity as a gluing map.

Proposition 3.10 ([42, Section 3]). *Let B be the building block constructed above.*

- *The space B is a non-orientable, complete, finite-volume hyperbolic 4-manifold with non-compact totally geodesic boundary.*
- *The boundary ∂B of B has exactly five connected components $\partial_1, \dots, \partial_5$, each tessellated by the facets F_i' of the copies of R in B .*
- *The hyperbolic manifold B has exactly ten cusps C_{ij} , $i < j \in \{1, \dots, 5\}$. Each cusp section S_{ij} is diffeomorphic to $K \times [0, 1]$, where K denotes the Klein bottle. One boundary component of S_{ij} is contained in ∂_i , and the other in ∂_j .*

Now, consider again the edge-labelled graph K_6 in Figure 8. For each vertex of K_6 , take a copy of B . If two vertices of K_6 are joined by

an edge of label $i \in \{1, \dots, 5\}$, then we glue together the boundary components ∂_i of the two corresponding copies of B using the identity as a gluing map. Let us call M' the resulting cusped hyperbolic manifold (without boundary), and M its orientable double cover.

Proposition 3.11. *The orientable hyperbolic 4-manifold M has exactly ten cusps, each with section diffeomorphic to the 3-torus.*

Proof: We obtain the cusps of M' by gluing together the cusps of the copies of B , and the gluing maps are induced by the identity. Hence, each cusp section of M' must be diffeomorphic to $K \times \mathbb{S}^1$. By construction, for each pair $i < j \in \{1, \dots, 5\}$, the simple cycle (of length 6) in the graph K_6 with edges labelled alternately by i and j corresponds to a cusp of M' , thus M' has exactly ten cusps. Since the cusps of M' are non-orientable, each cusp section of M is the orientable double covering of a cusp section of M' . In particular, M has precisely ten cusps, each with 3-torus section. \square

Remark 3.12. The natural map $M \rightarrow \mathcal{O}_R$ is an orbifold covering of degree $6 \cdot 6 \cdot 2 = 72$, where \mathcal{O}_R denotes the orbifold associated with R .

3.4. Topology of the filling. In this subsection, we build the manifold X of Theorem A topologically. The desired (singular) geometric structures on X will be given in the next subsection.

Let $Q \subset \mathbb{R}^4$ be a bitruncated 4-simplex. We define X in the same way as the manifold M (see Subsection 3.3), but substituting the ideal rectified simplex R with the bitruncated simplex Q .

Proposition 3.13. *The space X is a closed, orientable, smooth 4-manifold, containing ten pairwise disjoint embedded 2-tori whose complement is diffeomorphic to the cusped hyperbolic manifold M .*

Proof: To prove that the polyhedral complex X is a smooth manifold, it suffices to show that the link of each vertex is homeomorphic to the 3-sphere. Recall that the polytope Q is vertex transitive. The vertex link L of Q is the tetrahedron on the right in Figure 4.

When we glue the first six copies of Q to form B , for each vertex of Q , six copies of L are glued cyclically around one edge, to form a polyhedral complex L' homeomorphic to the closed 3-disc D^3 . Thus, B is a 4-manifold with boundary. Note that R is homeomorphic to the complement $Q \setminus \cup_{i < j} F'_i \cap F'_j$ of the filling ridges; in particular, B is non-orientable (recall Proposition 3.10). When we glue six copies of B

to get X' , for each vertex of the complex, six copies of L' are glued cyclically around a circle $C \subset \partial L'$. The resulting polyhedral complex is clearly homeomorphic to the 3-sphere, so X' is a manifold. Its orientation cover X is thus a manifold.

Now, the cusped hyperbolic manifold M is diffeomorphic to the interior of a compact manifold \overline{M} with boundary $\partial\overline{M}$ consisting of ten 3-tori (recall Proposition 3.11). Let $\Sigma \subset X$ be the union of the copies of the filling ridges of Q . Clearly, M is diffeomorphic to $X \setminus \Sigma$. Moreover, $\partial\overline{M}$ is diffeomorphic to the boundary of a regular neighbourhood of Σ in X . Thus, Σ consists of ten 2-tori. \square

We conclude the subsection with some additional information about X .

Remark 3.14. The 4-manifold X has Euler characteristic $\chi(X) = 12$. Indeed, \mathcal{O}_R has orbifold Euler characteristic $\chi^{\text{orb}}(\mathcal{O}_R) = 1/6$ [42, Appendix 1], and the covering $M \rightarrow \mathcal{O}_R$ has degree 72 (recall Remark 3.12), so $\chi(M) = 12$. Since $\chi(\partial\overline{M}) = 0$ and $\chi(\Sigma) = 0$, we have $\chi(X) = \chi(M) = 12$.

Remark 3.15. The manifold X is a filling of M , which has a maximal cusp section S such that each filling curve in S has length 6. The reason is that the maximal, and maximally symmetric, cusp section of \mathcal{O}_R consists of ten Euclidean prisms, each with all edges of length 1 [42, Section 3.2]. Each of the filling curves of X is made of six copies of the height of these prisms. Since $6 < 2\pi$, one cannot apply directly the Gromov–Thurston 2π theorem to conclude, for instance, that X is aspherical. We will see later that, being convex projective, X is in fact aspherical.

Remark 3.16. The 6 theorem, obtained independently by Agol [1] and Lackenby [44], shows that a filling of a cusped hyperbolic 3-manifold carries a hyperbolic structure as soon as the filling curves are of length *strictly* greater than 6. This is thus an improvement of Gromov and Thurston’s 2π theorem in dimension three. It is an open question whether or not it is possible to generalise the 6 theorem to higher dimension as follows: “The fundamental group of a filling of a cusped hyperbolic n -manifold is relatively hyperbolic with respect to the collection of subgroups associated with the inserted $(n - 2)$ -submanifolds, as soon as the filling curves are of length > 6 .”

Note that in dimension $n = 3$ the bound of 6 is sharp, as shown by Agol [1, Section 7]. Remark 3.15 shows that the same bound would be sharp in dimension $n = 4$.

3.5. Cone-manifold structures on the filling. We now conclude the proof of Theorem A. Let $\Sigma \subset X$ be the union of the copies of the filling ridges $F'_i \cap F'_j$. We first show item (5), giving a path of projective cone-manifold structures on X with the desired properties.

In Proposition 3.1, we built a path of mirror polytope structures $(0, \frac{\pi}{3}] \ni \alpha \mapsto P_\alpha$ on the bitruncated simplex Q . Since the manifold X is built by pairing the facets of some copies of Q through the map induced by the identity, for each α we have a well-defined projective structure on the complement in X of the ridges of the copies of Q . Indeed, the projective structures of the copies of $P_\alpha \setminus \cup(\text{ridges})$ match well via the projective reflections associated with the facets of P_α . We want to show that this projective structure extends to a projective structure on $X \setminus \Sigma$, and on the whole X as projective cone-manifold structures with cone angle $\theta = 6 \cdot \alpha$ along Σ .

The link L of a vertex of P_α is a mirror tetrahedron. Its non-right dihedral angles are α and $\frac{\pi}{3}$ along its two opposite edges $\{i'_1, i'_2\}$ and $\{i_3, i_4\}$, respectively (see Figure 6-right). Recall Subsection 2.1 about projective cone-manifolds. Some copies of L are glued together in X to form a 3-sphere, which we now show to be the projective cone 3-manifold $\mathbb{S}^1 * C_\theta$, where $\theta = 6 \cdot \alpha$.

The manifold X was built in three steps. At the first step (when gluing six copies of Q to build the block B), six copies of L are glued cyclically along its edge $\{i_3, i_4\}$. The resulting space L' is the intersection of two half-spaces of \mathbb{S}^3 with “dihedral angle”²¹ α . At the second step (when gluing six copies of B to build X'), six copies of L' are glued cyclically along its edge. Indeed, the cycle in the graph K_6 with edges labelled alternately by i_1 and i_2 has length 6. The resulting space is thus $\mathbb{S}^1 * C_\theta$. The third step is just the orientation double covering $X \rightarrow X'$, so the local structure of X is the same as that of X' .

The union of the copies of the filling ridge $F'_{i_1} \cap F'_{i_2}$ of Q in X is a component T of Σ (a 2-torus). The holonomy of a meridian of T in X is $(\sigma'_{i_1} \sigma'_{i_2})^3$, where σ'_i is the projective reflection (depending on α) associated with the facet F'_i of the mirror polytope P_α .

Since (X, Σ) is locally modelled on $(\mathbb{S}^2 * C_\theta, \mathbb{S}^2)$, we have a projective cone structure σ_θ for each $\theta \in (0, 2\pi]$. Since $\mathbb{S}^2 * C_{2\pi} = \mathbb{S}^4$, the projective structure $\sigma_{2\pi}$ is non-singular, and each component of Σ is totally geodesic in X (see Remark 2.2). The associated path of projective cone-manifold structures $\theta \mapsto \sigma_\theta$ on X is analytic because the path of Cartan matrices $t \mapsto C_t$ is analytic.

²¹Since to each hyperplane we associate a reflection, it makes sense to talk about the dihedral angle, even if L' is not properly convex.

We have shown item (5) of Theorem A. Clearly, item (1) follows from Proposition 3.11. Item (2) is shown in Remark 3.14, and item (3) in Remark 3.15.

Let us now fix an integer $p \geq 3$. By Corollary 3.7, the mirror polytope $P_{\pi/p}$ is a convex projective orbifold. If moreover $p = 3m$ (i.e. $\theta = 2\pi/m$), the natural map $X \rightarrow P_{\pi/p}$ gives X a convex projective orbifold structure Ω_m/Γ_m with cone structure $\sigma_{2\pi/m}$. We have thus shown the first part of item (6).

Remark 3.17. We already know that the convex projective manifold X is indecomposable. Another way to see this is as follows. Since X covers the Coxeter polytope $P_{\pi/3}$, and the Coxeter group W_3 acts strongly irreducibly on \mathbb{R}^5 because the Cartan matrix of $P_{\pi/3}$ is indecomposable and W_3 is not virtually abelian (see e.g. [49, Theorem 2.18]), so does $\Gamma_1 \cong \pi_1(X)$.

Let T_1, \dots, T_{10} be the components of Σ . Being totally geodesic for each T_i , the natural map $\pi_1 T_i \rightarrow \Gamma_m$ induced by the inclusion $T_i \subset X$ is injective. For $m \geq 2$, the group Γ_m is relatively hyperbolic with respect to $\{\pi_1 T_i\}_i$. Indeed, by Proposition 3.8, the Coxeter group W_{3m} , which is the orbifold fundamental group of $P_{\pi/3m}$, is relatively hyperbolic with respect to the collection $\{(W_{3m})_U \mid U \in \mathcal{T}_{3m}\}$ which corresponds to the fundamental groups of the T_1, \dots, T_{10} . The proof of item (6) is complete.

It remains to show item (4). For $m = 1$ (i.e. $\alpha = \pi/3$ and $\theta = 2\pi$) there is another collection of totally geodesic tori T'_1, \dots, T'_{10} tiled by the ridges $F_i \cap F_j$. Being totally geodesic, also T'_i is π_1 -injective. This time, the group $\Gamma_1 \cong \pi_1 X$ is relatively hyperbolic with respect to $\{\pi_1 T_i, \pi_1 T'_i\}_i$. Indeed, by Proposition 3.8, the Coxeter group W_3 , which is the orbifold fundamental group of $P_{\pi/3}$, is relatively hyperbolic with respect to the collection $\{(W_3)_U \mid U \in \mathcal{T}_3\}$ which corresponds to the fundamental groups of the T_1, \dots, T_{10} and T'_1, \dots, T'_{10} . Finally, for every $T \in \{T_1, \dots, T_{10}\}$ and $T' \in \{T'_1, \dots, T'_{10}\}$, the tori T and T' are transverse (sometimes $T \cap T' = \emptyset$) in X , since the ridges $F'_i \cap F'_j$ and $F_k \cap F_\ell$ are so in \mathbb{S}^4 . Also, in the universal cover Ω , the lifts of T and T' are transverse.

The proof of Theorem A is complete.

Appendix A. The spherical and affine Coxeter diagrams

For the reader’s convenience, we reproduce below the list of the irreducible spherical and irreducible affine Coxeter diagrams.




TABLE 2. The irreducible spherical Coxeter diagrams.




TABLE 3. The irreducible affine Coxeter diagrams.

References

- [1] I. AGOL, Bounds on exceptional Dehn filling II, *Geom. Topol.* **14(4)** (2010), 1921–1940. DOI: 10.2140/gt.2010.14.1921.
- [2] M. T. ANDERSON, Dehn filling and Einstein metrics in higher dimensions, *J. Differential Geom.* **73(2)** (2006), 219–261. DOI: 10.4310/jdg/1146169911.
- [3] R. H. BAMLER, Construction of Einstein metrics by generalized Dehn filling, *J. Eur. Math. Soc. (JEMS)* **14(3)** (2012), 887–909. DOI: 10.4171/JEMS/321.
- [4] S. A. BALLAS, J. DANCIGER, AND G.-S. LEE, Convex projective structures on nonhyperbolic three-manifolds, *Geom. Topol.* **22(3)** (2018), 1593–1646. DOI: 10.2140/gt.2018.22.1593.
- [5] T. BARBOT, F. BONSANTE, AND J.-M. SCHLENKER, Collisions of particles in locally AdS spacetimes I. Local description and global examples, *Comm. Math. Phys.* **308(1)** (2011), 147–200. DOI: 10.1007/s00220-011-1318-6.

- [6] Y. BENOIST, Convexes divisibles II, *Duke Math. J.* **120**(1) (2003), 97–120. DOI: 10.1215/S0012-7094-03-12014-1.
- [7] Y. BENOIST, Convexes divisibles I, in: “*Algebraic Groups and Arithmetic*”, Tata Inst. Fund. Res., Mumbai, 2004, pp. 339–374.
- [8] Y. BENOIST, Convexes divisibles IV. Structure du bord en dimension 3, *Invent. Math.* **164**(2) (2006), 249–278. DOI: 10.1007/s00222-005-0478-4.
- [9] Y. BENOIST, Convexes hyperboliques et quasiisométries, *Geom. Dedicata* **122** (2006), 109–134. DOI: 10.1007/s10711-006-9066-z.
- [10] Y. BENOIST, A survey on divisible convex sets, in: “*Geometry, Analysis and Topology of Discrete Groups*”, Adv. Lect. Math. (ALM) **6**, Int. Press, Somerville, MA, 2008, pp. 1–18.
- [11] N. BERGERON AND T. GELANDER, A note on local rigidity, *Geom. Dedicata* **107** (2004), 111–131. DOI: 10.1023/B:GEOM.0000049122.75284.06.
- [12] S. A. BLEILER AND C. D. HODGSON, Spherical space forms and Dehn filling, *Topology* **35**(3) (1996), 809–833. DOI: 10.1016/0040-9383(95)00040-2.
- [13] M. BOILEAU, B. LEEB, AND J. PORTI, Geometrization of 3-dimensional orbifolds, *Ann. of Math. (2)* **162**(1) (2005), 195–290. DOI: 10.4007/annals.2005.162.195.
- [14] N. BOURBAKI, “*Groupes et algèbres de Lie*”, Chapitres 4, 5 et 6, Elements de Mathématique, Masson, Paris, 1981.
- [15] B. H. BOWDITCH, Relatively hyperbolic groups, *Internat. J. Algebra Comput.* **22**(3) (2012), 1250016, 66 pp. DOI: 10.1142/S0218196712500166.
- [16] M. R. BRIDSON AND A. HAEFLIGER, “*Metric Spaces of Non-Positive Curvature*”, Grundlehren der mathematischen Wissenschaften **319**, Springer-Verlag, Berlin, 1999. DOI: 10.1007/978-3-662-12494-9.
- [17] V. M. BUCHSTABER AND T. E. PANOV, “*Toric Topology*”, Mathematical Surveys and Monographs **204**, American Mathematical Society, Providence, RI, 2015. DOI: 10.1090/surv/204.
- [18] P.-E. CAPRACE, Buildings with isolated subspaces and relatively hyperbolic Coxeter groups, *Innov. Incidence Geom.* **10** (2009), 15–31. DOI: 10.2140/iig.2009.10.15.
- [19] P.-E. CAPRACE, Erratum to “Buildings with isolated subspaces and relatively hyperbolic Coxeter groups”, *Innov. Incidence Geom.* **14** (2015), 77–79. DOI: 10.2140/iig.2015.14.77.
- [20] S. CHOI, G.-S. LEE, AND L. MARQUIS, Deformations of convex real projective manifolds and orbifolds, in: “*Handbook of Group Actions*”, Vol. III, Adv. Lect. Math. (ALM) **40**, Int. Press, Somerville, MA, 2018, pp. 263–310.
- [21] S. CHOI, G.-S. LEE, AND L. MARQUIS, Convex projective generalized Dehn filling, *Ann. Sci. Éc. Norm. Supér. (4)* **53**(1) (2020), 217–266. DOI: 10.24033/asens.2421.
- [22] S. CHOI, G.-S. LEE, AND L. MARQUIS, Deformation spaces of Coxeter truncation polytopes, Preprint (2021). [arXiv:2108.11689](https://arxiv.org/abs/2108.11689).
- [23] M. CONDER AND C. MACLACHLAN, Compact hyperbolic 4-manifolds of small volume, *Proc. Amer. Math. Soc.* **133**(8) (2005), 2469–2476. DOI: 10.1090/S0002-9939-05-07634-3.
- [24] D. COOPER, D. D. HODGSON, AND S. P. KERCKHOFF, “*Three-dimensional Orbifolds and Cone-Manifolds*”, With a postface by Sadayoshi Kojima, MSJ Memoirs **5**, Mathematical Society of Japan, Tokyo, 2000. DOI: 10.2969/msjmemoirs/005010000.

- [25] D. COOPER, D. D. LONG, AND M. B. THISTLETHWAITE, Flexing closed hyperbolic manifolds, *Geom. Topol.* **11** (2007), 2413–2440. DOI: 10.2140/gt.2007.11.2413.
- [26] H. S. M. COXETER, “*Regular Polytopes*”, Third edition, Dover Publications, Inc., New York, 1973.
- [27] J. DANCIGER, Geometric Transitions: From Hyperbolic to Ads Geometry, Thesis (Ph.D.)-Stanford University (2011). Available on ProQuest.
- [28] J. DANCIGER, A geometric transition from hyperbolic to anti-de Sitter geometry, *Geom. Topol.* **17(5)** (2013), 3077–3134. DOI: 10.2140/gt.2013.17.3077.
- [29] J. DANCIGER, F. GUÉRITAUD, AND F. KASSEL, Convex cocompact actions in real projective geometry, Preprint (2017). arXiv:1704.08711.
- [30] J. DANCIGER, F. GUÉRITAUD, AND F. KASSEL, Convex cocompactness in pseudo-Riemannian hyperbolic spaces, *Geom. Dedicata* **192** (2018), 87–126. DOI: 10.1007/s10711-017-0294-1.
- [31] B. FARB, Relatively hyperbolic groups, *Geom. Funct. Anal.* **8(5)** (1998), 810–840. DOI: 10.1007/s000390050075.
- [32] K. FUJIWARA AND J. F. MANNING, CAT(0) and CAT(−1) fillings of hyperbolic manifolds, *J. Differential Geom.* **85(2)** (2010), 229–269. DOI: 10.4310/jdg/1287580965.
- [33] K. FUJIWARA AND J. F. MANNING, Simplicial volume and fillings of hyperbolic manifolds, *Algebr. Geom. Topol.* **11(4)** (2011), 2237–2264. DOI: 10.2140/agt.2011.11.2237.
- [34] H. GARLAND AND M. S. RAGHUNATHAN, Fundamental domains for lattices in (R-)rank 1 semisimple Lie groups, *Ann. of Math. (2)* **92(2)** (1970), 279–326. DOI: 10.2307/1970838.
- [35] W. M. GOLDMAN, Geometric structures on manifolds, University of Maryland, Lecture Notes, 2018.
- [36] D. H. GOTTLIEB, A certain subgroup of the fundamental group, *Amer. J. Math.* **87(4)** (1965), 840–856. DOI: 10.2307/2373248.
- [37] D. GROVES AND J. F. MANNING, Dehn filling in relatively hyperbolic groups, *Israel J. Math.* **168** (2008), 317–429. DOI: 10.1007/s11856-008-1070-6.
- [38] O. GUICHARD AND A. WIENHARD, Anosov representations: domains of discontinuity and applications, *Invent. Math.* **190(2)** (2012), 357–438. DOI: 10.1007/s00222-012-0382-7.
- [39] G. C. HRUSKA, Relative hyperbolicity and relative quasiconvexity for countable groups, *Algebr. Geom. Topol.* **10(3)** (2010), 1807–1856. DOI: 10.2140/agt.2010.10.1807.
- [40] M. KAPOVICH, Convex projective structures on Gromov–Thurston manifolds, *Geom. Topol.* **11** (2007), 1777–1830. DOI: 10.2140/gt.2007.11.1777.
- [41] A. KOLPAKOV, A. W. REID, AND S. RIOLO, Many cusped hyperbolic 3-manifolds do not bound geometrically, *Proc. Amer. Math. Soc.* **148(5)** (2020), 2233–2243. DOI: 10.1090/proc/14573.
- [42] A. KOLPAKOV AND L. SLAVICH, Symmetries of hyperbolic 4-manifolds, *Int. Math. Res. Not. IMRN* **2016(9)** (2016), 2677–2716. DOI: 10.1093/imrn/rnv210.
- [43] F. LABOURIE, Anosov flows, surface groups and curves in projective space, *Invent. Math.* **165(1)** (2006), 51–114. DOI: 10.1007/s00222-005-0487-3.
- [44] M. LACKENBY, Word hyperbolic Dehn surgery, *Invent. Math.* **140(2)** (2000), 243–282. DOI: 10.1007/s002220000047.

- [45] G.-S. LEE, L. MARQUIS, S. RIOLO, AND T. YUKITA, Dehn filling all the cusps of a hyperbolic 4-manifold, In preparation.
- [46] C. LONG, Small volume closed hyperbolic 4-manifolds, *Bull. Lond. Math. Soc.* **40(5)** (2008), 913–916. DOI: 10.1112/blms/bdn077.
- [47] L. MARQUIS, Espace des modules de certains polyèdres projectifs miroirs, *Geom. Dedicata* **147** (2010), 47–86. DOI: 10.1007/s10711-009-9438-2.
- [48] L. MARQUIS, Around groups in Hilbert geometry, in: “*Handbook of Hilbert Geometry*”, IRMA Lect. Math. Theor. Phys. **22**, Eur. Math. Soc., Zürich, 2014, pp. 207–261.
- [49] L. MARQUIS, Coxeter group in Hilbert geometry, *Groups Geom. Dyn.* **11(3)** (2017), 819–877. DOI: 10.4171/GGD/416.
- [50] B. MARTELLI, “*An Introduction to Geometric Topology*”, CreateSpace Independent Publishing Platform, 2016.
- [51] B. MARTELLI, Hyperbolic four-manifolds, in: “*Handbook of Group Actions*”, Vol. III, Adv. Lect. Math. (ALM) **40**, Int. Press, Somerville, MA, 2018, pp. 37–58.
- [52] B. MARTELLI AND S. RIOLO, Hyperbolic Dehn filling in dimension four, *Geom. Topol.* **22(3)** (2018), 1647–1716. DOI: 10.2140/gt.2018.22.1647.
- [53] C. T. MCMULLEN, The Gauss–Bonnet theorem for cone manifolds and volumes of moduli spaces, *Amer. J. Math.* **139(1)** (2017), 261–291. DOI: 10.1353/ajm.2017.0005.
- [54] D. B. MCREYNOLDS, A. W. REID, AND M. STOVER, Collisions at infinity in hyperbolic manifolds, *Math. Proc. Cambridge Philos. Soc.* **155(3)** (2013), 459–463. DOI: 10.1017/S0305004113000364.
- [55] G. MOUSSONG, Hyperbolic Coxeter groups, Thesis (Ph.D.)- The Ohio State University (1988).
- [56] D. V. OSIN, Peripheral fillings of relatively hyperbolic groups, *Invent. Math.* **167(2)** (2007), 295–326. DOI: 10.1007/s00222-006-0012-3.
- [57] S. RIOLO AND A. SEPPI, Geometric transition from hyperbolic to Anti-de Sitter structures in dimension four, Preprint (2019). arXiv:1908.05112.
- [58] S. RIOLO AND L. SLAVICH, New hyperbolic 4-manifolds of low volume, *Algebr. Geom. Topol.* **19(5)** (2019), 2653–2676. DOI: 10.2140/agt.2019.19.2653.
- [59] V. SCHROEDER, A cusp closing theorem, *Proc. Amer. Math. Soc.* **106(3)** (1989), 797–802. DOI: 10.2307/2047438.
- [60] L. SLAVICH, The complement of the figure-eight knot geometrically bounds, *Proc. Amer. Math. Soc.* **145(3)** (2017), 1275–1285. DOI: 10.1090/proc/13272.
- [61] W. THURSTON, “*The Geometry and Topology of Three-Manifolds*”, Electronic version 1.1, March 2002. <http://www.msri.org/publications/books/gt3m/>.
- [62] W. P. THURSTON, Shapes of polyhedra and triangulations of the sphere, in: “*The Epstein Birthday Schrift*”, *Geom. Topol. Monogr.* **1**, Geom. Topol. Publ., Coventry, 1998, pp. 511–549. DOI: 10.2140/gtm.1998.1.511.
- [63] C. VERNICOS, Introduction aux géométries de Hilbert, in: “*Actes de Séminaire de théorie spectrale et géométrie*”, Vol. 23, Année 2004–2005, Sémin. Théor. Spectr. Géom. **23**, Univ. Grenoble I, Saint-Martin-d’Hères, 2005, pp. 145–168. DOI: 10.5802/tsg.236.
- [64] J. VEY, Sur les automorphismes affines des ouverts convexes saillants, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* **24** (1970), 641–665.

- [65] È. B. VINBERG, Discrete linear groups that are generated by reflections (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **35** (1971), 1072–1112.

Gye-Seon Lee

Department of Mathematics, Sungkyunkwan University, Suwon, South Korea

E-mail address: `gyeseonlee@skku.edu`

Ludovic Marquis

Univ Rennes, CNRS, IRMAR - UMR 6625, F-35000 Rennes, France

E-mail address: `ludovic.marquis@univ-rennes1.fr`

Stefano Riolo

Dipartimento di Matematica, Università di Pisa, Italy

E-mail address: `stefano.riolo@dm.unipi.it`

Received on June 17, 2020.

Accepted on April 6, 2021.