

On the Local-indicability Cohen-Lyndon Theorem

Yago Antolín, Warren Dicks and Peter A. Linnell

February 10, 2010

Abstract

For a group H and a subset X of H , we let ${}^H X$ denote the set $\{h x h^{-1} \mid h \in H, x \in X\}$, and when X is a free-generating set of H , we say that the set ${}^H X$ is a *Whitehead subset* of H .

For a group F and an element r of F , we say that r is *Cohen-Lyndon aspherical* in F if ${}^F \{r\}$ is a Whitehead subset of the subgroup of F that is generated by ${}^F \{r\}$.

In 1963, D. E. Cohen and R. C. Lyndon independently showed that in each free group each non-trivial element is Cohen-Lyndon aspherical. Their proof used the celebrated induction method devised by W. Magnus in 1930 to study one-relator groups.

In 1987, M. Edjvet and J. Howie showed that if A and B are locally indicable groups, then each cyclically reduced element of $A*B$ that does not lie in $A \cup B$ is Cohen-Lyndon aspherical in $A*B$. Their proof used the Cohen-Lyndon Theorem.

Using Bass-Serre Theory and the Edjvet-Howie Theorem, one can deduce the Local-indicability Cohen-Lyndon Theorem: if F is a locally indicable group and T is an F -tree with trivial edge stabilizers, then each element of F that fixes no vertex of T is Cohen-Lyndon aspherical in F . Conversely, the Cohen-Lyndon Theorem and the Edjvet-Howie Theorem are immediate consequences of the Local-indicability Cohen-Lyndon Theorem.

In this article, we give a detailed review of Howie induction and arrange the arguments of Edjvet and Howie into a Howie-inductive proof of the Local-indicability Cohen-Lyndon Theorem that does not use Magnus induction or the Cohen-Lyndon Theorem.

We conclude with a review of some standard applications of Cohen-Lyndon asphericity.

2000 Mathematics Subject Classification. Primary: 20E08; Secondary: 20E05, 20E07, 20J06.

Key words. Cohen-Lyndon asphericity, locally indicable groups, groups acting on trees, staggerability

1 Outline

1.1 Notation. Let F be a multiplicative group, fixed throughout the article.

The disjoint union of two sets X and Y will be denoted by $X \vee Y$.

By a *transversal* for a (left or right) F -action on a set X we mean a subset of X which contains exactly one element of each F -orbit of X ; by the axiom of choice, transversals always exist.

For elements x, y of F , we write $\bar{x} := x^{-1}$, ${}^x y := xyx$, $[x, y] := xyx\bar{y}$, and $\mathbf{C}_F(x) := \{f \in F \mid f x = x\}$. For any subgroup H of F , we say that x, y are *H -conjugate* if there exists some $h \in H$ such that ${}^h x = y$. Conjugation actions will always be left actions in this article.

If R and X are subsets of F , we let $\langle R \rangle$ denote the subgroup of F generated by R , and we write ${}^X R := \{x r \mid r \in R, x \in X\}$ and $F/\langle R \rangle := F/\langle {}^F R \rangle$. If R consists of a single element r , we write simply $\langle r \rangle$, ${}^X r$ and $F/\langle r \rangle$, respectively. We say that X is a *free-generating set* of F if the induced group homomorphism $\langle X \mid \rangle \rightarrow F$ is bijective.

A subset Y of F is said to be a *Whitehead subset* of F if there exists some free-generating set X of F such that ${}^F X = Y$, that is, Y is closed under the F -conjugation action and some transversal for the F -conjugation action on Y is a free-generating set of F .

Borrowing terminology from [3], we say that an element r of F is *Cohen-Lyndon aspherical* in F if ${}^F r$ is a Whitehead subset of $\langle {}^F r \rangle$. Thus, r is Cohen-Lyndon aspherical in F if and only if some transversal for the $\langle {}^F r \rangle$ -conjugation action on ${}^F r$ is a free-generating set of $\langle {}^F r \rangle$. Thus, r is Cohen-Lyndon aspherical in F if and only if there exists some subset X of F such that X is a transversal for the $(\langle {}^F r \rangle \mathbf{C}_F(r))$ -action on F by multiplication on the right and, moreover, ${}^X r$ is a free-generating set of $\langle {}^F r \rangle$. Here, $X \neq \emptyset$ and $r \neq 1$ and the map $X \rightarrow {}^X r$, $x \mapsto x r$, is bijective.

We say that a subset R of F is *Cohen-Lyndon aspherical in F* if no two distinct elements of R are F -conjugate and ${}^F R$ is a Whitehead subset of $\langle {}^F R \rangle$. Thus, R is Cohen-Lyndon aspherical in F if and only if no two distinct elements of R are F -conjugate and some transversal for the $\langle {}^F R \rangle$ -conjugation action on ${}^F R$ is a free-generating set of $\langle {}^F R \rangle$. Thus, R is Cohen-Lyndon aspherical in F if and only if no two distinct elements of R are F -conjugate and there exists some family $(X_r \mid r \in R)$ of subsets of F with the properties that, for each $r \in R$, X_r is a transversal for the $(\langle {}^F R \rangle \mathbf{C}_F(r))$ -action on F by multiplication on the right and, moreover, $\bigcup_{r \in R} ({}^{X_r} r)$ is a free-generating set of $\langle {}^F R \rangle$.

A group is said to be *indicable* if it is trivial or it has some quotient that is infinite and cyclic. A group is said to be *locally indicable* if all its finitely generable subgroups are indicable. For example, all free groups are locally indicable. \square

In [4, Theorem 4.1], D. E. Cohen and R. C. Lyndon proved the following.

The Cohen-Lyndon Theorem. *If F is a free group, then each non-trivial element of F is Cohen-Lyndon aspherical in F .* \square

Both the proof in [4] and its simplification by A. Karrass and D. Solitar [12, Theorem 2] use the famous induction method that was devised by W. Magnus in 1930 to study one-relator groups.

In [6, Theorem 1.1], M. Edjvet and J. Howie used the Cohen-Lyndon Theorem and Karrass-Solitar reduction to prove the following.

The Edjvet-Howie Theorem. *If A and B are locally indicable groups, then each cyclically reduced element of $A*B$ that does not lie in $A \cup B$ is Cohen-Lyndon aspherical in $A*B$.* \square

By applying Bass-Serre Theory, one can deduce the following.

The Local-indicability Cohen-Lyndon Theorem. *If F is a locally indicable group and T is an F -tree with trivial edge stabilizers, then each element of F that fixes no vertex of T is Cohen-Lyndon aspherical in F .* \square

1.2 Example. Let X be a set and let $F = \langle X \mid \ \rangle$.

Let T be the F -graph with vertex set F and edge set $F \times X$ such that each edge $(f, x) \in F \times X$ has initial vertex f and terminal vertex fx . In a natural way, F acts on T . The stabilizers are trivial. It is well known that T is a tree; see, for example, [5, Theorem I.7.6].

Thus, the Cohen-Lyndon Theorem is the case of the Local-indicability Cohen-Lyndon Theorem where F acts freely on T . \square

1.3 Example. Let A and B be locally indicable groups and let $F = A*B$.

Let T be the F -graph with vertex set $(F/A) \vee (F/B)$ and edge set F such that each edge $f \in F$ has initial vertex fA and terminal vertex fB . In a natural way F acts on T . The edge stabilizers are trivial and the elements of F which fix vertices of T are the F -conjugates of the elements of $A \cup B$. It can be shown that T is a tree; see, for example, [5, Theorem I.7.6].

Thus, the Edjvet-Howie Theorem is the case of the Local-indicability Cohen-Lyndon Theorem where T has only one edge F -orbit and two vertex F -orbits. \square

In his work on locally indicable groups in [8], [9], [11], Howie developed a powerful induction technique that amounts to being given certain information about a group,

and then choosing an appropriate finitely generable subgroup which contains the given information,

and then choosing an appropriate normal subgroup of the finitely generable subgroup for which the quotient group is infinite and cyclic, and translating the given information into information about the normal subgroup,

and repeating this two-step cycle as often as possible. This simple procedure has many applications. Magnus induction has a formally similar format but requires a free-generating set and more careful choices at each step.

In this article, we give a Bass-Serre-theoretical survey of Howie induction and re-arrange the arguments used by Edjvet and Howie in [6] into a Howie-inductive proof of the Local-indicability Cohen-Lyndon Theorem that does not use Magnus induction or the Cohen-Lyndon Theorem.

In Section 2, we introduce definitions concerning staggerable subsets, strongly staggerable subsets, and other concepts that we shall be using.

In Section 3, we describe the finite descending chain of subgroups of F used in Howie induction. We then see that the staggerable conditions can be moved all the way down the chain.

In Section 4, we find that at the bottom of any of the chains of Section 3 the conditions of staggerability and local indicability interact to produce Whitehead subsets. This section reproduces parts of Appendix A of [1] with some modifications to suit the current applications.

In Section 5, we recall some technical results of Cohen-Lyndon, of Karrass-Solitar, and of Edjvet-Howie. These results then allow us to take information that was spontaneously generated at the bottom of the chain and move it all the way back up the chain; the Local-indicability Cohen-Lyndon Theorem then follows.

Corollary 2.2 of [4] gives a sufficient condition for a subset of a free group F to be Cohen-Lyndon aspherical in F . In Section 6, we find that the preceding machinery implies the more general result that, for any locally indicable group F and any subset R of F , if there exists some F -tree T with trivial edge stabilizers such that R is strongly T -staggerable modulo F , then R is Cohen-Lyndon aspherical in F . Standard arguments then yield consequences concerning the quotient group $G := F/\langle\langle R \rangle\rangle$; for example, one obtains information about the torsion subgroups of G and the higher homology groups of G .

2 Staggerability

In this section, we introduce definitions concerning staggerability and other concepts that we shall be using. Recall that F is a multiplicative group.

2.1 Notation. For any set X , we let $|X|$ denote the cardinal of X , and we let $\mathbb{Z}X$ and $\mathbb{Z}[X]$ denote the \mathbb{Z} -module that is free on X .

By an *ordering* of a set X , we shall mean a binary relation which *totally* orders X .

We will find it useful to have a notation for intervals in \mathbb{Z} that is different from the notation for intervals in \mathbb{R} . Let $i, j \in \mathbb{Z}$. We write $[i \uparrow j] := \{k \in \mathbb{Z} \mid k \geq i \text{ and } k \leq j\}$, and $]-\infty \uparrow j] := \{k \in \mathbb{Z} \mid k \leq j\}$, and $[i \uparrow \infty[:= \{k \in \mathbb{Z} \mid k \geq i\}$.

We shall define families of subscripted symbols by using the following convention. Let v be a symbol. For each $k \in \mathbb{Z}$, we let v_k denote the ordered pair (v, k) , and, for each subset I of \mathbb{Z} , we let $v_I := (v_k \mid k \in I)$.

For two subsets Y, Z of a set X , the complement of $Y \cap Z$ in Y will be denoted by $Y - Z$ (and not by $Y \setminus Z$ since we let $F \setminus Z$ denote the set of F -orbits of a left F -set Z).

For any subset Y of a left F -set X , we write $\text{glue}(F, Y) := \{f \in F \mid fY \cap Y \neq \emptyset\}$.

We write $F' := \langle \{[x, y] \mid x, y \in F\} \rangle \trianglelefteq F$ and $F^{\text{ab}} := F/F'$, the abelianization of F . □

2.2 Definitions. Let r be an element of F .

We say that r has a unique root in F if $r = 1$ or r lies in a unique maximal infinite cyclic subgroup of F .

If $r \neq 1$ and r lies in a unique maximal infinite cyclic subgroup C of F , we define *the unique root of r in F* , denoted by $\sqrt[r]{r}$, to be the unique generator of C of which r is a positive power. We say that 1 is *the unique root of 1*, and we define $\sqrt[1]{1} := 1$.

If $r \neq 1$ and $\mathbf{C}_F(r)$ is infinite and cyclic, then r has a unique root in F and $\langle \sqrt[r]{r} \rangle = \mathbf{C}_F(r)$.

We say that a subset R of F has *unique roots in F* if every element of R has a unique root in F , in which case we let $\sqrt[R]{R}$ denote the set of these unique roots in F . □

We now define the staggerability concepts we need. We shall use [5] as our reference for Bass-Serre Theory.

2.3 Definitions. Let $T = (T, VT, ET, \iota, \tau)$ be an F -tree and suppose that ET is F -free.

(i). Let r be an element of F that fixes no vertex of T .

There exists a unique minimal $\langle r \rangle$ -subtree of T , which is denoted by $\text{axis}(r)$ and has the form of a real line shifted by r ; see, for example, [5, Proposition I.4.11]. We write $E\text{axis}(r) := E(\text{axis}(r))$, and we shall be particularly interested in the finite set $F \setminus (F(E\text{axis}(r))) := \{Fe \mid e \in E\text{axis}(r)\} \subseteq F \setminus ET$.

For all $f \in F$, $\text{axis}(f r) = f \text{axis}(r)$ and $F \setminus (F(E\text{axis}(f r))) = F \setminus (F(E\text{axis}(r)))$.

For all $n \in \mathbb{Z} - \{0\}$, $\text{axis}(r^n) = \text{axis}(r)$.

Since F acts freely on ET , it can be shown that $\mathbf{C}_F(r)$ acts freely on $\text{axis}(r)$, and, by Bass-Serre Theory, $\mathbf{C}_F(r)$ is infinite and cyclic. Here, r has a unique root in F and $\langle r \rangle \subseteq \langle \sqrt[r]{r} \rangle = \mathbf{C}_F(r) \subseteq \text{glue}(F, E\text{axis}(r))$; notice that if $\text{glue}(F, E\text{axis}(r)) = \langle r \rangle$, then $\sqrt[r]{r} = r$.

(ii). If $<$ is some (total) ordering of ET and R is some subset of F , we say that R is $(T, <)$ -staggered modulo F if the following three conditions hold.

- (S1). Each F -orbit in ET is an interval in $(ET, <)$; here, there exists a unique ordering of $F \setminus ET$, again denoted by $<$, with the property that for all $e, e' \in ET$, $Fe < Fe'$ if and only if $e < e'$ and $Fe \neq Fe'$.
- (S2). Each element of R fixes no vertex of T .
- (S3). For each $(r_1, r_2) \in R \times R$, exactly one of the following three conditions holds.
 - (a) ${}^F r_1 = {}^F r_2$.
 - (b) In $(F \setminus ET, <)$,
 $\min(F \setminus (F(E\text{axis}(r_1)))) < \min(F \setminus (F(E\text{axis}(r_2))))$ and
 $\max(F \setminus (F(E\text{axis}(r_1)))) < \max(F \setminus (F(E\text{axis}(r_2))))$.
 - (c) In $(F \setminus ET, <)$,
 $\min(F \setminus (F(E\text{axis}(r_2)))) < \min(F \setminus (F(E\text{axis}(r_1))))$ and
 $\max(F \setminus (F(E\text{axis}(r_2)))) < \max(F \setminus (F(E\text{axis}(r_1))))$.

We say that R is *strongly* $(T, <)$ -staggered modulo F if, moreover, the following four conditions hold.

- (S4). No two distinct elements of R are F -conjugate.
- (S5). $F \setminus T$ has a unique maximal subtree.
- (S6). $(F \setminus ET, <)$ is order isomorphic to an interval in \mathbb{Z} .
- (S7). Any two $<$ -consecutive edges in $(F \setminus ET, <)$ have a vertex in common in $F \setminus T$.

We have two types of examples in mind satisfying (S5), (S6) and (S7): cases where $F \setminus T$ has only one vertex and cases where $F \setminus T$ has the form of the real line.

(iii). A subset R of F is said to be T -staggerable modulo F if there exists some ordering $<$ of ET such that R is $(T, <)$ -staggered modulo F .

Notice that if R is T -staggerable modulo F , then R has unique roots in F .

Notice that if $|R| = 1$, then R is T -staggerable modulo F if and only if the unique element of R fixes no vertex of T .

A subset R of F is said to be *strongly* T -staggerable modulo F if there exists some ordering $<$ of ET such that R is strongly $(T, <)$ -staggered modulo F . \square

2.4 Remarks. In the case where F is free, the concept of a staggered subset appeared in Howie's article [11, p.642] and was a generalization, from the case where $|F \setminus VT| = 1$ and F is free, of a definition of a staggered presentation that is given in Lyndon and Schupp's book [14, p.152]. In the case where $|F \setminus VT| = 1$ and F is free, the above concept of a strongly staggered subset corresponds to a definition that is given in [14, p.104] of what could reasonably be called a strongly staggered presentation but is somewhat confusingly called a "staggered presentation"; such strongly staggered presentations arise unnamed in the hypotheses of Lyndon's (Non-Simple) Identity Theorem [13, Section 7] and Cohen and Lyndon's Corollary 2.2 [4].

In Corollary 6.2 below, we show that, for any locally indicable group F and any subset R of F , if there exists some F -tree T with trivial edge stabilizers such that R is strongly T -staggerable modulo F , then R is Cohen-Lyndon aspherical in F . The case of Corollary 6.2 where $|F \setminus VT| = 1$ and F is free is precisely the Cohen-Lyndon result for strongly staggered presentations [4, Corollary 2.2]; see also [14, Proposition III.11.1]. \square

We shall be using the following well-known observation.

2.5 Remark. Let X be a generating set of F , let T be an F -tree, and let Y be a subtree of T such that, for some vertex v of T , $Y \supseteq \{v\} \cup Xv$, or, more generally, such that $X \subseteq \text{glue}(F, Y)$.

Consider the F -subforest FY of T , and let T' denote the component of FY that contains Y . The set $\{f \in F \mid fY \subseteq T'\}$ is closed under right multiplication by elements of $X \cup X^{-1}$ and contains 1, hence it is all of F . Thus $FY = T'$. Hence FY is an F -subtree of T . Moreover, the map $Y \rightarrow F \setminus T'$, $y \mapsto Fy$, is surjective. \square

3 Moving information down a chain

In this section, without mentioning local indicability, we shall see how to move a staggerable subset S down a finite chain of subgroups to a finitely generable group whose abelianization is virtually generated by S . The finite descending chain of subgroups is essentially the same chain of subgroups considered by Howie in his tower arguments in [9] and [10].

We shall often consider the following situation.

3.1 Setting. Let S and Φ be finite subsets of F such that $\langle S \cup \Phi \rangle / \langle S \rangle$ has no infinite, cyclic quotients.

Let T be an F -tree with trivial edge stabilizers.

Choose an arbitrary vertex v of T . Let Y be the smallest subtree of T which contains $\{v\} \cup Sv \cup \Phi v$. Then Y is finite. Since F acts freely on ET , $\text{glue}(F, EY)$ is finite. We set $S^+ := S \cup \Phi \cup \text{glue}(F, EY)$. Then S^+ is a finite subset of $\text{glue}(F, Y)$. We set $\nu := |S^+| + 1$. For each subgroup H of F , we set $H^\dagger := \langle S^+ \cap H \rangle$. Notice that if H contains $S \cup \Phi$, then H^\dagger also contains $S \cup \Phi$.

We set $F_0 := F$. Suppose that, for some $n \in [0 \uparrow(\nu-1)]$, we have a subgroup F_n of F containing S . If $F_n^\dagger / \langle S \rangle$ has no infinite, cyclic quotients, we choose $F_{n+1} := F_n^\dagger$; otherwise, we choose F_{n+1} to be an arbitrary normal subgroup of F_n^\dagger which contains S such that F_n^\dagger / F_{n+1} is infinite and cyclic. After ν such steps, we will have recursively chosen a finite, monotonically decreasing sequence $F_{[0 \uparrow \nu]}$ of subgroups of F that contain S . \square

3.2 Proposition. *In Setting 3.1, the following hold.*

- (i). $F_\nu = \langle S^+ \cap F_\nu \rangle$ and $S \cup \Phi \subseteq F_\nu$ and $F_\nu / \langle S \rangle$ has no infinite, cyclic quotients.
- (ii). If S is T -staggerable modulo F , then S is T -staggerable modulo F_ν .

Proof. (i). If $n \in [0 \uparrow(\nu-1)]$ and $S \cup \Phi \subseteq F_n$, then $S \cup \Phi \subseteq F_n^\dagger$ and, since $\langle S \cup \Phi \rangle / \langle S \rangle$ has no infinite, cyclic quotients, $S \cup \Phi \subseteq F_{n+1}$. By induction, $S \cup \Phi \subseteq F_\nu$.

Consider any $n \in [0 \uparrow(\nu-1)]$. If $F_{n+1} < F_n^\dagger$ ($= \langle S^+ \cap F_n \rangle \leq F_n$), then $F_{n+1} \not\subseteq S^+ \cap F_n$, and, hence, $S^+ \cap F_{n+1} \subset S^+ \cap F_n$; notice that the number of those n for which the latter happens is at most $|S^+| = \nu - 1$. It follows that there exists some $\mu \in [0 \uparrow(\nu-1)]$ such that $F_{\mu+1} = F_\mu^\dagger$. Then $F_\mu^\dagger / \langle S \rangle$ has no infinite, cyclic quotients, and

$$F_\mu^\dagger = \langle S^+ \cap F_\mu \rangle = \langle S^+ \cap F_\mu^\dagger \rangle = \langle S^+ \cap F_{\mu+1} \rangle = F_{\mu+1}^\dagger.$$

It then follows that the sequence $F_{[\mu \uparrow \nu]}^\dagger$ is constant, and then (i) follows.

(ii). By hypothesis, S is T -staggerable modulo F_0 ($= F$).

Let n be an element of $[0 \uparrow(\nu-1)]$ such that S is T -staggerable modulo F_n . By induction, it suffices to show that S is T -staggerable modulo F_{n+1} .

There exists some ordering $<$ of ET such that S is $(T, <)$ -staggered modulo F_n . By altering the ordering $<$ of ET within each F_n -orbit, we can arrange that every F_n^\dagger -orbit is an interval with respect to $<$. We claim that S is now $(T, <)$ -staggered modulo F_n^\dagger . It suffices to show that if two elements of S are F_n -conjugate, then they are F_n^\dagger -conjugate. Suppose then that we have $f \in F_n$ and $r, {}^f r \in S \subseteq F_n^\dagger$. Now $\text{axis}(r) \subseteq \langle r \rangle(Y) \subseteq F_n^\dagger(Y)$ and $f \text{axis}(r) = \text{axis}({}^f r) \subseteq \langle {}^f r \rangle(Y) \subseteq F_n^\dagger(Y)$. Hence, $f \in \text{glue}(F_n, F_n^\dagger(EY))$. Thus $F_n^\dagger f F_n^\dagger$ meets

$$\text{glue}(F_n, EY) \subseteq S^+ \cap F_n \subseteq \langle S^+ \cap F_n \rangle = F_n^\dagger.$$

Hence $f \in F_n^\dagger$. Thus S is $(T, <)$ -staggered modulo F_n^\dagger .

If $F_{n+1} = F_n^\dagger$, then S is $(T, <)$ -staggered modulo F_{n+1} as desired. Thus, we may assume that F_n^\dagger / F_{n+1} is infinite and cyclic. There exists some $z \in F_n^\dagger$ such that $z F_{n+1}$ generates F_n^\dagger / F_{n+1} . Then $F_n^\dagger = \langle z \rangle F_{n+1}$. Since F_n^\dagger / F_{n+1} is infinite, $\langle z \rangle \cap F_{n+1} = \{1\}$. Within each F_n^\dagger -orbit in ET , the F_{n+1} -orbits are permuted by z and form a single $\langle z \rangle$ -orbit. By altering the ordering $<$ of ET within each F_n^\dagger -orbit, we can arrange that every F_{n+1} -orbit is an interval with respect to $<$, and such that z moves each F_{n+1} -orbit to the next $<$ -largest orbit. It suffices to examine two elements of S which are F_n^\dagger -conjugate. Suppose then that we have $f \in F_{n+1}$ and $i \in \mathbb{Z}$ and $r \in S$ such that $z^i f r \in S$. If $i = 0$, then r and $z^i f r$ are F_{n+1} -conjugate, which is one of the desired possibilities. We now assume that $i > 0$; the case where $i < 0$ is similar. Now

$$F_{n+1} \setminus (F_{n+1}(E \text{axis}(z^i f r))) = F_{n+1} \setminus (F_{n+1}(z^i f E \text{axis}(r))) = z^i (F_{n+1} \setminus (F_{n+1}(E \text{axis}(r)))).$$

Thus,

$$\min(F_{n+1} \setminus (F_{n+1}(E \text{axis}(z^i f r)))) = z^i \min(F_{n+1} \setminus (F_{n+1}(E \text{axis}(r)))) > \min(F_{n+1} \setminus (F_{n+1}(E \text{axis}(r)))),$$

$$\max(F_{n+1} \setminus (F_{n+1}(E\text{axis}(z^i f_r)))) = z^i \max(F_{n+1} \setminus (F_{n+1}(E\text{axis}(r)))) > \max(F_{n+1} \setminus (F_{n+1}(E\text{axis}(r)))).$$

Hence, S is $(T, <)$ -staggered modulo F_{n+1} , as desired. \square

4 New information generated at the bottom of a chain

In this section, we review, with some changes, the main results of [1, Appendix A]. The following is a minor extension of [1, Lemma A.2.1].

4.1 Lemma. *Let F be a finitely generable, locally indicable group, let T be an F -tree with trivial edge stabilizers, and let S be a subset of F such that S is T -staggerable modulo F and $\sqrt[F]{S} = S$. Then the following hold.*

- (i). $F/\langle\langle S \rangle\rangle$ is indicable.
- (ii). If $F/\langle\langle S \rangle\rangle$ is trivial, then, for each $r \in S$, $\text{glue}(F, E\text{axis}(r)) = \langle r \rangle$, and F acts freely on T , and ${}^F S$ is a Whitehead subset of F .

Proof. Recall from Definitions 2.3 that S has unique roots in F .

Without loss of generality, we may replace S with ${}^F S$, and we then have ${}^F S = S$.

By hypothesis, there exists some ordering $<$ of ET such that S is $(T, <)$ -staggered modulo F .

Let X be a finite generating set of F , let v be a vertex of T , let Y be the smallest subtree of T containing $\{v\} \cup Xv$, and let $T' := FY$. By Remark 2.5, T' is an F -subtree of T and $F \setminus T'$ is finite. Consider any $r \in S$. Then T' is an $\langle r \rangle$ -subtree of T . Since $\text{axis}(r)$ is the unique smallest $\langle r \rangle$ -subtree of T , we see that T' contains $\text{axis}(r)$. It follows that S is $(T', <)$ -staggered modulo F . It now suffices to prove the result with T' in place of T . Thus, we may assume that $F \setminus T$ is finite.

By induction, we may assume that the result holds for all smaller values of $|F \setminus ET|$.

Since F is a finitely generable, locally indicable group, F is indicable, and it can be seen that (i) and (ii) hold when S is empty. Thus we may assume that S is non-empty. It then follows from Definitions 2.3 that $|F \setminus ET| \geq 1$. In particular, there exists some $e_{\max} \in \bigcup_{r \in S} E\text{axis}(r)$ such that, in $(F \setminus ET, <)$,

$$Fe_{\max} = \max\{Fe \mid e \in \bigcup_{r \in S} E\text{axis}(r)\}.$$

There then exists some $r_{\max} \in S$ such that $e_{\max} \in E\text{axis}(r_{\max})$, and, by the definition of $(T, <)$ -staggered modulo F , Fe_{\max} does not meet the axis of any element of $S - {}^F(r_{\max})$. Thus there exists some pair (r, e) , for example $(r, e) = (r_{\max}, e_{\max})$, such that the following hold.

$$r \in S, e \in E\text{axis}(r), \text{ and } Fe \text{ does not meet the axis of any element of } S - {}^F r. \quad (1)$$

$$\text{If } \text{glue}(F, E\text{axis}(r)) \neq \langle r \rangle, \text{ then } (r, e) = (r_{\max}, e_{\max}). \quad (2)$$

We consider the F -forest $T - Fe$. Let T_l denote the component of $T - Fe$ containing ιe , and let T_τ denote the component of $T - Fe$ containing τe . Let F_l denote the F -stabilizer of $\{T_l\}$, and let F_τ denote the F -stabilizer of $\{T_\tau\}$, where we are using set brackets to emphasize that we want to consider each component of $T - Fe$ as a single element. Let $S_l := S \cap F_l$ and $S_\tau := S \cap F_\tau$. By (1), for each $r' \in S - {}^F r$, $\text{axis}(r')$ lies in $T - Fe$ and hence lies in a component of $T - Fe$. It follows that ${}^F S_l \cup {}^F S_\tau \cup {}^F r$ is all of S . Notice that if $F\{T_l\} = F\{T_\tau\}$ then ${}^F S_l = {}^F S_\tau$.

By applying the Bass-Serre Structure Theorem to the F -tree whose vertices are the components of $T - Fe$ and whose edge set is Fe , with fe joining fT_l to fT_τ ($f \in F$), we find that

$$F = \begin{cases} F_l * F_\tau & \text{if } F\{T_l\} \neq F\{T_\tau\}, \\ F_l * \langle f \mid \rangle & \text{if } f \in F \text{ and } fT_l = T_\tau. \end{cases} \quad (3)$$

Hence,

$$F/\langle\langle S - {}^F r \rangle\rangle = \begin{cases} (F_l/\langle\langle S_l \rangle\rangle) * (F_\tau/\langle\langle S_\tau \rangle\rangle) & \text{if } F\{T_l\} \neq F\{T_\tau\}, \\ (F_l/\langle\langle S_l \rangle\rangle) * \langle f \mid \rangle & \text{if } f \in F \text{ and } fT_l = T_\tau. \end{cases} \quad (4)$$

Case 1. Both $F_l/\langle S_l \rangle$ and $F_\tau/\langle S_\tau \rangle$ have infinite, cyclic quotients.

By (4), $F/\langle S - F_r \rangle$ has a rank-two, free-abelian quotient. On incorporating r , we see that $F/\langle S \rangle$ has an infinite, cyclic quotient, and, hence, (i) and (ii) hold in this case.

Case 2. One of $F_l/\langle S_l \rangle$, $F_\tau/\langle S_\tau \rangle$ does not have an infinite, cyclic quotient.

By reversing the orientation of every edge of T , if necessary, we may assume that $F_l/\langle S_l \rangle$ does not have an infinite, cyclic quotient.

By the induction hypothesis applied to (F_l, T_l, S_l) , we see that $F_l/\langle S_l \rangle$ is trivial, and that F_l acts freely on T_l , and that some transversal for that F_l -action on S_l by conjugation is a (finite) free-generating set of F_l , and that, for each $r_l \in S_l$, $\text{glue}(F_l, E\text{axis}(r_l)) = \langle r_l \rangle$, and, hence, $\text{glue}(F, E\text{axis}(r_l)) = \langle r_l \rangle$.

By inverting every element of S , if necessary, we may assume the following.

$$\text{There exists some segment of axis}(r) \text{ of the form } e, p, re. \quad (5)$$

Consider the case where $F\{T_l\} \neq F\{T_\tau\}$. By (5), we have a path $\bar{r}p$ in $\text{axis}(r)$ from $\bar{r}\tau e \in \bar{r}T_\tau \neq T_l$ to $\iota e \in T_l$. Now $\bar{r}p$ necessarily enters T_l through an oriented edge of the form ge^{-1} for some $g \in F_l$, and g then lies in $\text{glue}(F_l, E\text{axis}(r)) = \langle r \rangle$. This proves the following.

$$\text{If } \text{glue}(F_l, E\text{axis}(r)) \subseteq \langle r \rangle \text{ then } F\{T_l\} = F\{T_\tau\}. \quad (6)$$

Consider the case where S_l is empty. Here, $F_l = F_l/\langle S_l \rangle = \{1\}$. By (6), $F\{T_l\} = F\{T_\tau\}$, and, then, by (3), there exists some $t \in F$ such that $F = \langle t \mid \ \rangle$. Now $\{t, \bar{t}\} = \sqrt[F]{F} - \{1\} \supseteq S = \{r\}$. Hence, $F = \langle r \mid \ \rangle$, and, hence, $\text{glue}(F, E\text{axis}(r)) = \langle r \rangle$. This proves the following.

$$\text{If } \text{glue}(F, E\text{axis}(r)) \neq \langle r \rangle \text{ then } S_l \text{ is non-empty}. \quad (7)$$

Case 2.1. $\text{glue}(F, E\text{axis}(r)) = \langle r \rangle$.

Here, by (6), $F\{T_l\} = F\{T_\tau\}$. Hence, $F(VT_l) = VT$ and $S = {}^F S_l \cup F_r$. Consider any $v \in VT$. We wish to show that $F_v = 1$, and, we may assume that $v \in VT_l$. Here, $F_v \leq F_l$, and, since F_l acts freely on T_l , $F_v = 1$, as desired. Thus, F acts freely on T . In (5), the path p from τe to $r\tau e$ in $\text{axis}(r)$ does not meet Fe since $\text{glue}(F, E\text{axis}(r)) = \langle r \rangle$, and, hence, p stays within T_τ , and, hence, $r\tau e \in T_\tau$. Thus, $rT_l = T_\tau$, and, by (3), $F = F_l * \langle r \mid \ \rangle$. Since $F_l/\langle S_l \rangle$ is trivial, we see that $F/\langle S \rangle$ is trivial, and, hence, $F/\langle S \rangle$ is indicable. Here all the required conclusions hold.

Case 2.2. $\text{glue}(F, E\text{axis}(r)) \neq \langle r \rangle$.

By (7), S_l is non-empty, and, hence, there exists some $e_l \in \bigcup_{r_l \in S_l} E\text{axis}(r_l)$ such that

$$Fe_l = \min\{Fe \mid e \in \bigcup_{r_l \in S_l} E\text{axis}(r_l)\}.$$

There then exists some $r_l \in S_l$ such that $e_l \in E\text{axis}(r_l)$, and we then know that $\text{glue}(F, E\text{axis}(r_l)) = \langle r_l \rangle$. By (2), $(r, e) = (r_{\max}, e_{\max})$. Using the definition of $(T, <)$ -staggered modulo F , one can show that Fe_l does not meet the axis of any element of ${}^F r$, and, similarly, Fe_l does not meet the axis of any element of ${}^F S_l - F_{r_l}$. It is clear that if ${}^F S_\tau \neq {}^F S_l$, then Fe_l does not meet the axis of any element of ${}^F S_\tau$. Hence, Fe_l does not meet the axis of any element of $S - F_{r_l}$. We now replace (r, e) with (r_l, e_l) in (1), (2), and we then find that the same argument as before now terminates in Case 1 or Case 2.1. Hence, all the desired conclusions hold.

This completes the proof. \square

4.2 Corollary. *In Setting 3.1, suppose that F is locally indicable and that S is T -staggerable modulo F and that $\sqrt[F]{S} = S$. Then $S \cup \Phi \subseteq F_\nu$ and F_ν acts freely on T and ${}^{F_\nu} S$ is a Whitehead subset of F_ν .*

Proof. Recall from Definitions 2.3 that S has unique roots in F .

By Proposition 3.2(i), F_ν is finitely generated and $S \cup \Phi \subseteq F_\nu$ and $F_\nu/\langle S \rangle$ has no infinite, cyclic quotients. By Proposition 3.2(ii), S is T -staggerable modulo F_ν . By Lemma 4.1(i), $F_\nu/\langle S \rangle$ is indicable, and, hence, trivial. Now the result holds by Lemma 4.1(ii). \square

The next result is repeated from Corollary A.2.3 of [1]; Section A.3 of [1] describes how these results continue work of Magnus, Brodskii, Howie, Short and others.

4.3 Theorem. *Let F be a locally indicable group and let T be an F -tree with trivial edge stabilizers and let R be a subset of F that is T -staggerable modulo F .*

Then the following hold.

- (i). (The Local-indicability Freiheitssatz) $\langle {}^F R \rangle$ acts freely on T , and, hence, $\langle {}^F R \rangle$ is a free group and the F -stabilizer of each vertex of T embeds in $F/\langle R \rangle$ under the natural map.
- (ii). $F/\langle \sqrt[F]{R} \rangle$ is locally indicable.

Proof. Since $\langle {}^F R \rangle \leq \langle {}^F(\sqrt[F]{R}) \rangle$, we may replace R with $\sqrt[F]{R}$ and then we have $\sqrt[F]{R} = R$.

(i). Let S be an arbitrary finite subset of ${}^F R$ and let Φ be the empty set. We may then assume that we are in Setting 3.1. By Corollary 4.2, $\langle S \rangle \leq F_\nu$ and F_ν acts freely on T . Hence, $\langle S \rangle$ acts freely on T , and, since S is an arbitrary finite subset of ${}^F R$, we see that $\langle {}^F R \rangle$ acts freely on T , that is, the F -stabilizer of each vertex of T embeds in $F/\langle R \rangle$ under the natural map. By Reidemeister's Theorem, or by Bass-Serre Theory, $\langle {}^F R \rangle$ is a free group.

(ii). Let H be an arbitrary finitely generable subgroup of $F/\langle {}^F R \rangle$. It suffices to show that H is indicable. We may assume that H has no infinite, cyclic quotients. Then H^{ab} is finite. Let d denote the exponent of H^{ab} . There exists some finite subset Φ of F such that $H = (\langle \Phi \rangle \langle {}^F R \rangle) / \langle {}^F R \rangle$. Now $H' = (\langle \Phi \rangle' \langle {}^F R \rangle) / \langle {}^F R \rangle$ and we see that $\Phi^d \subseteq \langle \Phi \rangle' \langle {}^F R \rangle$. Hence, there exists some finite subset S of ${}^F R$ such that the finite set Φ^d lies in $\langle \Phi \rangle' \langle S \rangle$. Then, the abelian group $(\langle \Phi \cup S \rangle / \langle S \rangle)^{\text{ab}}$ has exponent at most d , and, hence, $\langle \Phi \cup S \rangle / \langle S \rangle$ has no infinite, cyclic quotients. We may then assume that we are in Setting 3.1. By Corollary 4.2, $\Phi \subseteq F_\nu = \langle {}^{F_\nu} S \rangle \leq \langle {}^F R \rangle$, and, hence, H is trivial, and hence, H is indicable. \square

4.4 Corollary (Brodskiĭ-Howie-Short). *Suppose that A and B are locally indicable groups and that r is an element of $A*B$. If no $A*B$ -conjugate of r lies in A , then the natural map embeds A in $(A*B)/\langle r \rangle$. \square*

4.5 Corollary (Magnus' Freiheitssatz). *Suppose that A and B are free groups and that r is an element of $A*B$. If no $A*B$ -conjugate of r lies in A , then the natural map embeds A in $(A*B)/\langle r \rangle$. \square*

5 Moving the new information back up the chain

In this section, we recover the Local-indicability Cohen-Lyndon Theorem.

5.1 Lemma. *Let Y be a non-empty subset of F .*

Then Y is a Whitehead subset of F if and only if all of the following hold: each $y \in Y$ freely generates $\mathbf{C}_F(y)$; Y generates F ; Y is closed under the F -conjugation action; and, Y with the F -conjugation action is the vertex set of an F -tree with trivial edge stabilizers.

Proof. Suppose that Y is a Whitehead subset of F , and let X be a free-generating set of F such that $Y = {}^F X$. Then Y generates F , Y is closed under the F -conjugation action, and each $y \in Y$ generates a non-trivial free factor of F and, hence, freely generates $\mathbf{C}_F(y)$. We can express F as the fundamental group of a tree of groups in which the edge groups are trivial and the family of vertex groups is $(\langle x \mid \ \ \ \rangle \mid x \in X)$. Let T denote the corresponding Bass-Serre tree. Thus $VT = \bigvee_{x \in X} F/\langle x \rangle \simeq \bigvee_{x \in X} {}^F x = {}^F X = Y$, and ET is some free F -set.

Conversely, suppose that Y generates F and that each $y \in Y$ freely generates $\mathbf{C}_F(y)$, and that Y is closed under the F -conjugation action, and that Y with the F -conjugation action is the vertex set of an F -tree T with trivial edge stabilizers. By the Bass-Serre Structure Theorem, there exists some graph of groups (\mathcal{F}, \bar{T}) and some maximal subtree \bar{T}_0 of \bar{T} such that F is the fundamental group of $(\mathcal{F}, \bar{T}, \bar{T}_0)$. By the centralizer condition, the family of vertex groups can be expressed as $(\langle x \mid \ \ \ \rangle \mid x \in X)$ for some transversal X for the F -conjugation action on Y . Since F is then generated by the F -conjugates of the vertex groups, it follows that $\bar{T} = \bar{T}_0$. Hence, F is the free product of the family of vertex groups. Thus, X is a free-generating set of F and $Y = {}^F X$, as desired. \square

5.2 Corollary. *Let R be a non-empty subset of F .*

Then R is Cohen-Lyndon aspherical in F if and only if all of the following hold: no two distinct elements of R are F -conjugate; each $r \in R$ freely generates $\mathbf{C}_{\langle {}^F R \rangle}(r)$; and, ${}^F R$ with the $\langle {}^F R \rangle$ -conjugation action is the vertex set of an $\langle {}^F R \rangle$ -tree with trivial edge stabilizers. \square

We now give a generalization of results of Cohen and Lyndon [4, Lemma 2.1] and Karrass and Solitar [12, Theorem 1] that is similar to a result of Chiswell, Collins and Huebschmann [3, Theorem 4.6].

5.3 Theorem. *Let (\mathcal{F}, Y) be a graph of groups with family of groups $(\mathcal{F}(y) \mid y \in Y)$ and family of edge maps $(\bar{t}_e: \mathcal{F}(e) \rightarrow \mathcal{F}(\tau_Y e) \mid e \in EY)$, let Y_0 be a maximal subtree of Y , and let F be the fundamental group of (\mathcal{F}, Y, Y_0) .*

Let $R := (r_v \mid v \in VY)$ be a family of elements of F with the property that, for each $v \in VY$, all of the following hold: $r_v \in \mathcal{F}(v)$; for each $e \in \iota_Y^{-1}(\{v\})$, $\langle \mathcal{F}^{(v)} r_v \rangle \cap \mathcal{F}(e) = \{1\}$; and, for each $e \in \tau_Y^{-1}(\{v\})$, $\langle \mathcal{F}^{(v)} r_v \rangle \cap \bar{t}_e \mathcal{F}(e) = \{1\}$.

Then, for each $v \in VY$, $\langle F R \rangle \cap \mathcal{F}(v) = \langle \mathcal{F}^{(v)} r_v \rangle$.

Let $U := \{v \in VY \mid r_v \neq 1\}$. Then $\{r_u \mid u \in U\}$ is Cohen-Lyndon aspherical in F if and only if, for each $u \in U$, r_u is Cohen-Lyndon aspherical in $\mathcal{F}(u)$.

Proof. For each $v \in VY$, let $\bar{\mathcal{F}}(v) := \mathcal{F}(v)/\langle r_v \rangle$, and, for each $e \in EY$, let $\bar{\mathcal{F}}(e) := \mathcal{F}(e)$. By the hypotheses on R , the edge maps for \mathcal{F} then induce injective maps which give a graph of groups $(\bar{\mathcal{F}}, Y)$ with family of groups $(\bar{\mathcal{F}}(y) \mid y \in Y)$. Let \bar{F} denote the fundamental group of $(\bar{\mathcal{F}}, Y, Y_0)$. Using the definition of fundamental groups, we then construct a natural group homomorphism $F \mapsto \bar{F}$, which is surjective and has kernel $\langle F R \rangle$. In particular, for each $v \in VY$, $\langle F R \rangle \cap \mathcal{F}(v) = \langle \mathcal{F}^{(v)} r_v \rangle$.

Let T denote the Bass-Serre tree for (\mathcal{F}, Y, Y_0) , and let \bar{T} denote the Bass-Serre tree for $(\bar{\mathcal{F}}, Y, Y_0)$. Using the definition of Bass-Serre trees, we construct a natural identification $\langle F R \rangle \backslash T = \bar{T}$. Using the Bass-Serre Structure Theorem for $\langle F R \rangle$ acting on T , we see that $\langle F R \rangle$ is the fundamental group of a tree of groups over \bar{T} , with trivial edge groups, and with family of vertex groups of the form $(\langle F R \rangle \cap {}^z \mathcal{F}(v) \mid v \in VY, z \in Z_v)$, where, for each $v \in VY$, Z_v is some transversal for the $\langle F R \rangle \mathcal{F}(v)$ -action on F by multiplication on the right. Thus $\langle F R \rangle = \ast_{v \in VY} \ast_{z \in Z_v} \langle {}^z \mathcal{F}(v) r_v \rangle = \ast_{u \in U} \ast_{z \in Z_u} \langle {}^z \mathcal{F}(u) r_u \rangle$.

We claim that, for each $u \in U$, $\mathbf{C}_F(r_u) = \mathbf{C}_{\mathcal{F}(u)}(r_u)$. If $f \in \mathbf{C}_F(r_u)$, then r_u fixes the path in T from $1\mathcal{F}(u)$ to $f\mathcal{F}(u)$, and, by the hypotheses, r_u fixes no edges of T . Hence the path under consideration is trivial, and, hence, $f \in \mathcal{F}(u)$, as claimed.

Suppose that, for each $u \in U$, r_u is Cohen-Lyndon aspherical in $\mathcal{F}(u)$, and let Y_u be a transversal for the $\langle \mathcal{F}^{(u)} r_u \rangle \mathbf{C}_{\mathcal{F}(u)}(r_u)$ -action on $\mathcal{F}(u)$ by multiplication on the right such that $Y_u r_u$ is a free-generating set of $\langle \mathcal{F}^{(u)} r_u \rangle$. Then $Z_u Y_u$ is a transversal for the $\langle F R \rangle \mathbf{C}_F(r_u)$ -action on F by multiplication on the right and $\bigvee_{u \in U} Z_u Y_u r_u$ is a free-generating set of $\langle F R \rangle$. Thus R is Cohen-Lyndon aspherical in F .

Conversely, suppose that R is Cohen-Lyndon aspherical in F and that u is an element of U . By Corollary 5.2, each $r \in R$ freely generates $\mathbf{C}_{\langle F R \rangle}(r)$, no two distinct elements of R are F -conjugate, and there exists some $\langle F R \rangle$ -tree T' with vertex set ${}^F R$ and with trivial edge stabilizers. By Corollary 5.2, it remains to show that $\mathcal{F}^{(u)} r_u$ is the vertex set of some $\langle \mathcal{F}^{(u)} r_u \rangle$ -tree with trivial edge stabilizers. The subgroup $\langle \mathcal{F}^{(u)} r_u \rangle$ of $\langle F R \rangle$ acts on T' and for each vertex of T' , say ${}^f r_{u'}$ with $f \in F$ and $u' \in U$, the $\langle \mathcal{F}^{(u)} r_u \rangle$ -stabilizer is $\langle \mathcal{F}^{(u)} r_u \rangle \cap {}^f (\mathbf{C}_{\mathcal{F}(u')}(r_{u'}))$. The latter intersection fixes the path in T from $1\mathcal{F}(u)$ to $f\mathcal{F}(u')$, and, hence, is trivial unless $u' = u$ and $f \in \mathcal{F}(u)$. Thus the set of vertices of T' with non-trivial $\langle \mathcal{F}^{(u)} r_u \rangle$ -stabilizer is the subset $\mathcal{F}^{(u)} r_u$ of ${}^F R (= VT')$. Since the vertex stabilizers then generate $\langle \mathcal{F}^{(u)} r_u \rangle$ we see that $\langle \mathcal{F}^{(u)} r_u \rangle \backslash T'$ is a tree. Hence, successively $\langle \mathcal{F}^{(u)} r_u \rangle$ -equivariantly contracting suitable edges of T' produces an $\langle \mathcal{F}^{(u)} r_u \rangle$ -tree with vertex set $\mathcal{F}^{(u)} r_u$ and trivial edge stabilizers, as desired. \square

5.4 Remark. If A and B are groups and r is an element of A , it follows from Theorem 5.3 that r is Cohen-Lyndon aspherical in A if and only if r is Cohen-Lyndon aspherical in $A*B$. \square

5.5 Corollary. *Suppose that A and B are groups and that A_1 is a subgroup of A and that B_1 is a subgroup of B and that r is an element of A_1*B_1 and that the natural maps embed A_1 and B_1 in $(A_1*B_1)/\langle r \rangle$.*

*If r is Cohen-Lyndon aspherical in A_1*B_1 , then r is Cohen-Lyndon aspherical in $A*B$.*

Proof. By applying Theorem 5.3 to $(A_1*B_1)_{*B_1} B (= A_1*B)$, we see that r is Cohen-Lyndon aspherical in A_1*B and also the natural map embeds $(A_1*B_1)/\langle r \rangle$ in $(A_1*B)/\langle r \rangle$. Hence, the natural map embeds A_1 in $(A_1*B)/\langle r \rangle$.

By applying Theorem 5.3 to $(A_1*B)_{*A_1} A (= A*B)$, we see that r is Cohen-Lyndon aspherical in $A*B$. \square

5.6 Corollary. *Suppose that F is locally indicable and that T is an F -tree with trivial edge stabilizers and that r is an element of F which fixes no vertex of T .*

Let H be a subgroup of F such that $H \supseteq \text{glue}(F, E\text{axis}(r))$ and $H = \langle \text{glue}(H, \text{axis}(r)) \rangle$. If r is Cohen-Lyndon aspherical in H , then r is Cohen-Lyndon aspherical in F .

Proof. Let e be an edge in $\text{axis}(r)$. Let \bar{T} denote the F -tree obtained from T by collapsing all edges in $ET - Fe$. Then $E\bar{T} = Fe$ and r shifts e and $E\text{axis}_{\bar{T}}(r) = Fe \cap E\text{axis}_T(r)$ and $\text{glue}(H, \text{axis}_T(r)) \subseteq \text{glue}(H, \text{axis}_{\bar{T}}(r))$. It follows that we can replace T with \bar{T} and assume that $ET = Fe$.

Let $T' = H(\text{axis}(r))$. By Remark 2.5, T' is an H -subtree of T .

For each $e' \in E\text{axis}(r)$, there exists some $f \in F$ such that $fe = e'$, and then $f \in \text{glue}(F, E\text{axis}(r)) \subseteq H$ and $e' = fe \in He$. Hence $E\text{axis}(r) \subseteq He$. Hence $ET' = H(E\text{axis}(r)) \subseteq He$. In summary, $ET = Fe$ and $ET' = He$.

Case 1. $|F \setminus (VT)| = 2$.

Here, we can write $F = A*B$ and no F -conjugate of r lies in $A \cup B$, and $H = A_1*B_1$ for some $A_1 \leq A$ and some $B_1 \leq B$. By Corollary 4.4 and Corollary 5.5, r is Cohen-Lyndon aspherical in $A*B$ ($= F$).

Case 2. $|F \setminus (VT)| = 1$.

Here, we can write $F = A * (b: \{1\} \rightarrow \{1\})$ and no F -conjugate of r lies in A .

Case 2.1. $|H \setminus (VT')| = 2$.

Here, $H = A_1 * {}^b A_2$ for some $A_1 \leq A$ and some $A_2 \leq A$.

Then r is Cohen-Lyndon aspherical in $A * {}^b A$, by Corollary 4.4 and Corollary 5.5.

Now r is Cohen-Lyndon aspherical in $(A * {}^b A) * (b: A \rightarrow {}^b A)$ ($= A * \langle b \mid \rangle = F$), by Corollary 4.4 and Theorem 5.3.

Case 2.2. $|H \setminus (VT')| = 1$.

Here, $H = A_1 * (ba: \{1\} \rightarrow \{1\})$ for some $A_1 \leq A$ and some $a \in A$.

If no F -conjugate of r lies in $\langle ba \mid \rangle$, then r is Cohen-Lyndon aspherical in $A * \langle ba \mid \rangle$ ($= A * \langle b \mid \rangle = F$), by Corollary 4.4 and Corollary 5.5.

Thus we may assume that r itself lies in $\langle ba \mid \rangle$ and r is then clearly Cohen-Lyndon aspherical in $\langle ba \mid \rangle$. Hence, r is Cohen-Lyndon aspherical in $A * \langle ba \mid \rangle$ ($= A * \langle b \mid \rangle = F$), by Remark 5.4.

Hence, in all cases, r is Cohen-Lyndon aspherical in F . \square

5.7 Notation. If a and b are elements of F , we shall let $\langle a^i \mid \rangle b := \langle a^i b \mid i \in \mathbb{Z} \rangle$. For any subset J of \mathbb{Z} , we let $a^J b := \langle a^j b \mid j \in J \rangle \subseteq \langle a^i \mid \rangle b$. \square

The following will be useful for simplifying calculations.

5.8 Lemma. *Suppose that f is an element of F and that N is a normal subgroup of F such that $F/N = \langle fN \mid \rangle$. Let z be a symbol, let $\tilde{F} := F * \langle z \mid \rangle$, and let \tilde{N} denote the smallest normal subgroup of \tilde{F} containing $N \cup \{f\bar{z}\}$. Then $\tilde{F}/\tilde{N} = \langle f\tilde{N} \mid \rangle = \langle z\tilde{N} \mid \rangle$ and N is a free factor of \tilde{N} .*

Proof. It is easy to see that $F = \langle f \mid \rangle \rtimes N$ and $\tilde{F}/\tilde{N} = \langle f\tilde{N} \mid \rangle = \langle z\tilde{N} \mid \rangle$. It is not difficult to use generators and relations to check that $\tilde{F} = F * \langle z \mid \rangle = \langle f \mid \rangle \rtimes (N * \langle \langle f \mid \rangle (f\bar{z}) \mid \rangle)$, and, hence, $\tilde{N} = N * \langle \langle f \mid \rangle (f\bar{z}) \mid \rangle$. \square

The following is proved but not stated in [6]; here, we give an argument with a different rewriting procedure, to provide some variety.

5.9 Lemma. (Edjvet-Howie) *Suppose that A and B are locally indicable groups, and that r is an element of $A*B$ such that no $A*B$ -conjugate of r lies in $A \cup B$.*

*Let N be a normal subgroup of $A*B$ such that $r \in N$ and $(A*B)/N$ is infinite and cyclic. If r is Cohen-Lyndon aspherical in N , then r is Cohen-Lyndon aspherical in $A*B$.*

Proof. Let $F = A*B$.

It follows from Lemma 5.8 and Remark 5.4 that, by adjoining an infinite, cyclic free factor to A if necessary, we may assume that there exists some $a \in A$ such that $F/N = \langle aN \mid \rangle$.

Similarly, by adjoining an infinite, cyclic free factor to B if necessary, we may further assume that there exists some $b \in B$ such that $F/N = \langle aN \mid \rangle = \langle bN \mid \rangle$.

Let $N_A := A \cap N$ and let $N_B := B \cap N$. Notice that ${}^a N_A = N_A$ and ${}^a N_B = {}^{a\bar{b}} N_B$. It is not difficult to use generators and relations to check that $F = \langle a \mid \ \rangle \rtimes (N_A * N_B * \langle {}^{a^l} \ \rangle \langle a\bar{b} \mid \ \rangle)$, and, hence, $N = N_A * N_B * \langle {}^{a^l} \ \rangle \langle a\bar{b} \mid \ \rangle$.

We may replace r with any F -conjugate of r , since any automorphism of N respects Cohen-Lyndon asphericity.

Let $j \in \mathbb{Z}$. Conjugation by a^j induces an automorphism of N and we obtain the free-product decomposition $N = N_A * {}^{a^j} N_B * \langle {}^{a^l} \ \rangle \langle a\bar{b} \mid \ \rangle$; here, we consider the resulting cyclically reduced expression for r , and we see that there exists some finite subset J_j in \mathbb{Z} such that this cyclically reduced expression lies in $N_A * {}^{a^j} N_B * \langle {}^{a^{J_j}} \ \rangle \langle a\bar{b} \mid \ \rangle$. We may assume that J_j is minimal and that r itself lies in this free factor of N . If J_j is empty, then, by replacing r with $\bar{a}^j r$, we may assume that $r \in N_A * N_B$, and then, by Remark 5.4, r is Cohen-Lyndon aspherical in $N_A * N_B$. By Corollary 4.4 and Corollary 5.5, r is Cohen-Lyndon aspherical in $A * B$ ($= F$).

Thus we may assume that, for each $j \in \mathbb{Z}$, J_j is non-empty.

Let $j \in \mathbb{Z}$. Then ${}^{a^{j+1}} N_B = ({}^{a^j(a\bar{b})})({}^{a^j} N_B)$, and we find that $J_j \subseteq J_{j+1} \cup \{j\}$. Similarly, $J_{j+1} \subseteq J_j \cup \{j\}$, and, hence, $J_j \cup \{j\} = J_{j+1} \cup \{j\}$.

It follows that $\{j \in \mathbb{Z} \mid \min J_j < j\} = \{j \in \mathbb{Z} \mid \min J_{j+1} = \min J_j < j\} = \{j \in \mathbb{Z} \mid \min J_{j+1} < j\}$. It is not difficult to see that this set cannot be all of \mathbb{Z} . Let $K := \{j \in \mathbb{Z} \mid \min J_j \geq j\}$. Then $K = \{j \in \mathbb{Z} \mid \min J_{j+1} \geq j\}$ and $K \neq \emptyset$. Let us choose $k \in K$ to minimize the pair $(|J_k|, (\min J_k) - k)$. If $k+1 \in K$, then $\min J_{k+1} \geq k+1$, and then $k \notin J_{k+1}$, and then $J_{k+1} \subseteq J_k$, and then, by the minimality property of k , $J_{k+1} = J_k$ and $(\min J_{k+1}) - (k+1) \geq (\min J_k) - k$, which is a contradiction. Thus $k+1 \in \mathbb{Z} - K$. It follows that $\min J_{k+1} = k$.

By replacing r with $\bar{a}^k r$, we may assume that $k = 0$, and, hence, $\min J_1 = 0$. Let $\nu := \max J_1$. Then $\nu \geq 0$ and

$$r \in N_A * {}^a N_B * \langle {}^{a^{0+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle = N_A * {}^{a\bar{b}} N_B * \langle {}^{a^{0+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle = N_A * N_B * \langle {}^{a^{0+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle.$$

Since $\min J_1 = 0$, no N -conjugate of r lies in $N_A * {}^a N_B * \langle {}^{a^{1+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle = N_A * {}^{a\bar{b}} N_B * \langle {}^{a^{1+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle$.

We claim that no N -conjugate of r lies in $N_A * N_B * \langle {}^{a^{0+(\nu-1)}} \ \rangle \langle a\bar{b} \mid \ \rangle$. The case of the claim where $\nu = 0$ holds because J_0 is non-empty. The case of the claim where $\nu \geq 1$ holds because $\max J_1 = \nu$ and, hence, no N -conjugate of r lies in

$$N_A * {}^a N_B * \langle {}^{a^{0+(\nu-1)}} \ \rangle \langle a\bar{b} \mid \ \rangle = N_A * {}^{a\bar{b}} N_B * \langle {}^{a^{0+(\nu-1)}} \ \rangle \langle a\bar{b} \mid \ \rangle = N_A * N_B * \langle {}^{a^{0+(\nu-1)}} \ \rangle \langle a\bar{b} \mid \ \rangle.$$

This proves the claim.

By Remark 5.4, r is Cohen-Lyndon aspherical in $N_A * N_B * \langle {}^{a^{0+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle$. By Corollary 4.4 and Theorem 5.3, r is Cohen-Lyndon aspherical in the HNN extension

$$(N_A * N_B * \langle {}^{a^{0+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle) * (a : (N_A * N_B * \langle {}^{a^{0+(\nu-1)}} \ \rangle \langle a\bar{b} \mid \ \rangle) \rightarrow (N_A * {}^{a\bar{b}} N_B * \langle {}^{a^{1+\nu}} \ \rangle \langle a\bar{b} \mid \ \rangle)),$$

which is $\langle a \mid \ \rangle \rtimes (N_A * N_B * \langle {}^{a^l} \ \rangle \langle a\bar{b} \mid \ \rangle)$ ($= F$). □

5.10 Corollary. *Suppose that F is locally indicable and that T is an F -tree with trivial edge stabilizers and that r is an element of F which fixes no vertex of T .*

Let N be a normal subgroup of F such that $r \in N$ and F/N is infinite and cyclic. If r is Cohen-Lyndon aspherical in N , then r is Cohen-Lyndon aspherical in F .

Proof. The argument is similar to the proof of Corollary 5.6. Let e be an edge in $\text{axis}(r)$. Let \bar{T} denote the F -tree obtained from T by contracting all edges in $ET - Fe$. Then $E\bar{T} = Fe$ and r shifts e . Hence r fixes no vertex of \bar{T} . Thus we may replace T with \bar{T} and assume that $ET = Fe$.

Case 1. $|F \setminus (VT)| = 2$.

Here, we can write $F = A * B$ and no F -conjugate of r lies in $A \cup B$. Then, r is Cohen-Lyndon aspherical in F by Lemma 5.9.

Case 2. $|F \setminus (VT)| = 1$.

Here, we can write $F = A * (b : \{1\} \rightarrow \{1\})$ and no F -conjugate of r lies in A .

If no F -conjugate of r lies in $\langle b \mid \ \rangle$, then r is Cohen-Lyndon aspherical in $A * \langle b \mid \ \rangle$ ($= F$) by Lemma 5.9.

Thus we may assume that r itself lies in $\langle b \mid \ \ \rangle$ and r is then clearly Cohen-Lyndon aspherical in $\langle b \mid \ \ \rangle$. Hence, r is Cohen-Lyndon aspherical in $A*\langle b \mid \ \ \rangle (= F)$, by Remark 5.4.

Thus, in all cases, r is Cohen-Lyndon aspherical in F . \square

5.11 The Local-indicability Cohen-Lyndon Theorem. *Suppose that F is a locally indicable group and that T is an F -tree with trivial edge stabilizers. If r is an element of F that fixes no vertex of T , then $\mathbf{C}_F(r)$ is infinite and cyclic, and r is Cohen-Lyndon aspherical in F .*

Proof. Recall from Definitions 2.3 that r has a unique root in F and that $\mathbf{C}_F(r) = \langle \sqrt[r]{r} \rangle$.

Let $S := \{ \sqrt[r]{r} \}$ and $\Phi := \emptyset$. We may then assume that we are in Setting 3.1, and that v is a vertex in $\text{axis}(\sqrt[r]{r})$. Then $\langle S \rangle Y = \text{axis}(\sqrt[r]{r}) = \text{axis}(r)$. We shall use decreasing induction to show that, for each $n \in [0 \uparrow \nu]$, r is Cohen-Lyndon aspherical in F_n .

By Corollary 4.2, $F_\nu S$ is a Whitehead subset of F_ν . Since $|S| = 1$, F_ν is cyclic, and, hence, $F_\nu S = S$. Thus, $F_\nu = \langle \sqrt[r]{r} \mid \ \ \rangle$. It is then clear that r is Cohen-Lyndon aspherical in F_ν .

Now suppose that $n \in [0 \uparrow (\nu-1)]$ and that r is Cohen-Lyndon aspherical in F_{n+1} .

First, we wish to show that r is Cohen-Lyndon aspherical in F_n^\dagger . This is trivial if $F_n^\dagger = F_{n+1}$, and we consider only the case where F_n^\dagger/F_{n+1} is infinite and cyclic. Here, r is Cohen-Lyndon aspherical in F_n^\dagger by Corollary 5.10.

We next want to show that r is Cohen-Lyndon aspherical in F_n .

Recall that F_n^\dagger is generated by $\text{glue}(F_n^\dagger, Y)$ and that $\text{glue}(F_n, EY) \subseteq F_n^\dagger$. Now

$$\text{glue}(F_n^\dagger, Y) \subseteq \text{glue}(F_n^\dagger, \text{axis}(r))$$

and

$$\text{glue}(F_n, E\text{axis}(r)) = \text{glue}(F_n, \langle S \rangle (EY)) \subseteq \langle S \rangle \text{glue}(F_n, EY) \langle S \rangle \subseteq F_n^\dagger.$$

Here, r is Cohen-Lyndon aspherical in F_n by Corollary 5.6.

By descending induction, r is Cohen-Lyndon aspherical in $F_0 (= F)$. \square

6 Applications

We can combine Theorem 5.3, Theorem 5.11 and Corollary 4.4 to obtain a sufficient condition for a subset of F to be Cohen-Lyndon aspherical in F .

6.1 Theorem. *Let (\mathcal{F}, Y) be a graph of groups with family of groups $(\mathcal{F}(y) \mid y \in Y)$ and family of edge maps $(\bar{t}_e: \mathcal{F}(e) \rightarrow \mathcal{F}(\tau_Y e) \mid e \in EY)$, let Y_0 be a maximal subtree of Y , and let F be the fundamental group of (\mathcal{F}, Y, Y_0) .*

Let U be a subset of VY and let $(r_v \mid v \in U)$ be a family of elements of F with the property that, for each $v \in U$, the following hold.

- (a). $r_v \in \mathcal{F}(v)$ and $\mathcal{F}(v)$ is locally indicable and $\mathcal{F}(v)$ acts on some tree with trivial edge stabilizers and no vertex fixed by r_v .
- (b). For each $e \in \iota_Y^{-1}(\{v\})$, $\mathcal{F}(e)$ is a free factor of $\mathcal{F}(v)$ that contains no $\mathcal{F}(v)$ -conjugate of r_v .
- (c). For each $e \in \tau_Y^{-1}(\{v\})$, $\bar{t}_e \mathcal{F}(e)$ is a free factor of $\mathcal{F}(v)$ that contains no $\mathcal{F}(v)$ -conjugate of r_v .

Then $(r_v \mid v \in U)$ is Cohen-Lyndon aspherical in F . \square

We shall consider only the case that corresponds to the strongly staggered conditions.

6.2 Corollary. *Suppose that F is a locally indicable group and that R is a subset of F . If there exists some F -tree T with trivial edge stabilizers such that R is strongly T -staggerable modulo F , then R is Cohen-Lyndon aspherical in F .*

Cohen and Lyndon [4, Corollary 2.2] proved the case of this result where F is free and $|F \setminus VT| = 1$.

Proof of Corollary 6.2. Applied to F acting on T , the Bass-Serre Structure Theorem presents F as the fundamental group of a certain graph of groups (\mathcal{F}, Y, Y_0) . Let $<$ be an ordering of ET such that R is strongly $(T, <)$ -staggered modulo F . Here $EY = F \backslash ET$ and, by the staggered conditions, the ordering $<$ of ET induces an ordering, again denoted by $<$, of EY . By the strongly staggered conditions, $(EY, <)$ is order isomorphic to an interval in \mathbb{Z} . Let us consider only the case where $(EY, <)$ is order isomorphic to \mathbb{Z} ; the case where $(EY, <)$ is order isomorphic to a proper interval in \mathbb{Z} is handled in a similar way. By choosing an order isomorphism we get an indexing $e_{\mathbb{Z}}$ of the elements of EY .

For any interval I in \mathbb{Z} , e_I is an interval in $e_{\mathbb{Z}}$, and we let Y_I denote the subgraph of Y with edge set e_I together with the vertices of Y that are incident to these edges. By the strongly staggered conditions, Y_I is connected and $Y_0 \cap Y_I$ is the unique maximal subtree of Y_I . Then the graph of groups obtained by restricting \mathcal{F} to Y_I has a well-defined fundamental group which we denote by F_I . It can be seen that F_I is a free factor of F .

By the staggered conditions, the ordering $<$ of EY induces an ordering, again denoted by $<$, of R . Let us consider only the case where R is order isomorphic to \mathbb{Z} ; the case where R is order isomorphic to a proper interval in \mathbb{Z} is handled in a similar way. By choosing an order isomorphism we get an indexing $r_{\mathbb{Z}}$ of the elements of R . For each $n \in \mathbb{Z}$, we set $\mu_n := \min(F \backslash (F(E\text{axis}(r_n))))$ and $\nu_n := \max(F \backslash (F(E\text{axis}(r_n))))$. Clearly $\mu_n \leq \nu_n$. The staggered conditions imply that $\mu_n < \mu_{n+1}$ and $\nu_n < \nu_{n+1}$.

We can then express F as the fundamental group of a second graph of groups whose family of vertex groups is $(F_{[\mu_n \uparrow \nu_n]} \mid n \in \mathbb{Z})$ and whose family of edge groups is $(F_{[\mu_n \uparrow \nu_n]} \cap F_{[\mu_{n+1} \uparrow \nu_{n+1}]} \mid n \in \mathbb{Z})$, with the edge maps being those suggested by the intersection notation. Here the underlying graph has the form of the real line.

For each $n \in \mathbb{Z}$, some F -conjugate of r_n lies in $F_{[\mu_n \uparrow \nu_n]}$ and we may assume that r_n itself lies in the vertex group $F_{[\mu_n \uparrow \nu_n]}$. By the definition of μ_n and ν_n and the staggered conditions, no F -conjugate of r_n lies in either of the two incident-edge groups.

By Theorem 6.1, R is Cohen-Lyndon aspherical in F . □

Cohen-Lyndon asphericity has many applications, and we conclude by mentioning two of them.

6.3 Setting. Suppose that F is locally indicable, and that T is an F -tree with trivial edge stabilizers, and that R_0 is a subset of F which is strongly T -staggerable modulo F .

By Definitions 2.3, R_0 has unique roots in F and $\sqrt[F]{R_0}$ is strongly T -staggerable modulo F . For each $r \in R_0$, we have $\mathbf{C}_F(r) = \langle \sqrt[F]{r} \rangle$.

By Corollary 6.2, R_0 and $\sqrt[F]{R_0}$ are Cohen-Lyndon aspherical in F .

Let $G := F / \langle R_0 \rangle$. For each $r \in R_0$, let $G_r := \mathbf{C}_F(r) / \langle r \rangle = \langle \sqrt[F]{r} \mid r \rangle$, a finite, cyclic subgroup of G .

Let $(Y_r \mid r \in R_0)$ be a family of subsets of F with the properties that, for each $r \in R_0$, Y_r is a transversal for the $\langle \sqrt[F]{R_0} \rangle$ -action on F by multiplication on the left (or right) and $\bigcup_{r \in R_0} (Y_r \langle \sqrt[F]{r} \rangle)$ is a free-generating set of $\langle \sqrt[F]{R_0} \rangle$.

Let $(X_r \mid r \in R_0)$ be a family of subsets of F with the properties that, for each $r \in R_0$, X_r is a transversal for the $(\langle \sqrt[F]{R_0} \rangle \langle \sqrt[F]{r} \rangle)$ -action on F by multiplication on the right and $\bigcup_{r \in R_0} (X_r r)$ is a free-generating set of $\langle \sqrt[F]{R_0} \rangle$. For each element v in the G -graph $\langle \sqrt[F]{R_0} \rangle \backslash T$, let G_v denote the G -stabilizer of v . Also, let E_0 , resp. V_0 , be a transversal for the natural G -action on $\langle \sqrt[F]{R_0} \rangle \backslash ET$, resp. $\langle \sqrt[F]{R_0} \rangle \backslash VT$. Here, F is the free product of a free group and the group $\ast_{v \in V_0} G_v$. □

We first discuss torsion.

6.4 Corollary. *In Setting 6.3, let N denote the (normal) subgroup of G generated by the elements of G of finite order. Then the following hold.*

- (i). *For each $r \in R_0$, Y_r may be viewed as a transversal for the N -action on G by multiplication on the left, and $N = \ast_{r \in R_0} \ast_{g \in Y_r} {}^g G_r$, and, hence, N is a free product of finite, cyclic groups.*
- (ii). *G/N is locally indicable, and every torsion-free subgroup of G is locally indicable.*
- (iii). *Each non-trivial, torsion subgroup of G lies in exactly one of the (finite, cyclic) subgroups of G of the form ${}^g G_r$, $r \in R_0$, $g \in G$, and then r is unique and the coset $gG_r \in G/G_r$ is unique.*

In the case where F acts freely on T , and $|R| = 1$, (iii) can be attributed to Magnus and Lyndon [13], (i) is due to Fischer, Karrass and Solitar [7], and (ii) can be attributed to Brodskiĭ [2].

In the case where $F \setminus T$ has one edge and two vertices, and $|R| = 1$, these results can be attributed to Edjvet and Howie.

Proof of Corollary 6.4. Let $\overline{G} := F / \langle \sqrt[r]{R_0} \rangle$.

By Theorem 4.3(ii), \overline{G} is locally indicable, and, in particular, \overline{G} is torsion free.

Let $N_1 := \langle \sqrt[r]{R_0} \rangle / \langle R_0 \rangle$. Then $N_1 \trianglelefteq G$ and $G/N_1 = \overline{G}$. Since \overline{G} is torsion free, every element of G of finite order lies in N_1 , that is, $N \leq N_1$. Also, for each $r \in R_0$, the (faithful) image of Y_r in G is a transversal for the N_1 -action on G by multiplication on the left.

Since $\bigcup_{r \in R_0} (\langle \sqrt[r]{R_0} \rangle^{Y_r}) = \bigcup_{r \in R_0} \langle R_0 \rangle = R_0$, it follows that

$$N_1 = \langle \sqrt[r]{R_0} \rangle / \langle R_0 \rangle = \langle \bigvee_{r \in R_0} (Y_r \langle \sqrt[r]{R_0} \rangle) \mid \bigvee_{r \in R_0} \langle R_0 \rangle \rangle = \bigstar_{r \in R_0} \bigstar_{y \in Y_r} \langle \sqrt[r]{R_0} \mid r \rangle = \bigstar_{r \in R_0} \bigstar_{y \in Y_r} {}^y G_r.$$

Thus, N_1 is a free product of finite, cyclic groups. In particular, N_1 is generated by some set of elements of finite order in G , that is, $N_1 \leq N$. Hence $N_1 = N$ and (i) holds. Also, every torsion-free subgroup of N is free. Hence (ii) holds.

Any non-trivial, torsion subgroup H of G lies in N , and, by well-known properties of free products, there exists some $n \in N$ such that $H \leq {}^{ny}G_r$ for a unique $r \in R_0$ and a unique $y \in Y_r$, and here the coset $n {}^y G_r \in N / {}^y G_r$ is unique. Since the (faithful) image of Y_r in G is a transversal for the N -action on G by multiplication on the left, we see that (iii) holds. \square

We now consider exact sequences and homology groups.

6.5 Corollary. *In Setting 6.3, the following hold.*

- (i). *The left F -action on $\langle R_0 \rangle$ by conjugation induces a left G -action on $\langle R_0 \rangle^{ab}$, and $\langle R_0 \rangle^{ab}$ is then a left $\mathbb{Z}G$ -module that is naturally isomorphic to $\bigoplus_{r \in R_0} \mathbb{Z}[G/G_r]$.*
- (ii). *There exist natural exact sequences of left $\mathbb{Z}G$ -modules that have the form*

$$0 \rightarrow \bigoplus_{r \in R_0} \mathbb{Z}[G/G_r] \rightarrow \bigoplus_{e \in E_0} \mathbb{Z}[G] \rightarrow \bigoplus_{v \in V_0} \mathbb{Z}[G/G_v] \rightarrow \mathbb{Z} \rightarrow 0, \quad (8)$$

$$0 \rightarrow \bigoplus_{r \in R_0} \mathbb{Z}[G] \rightarrow \bigoplus_{e \in R_0 \vee E_0} \mathbb{Z}[G] \rightarrow \bigoplus_{v \in R_0 \vee V_0} \mathbb{Z}[G/G_v] \rightarrow \mathbb{Z} \rightarrow 0. \quad (9)$$

- (iii). *For each $n \in [3 \uparrow \infty[$, the change-of-groups natural transformation $\bigoplus_{v \in R_0 \vee V_0} \mathbf{H}_n(G_v, -) \rightarrow \mathbf{H}_n(G, -)$, between functors from the category of right $\mathbb{Z}G$ -modules to the category of abelian groups, is an isomorphism of functors, and the change-of-groups natural transformation $\mathbf{H}^n(G, -) \rightarrow \prod_{v \in R_0 \vee V_0} \mathbf{H}^n(G_v, -)$, between functors from the category of left $\mathbb{Z}G$ -modules to the category of abelian groups, is an isomorphism of functors.*

In the case where F is free and $|F \setminus VT| = 1$, (i) is Lyndon's Identity Theorem [13, Section 7]; in the case where F acts freely and $|R_0| = 1$, (i) is Lyndon's Simple Identity Theorem [13, Section 7]. In the case where $|R_0| = 1$, and $F \setminus T$ has one edge and two vertices, (i) is Howie's Simple Identity Theorem [10, Theorem 11]. The results (ii), (iii) are straightforward consequences of (i); see [13, Theorem 11.1] and [10, Theorem 3].

Proof of Corollary 6.5. For each $r \in R_0$, we have bijective correspondences

$$X_r \simeq F / (\langle R_0 \rangle \langle \sqrt[r]{R_0} \rangle) \simeq G / \langle \sqrt[r]{R_0} \langle R_0 \rangle \rangle \simeq G/G_r.$$

Since $\langle R_0 \rangle = \langle \bigvee_{r \in R_0} \langle X_r \rangle \mid \rangle$, we then have isomorphisms of abelian groups

$$\langle R_0 \rangle^{ab} \simeq \mathbb{Z}[\bigvee_{r \in R_0} \langle X_r \rangle] \simeq \mathbb{Z}[\bigvee_{r \in R_0} \langle X_r \rangle] \simeq \mathbb{Z}[\bigvee_{r \in R_0} \langle G/G_r \rangle] \simeq \bigoplus_{r \in R_0} \mathbb{Z}[G/G_r].$$

The composite isomorphism of abelian groups $\langle R_0 \rangle^{ab} \simeq \bigoplus_{r \in R_0} \mathbb{Z}[G/G_r]$ is compatible with the G -actions, and we find that (i) holds.

By Theorem 4.3(i), $\langle {}^F R_0 \rangle$ acts freely on T . By [5, Definitions I.8.1], the fundamental group of $\langle {}^F R_0 \rangle \backslash T$ can be identified with $\langle {}^F R_0 \rangle$. Then (i) and [5, Theorem I.9.2] give a sequence as in (8). We leave it as an exercise to construct a sequence as in (9) which maps onto the above-constructed sequence in (8) with kernel a short exact sequence. Alternatively, one can arrange for the sequences in (8) and (9) to be the augmented cellular chain complexes of acyclic, simply connected, hence contractible, CW-complexes on which G acts by permuting the cells. Here, the second CW-complex is obtained from the first CW-complex by G -equivariantly drawing on each two-cell a point and a finite set of edges joining the new point to old points. This second CW-complex has the property that every non-trivial finite subgroup of G fixes exactly one point of the space and the fixed point is a zero-cell.

For the short exact sequence

$$0 \rightarrow \ker \epsilon \rightarrow \bigoplus_{v \in R_0 \vee V_0} \mathbb{Z}[G/G_v] \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0$$

of left $\mathbb{Z}G$ -modules where ϵ is the corresponding augmentation map, we see from (9) that $\ker \epsilon$ has a free $\mathbb{Z}G$ -resolution of length at most two. Resulting long exact sequences in homology then show that (iii) holds. \square

Acknowledgments

The research of the first- and second-named authors was jointly funded by the MCI (Spain) through project MTM2008-01550.

We are grateful to Ian Chiswell and Jim Howie for several helpful observations.

References

- [1] Yago Antolín, Warren Dicks and Peter A. Linnell, *Non-orientable surface-plus-one-relation groups*, J. Algebra, Karl Gruenberg Memorial Issue, to appear.
- [2] S. D. Brodskii, *Equations over groups, and groups with one defining relation*, Siberian Math. J. **25** (1984), 235–251.
- [3] Ian M. Chiswell, Donald J. Collins and Johannes Huebschmann, *Aspherical group presentations*, Math. Z. **178** (1981), 1–36.
- [4] D. E. Cohen and R. C. Lyndon, *Free bases for normal subgroups of free groups*, Trans. Amer. Math. Soc. **108** (1963), 526–537.
- [5] Warren Dicks and M. J. Dunwoody, *Groups acting on graphs*, Cambridge Stud. Adv. Math. **17**, CUP, Cambridge, 1989, xvi+283 pp.
Errata at: <http://mat.uab.cat/~dicks/DDerr.html>
- [6] Martin Edjvet and James Howie, *A Cohen-Lyndon theorem for free products of locally indicable groups*, J. Pure Appl. Algebra **45** (1987), 41–44.
- [7] J. Fischer, A. Karrass and D. Solitar, *On one-relator groups having elements of finite order*, Proc. Amer. Math. Soc. **33** (1972), 297–301.
- [8] James Howie, *On pairs of 2-complexes of equations over groups*, J. Reine Angew. Math. **324** (1981), 165–174.
- [9] James Howie, *On locally indicable groups*, Math. Z. **180** (1982), 445–461.
- [10] James Howie, *Cohomology of one-relator products of locally indicable groups*, J. London Math. Soc. **30** (1984), 419–430.
- [11] James Howie, *A short proof of a theorem of Brodskii*, Publ. Mat. **44** (2000), 641–647.
- [12] A. Karrass and D. Solitar, *On a theorem of Cohen and Lyndon about free bases for normal subgroups*, Canadian J. Math. **24** (1972), 1086–1091.
- [13] Roger C. Lyndon, *Cohomology theory of groups with a single defining relation*, Ann. of Math. **52** (1950), 650–665.
- [14] Roger C. Lyndon and Paul E. Schupp, *Combinatorial group theory*, Ergeb. Math. Grenzgeb. **89**, Springer-Verlag, Berlin, 1977, ix+339 pp.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, E-08193 BELLATERRA (BARCELONA), SPAIN
E-mail address: yagoap@mat.uab.cat

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, E-08193 BELLATERRA (BARCELONA), SPAIN
E-mail address: dicks@mat.uab.cat
URL: <http://mat.uab.cat/~dicks/>

DEPARTMENT OF MATHEMATICS, VIRGINIA TECH, BLACKSBURG, VA 24061-0123, USA
E-mail address: linnell@math.vt.edu
URL: <http://www.math.vt.edu/people/plinnell/>